



FS

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : C07D 239/42, A61K 31/505, A61P 7/00, C07D 401/12, A61K 31/495, C07D 405/12, 409/12, 285/10, A61K 31/41, C07D 417/12</p>	A1	<p>(11) International Publication Number: <b>WO 00/43372</b></p> <p>(43) International Publication Date: 27 July 2000 (27.07.00)</p>
<p>(21) International Application Number: PCT/US00/01686</p> <p>(22) International Filing Date: 21 January 2000 (21.01.00)</p> <p>(30) Priority Data: 60/116,923 22 January 1999 (22.01.99) US 60/160,999 21 October 1999 (21.10.99) US</p> <p>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 60/116,923 (CIP) Filed on 22 January 1999 (22.01.99)</p> <p>(71) Applicants (for all designated States except US): ELAN PHARMACEUTICALS, INC. [US/US]; 800 Gateway Boulevard, South San Francisco, CA 94080 (US). AMERICAN HOME PRODUCTS [US/US]; 5 Giraldi Farms, Madison, NJ 07940 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): KONRADI, Andrei [US/US]; 95 Cervantes #105, San Francisco, CA 94123 (US). PLEISS, Michael, A. [US/US]; 848 Stella Court,</p>		<p>Sunnyvale, CA 94087-1355 (US). THORSETT, Eugene, D. [US/US]; 571 Buena Vista Drive, Moss Beach, CA 94038 (US). ASHWELL, Susan [GB/US]; 1015 Aspen Drive, Plainsboro, NJ 08536 (US). WELMAKER, Gregory, S. [US/US]; 2 Jason Court, Jackson, NJ 08527 (US). KREFT, Anthony [US/US]; 43 Barley Court, Langhorne, PA 19047 (US). SARANTAKIS, Dimitrios [US/US]; 262 Sentinel Avenue, Newtown, PA 18940 (US). DRESSEN, Darren, B. [US/US]; 3110 Casa De Campo #2, San Mateo, CA 94403 (US). GRANT, Francine, S. [FR/US]; 66 Marilyn Drive, Milpitas, CA 95035 (US). SEMKO, Christopher [CA/US]; 2361 Carpenter Court, Fremont, CA 94539 (US). XU, Ying-Zi [CN/US]; 5136 Amberwood Drive, Fremont, CA 94555 (US).</p> <p>(74) Agents: HAGENAH, Jeffrey, A. et al.; Burns, Doane Swecker &amp; Mathis, LLP, P.O. Box 1404, Alexandria, VA 22313-1404 (US).</p> <p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
<p>(54) Title: ACYL DERIVATIVES WHICH TREAT VLA-4 RELATED DISORDERS</p> <p>(57) Abstract</p> <p>Disclosed are compounds such as Ia and Ib which bind VLA-4. Certain of these compounds also inhibit leukocyte adhesion and, in particular, leukocyte adhesion mediated by VLA-4. Such compounds are useful in the treatment of inflammatory diseases in a mammalian patient, e.g., human, such as asthma, Alzheimer's disease, atherosclerosis, AIDS dementia, diabetes, inflammatory bowel disease, rheumatoid arthritis, tissue transplantation, tumor metastasis and myocardial ischemia. The compounds can also be administered for the treatment of inflammatory brain diseases such as multiples sclerosis.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## ACYL DERIVATIVES WHICH TREAT VLA-4 RELATED DISORDERS

## CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Serial No. 60/116,923, filed January 22, 1999, and U.S. Serial No. 60/160,999, filed October 21, 1999; the disclosures of which are incorporated herein by reference in their entirety.

## BACKGROUND OF THE INVENTION

Field of the Invention

This invention relates to compounds which inhibit leukocyte adhesion and, in particular, leukocyte adhesion mediated by VLA-4.

References

The following publications, patents and patent applications are cited in this application as superscript numbers:

- <sup>1</sup> Hemler and Takada, *European Patent Application Publication No. 330,506*, published August 30, 1989
- <sup>2</sup> Elices, et al., *Cell*, 60:577-584 (1990)
- <sup>3</sup> Springer, *Nature*, 346:425-434 (1990)
- <sup>4</sup> Osborn, *Cell*, 62:3-6 (1990)
- <sup>5</sup> Vedder, et al., *Surgery*, 106:509 (1989)
- <sup>6</sup> Pretolani, et al., *J. Exp. Med.*, 180:795 (1994)
- <sup>7</sup> Abraham, et al., *J. Clin. Invest.*, 93:776 (1994)

-- 2 --

- 8 Mulligan, et al., *J. Immunology*, 150:2407 (1993)
- 9 Cybulsky, et al., *Science*, 251:788 (1991)
- 5 10 Li, et al., *Arterioscler. Thromb.*, 13:197 (1993)
- 11 Sasseville, et al., *Am. J. Path.*, 144:27 (1994)
- 10 12 Yang, et al., *Proc. Nat. Acad. Science (USA)*, 90:10494 (1993)
- 13 Burkly, et al., *Diabetes*, 43:529 (1994)
- 14 Baron, et al., *J. Clin. Invest.*, 93:1700 (1994)
- 15 15 Hamann, et al., *J. Immunology*, 152:3238 (1994)
- 16 Yednock, et al., *Nature*, 356:63 (1992)
- 20 17 Baron, et al., *J. Exp. Med.*, 177:57 (1993)
- 18 van Dinther-Janssen, et al., *J. Immunology*, 147:4207 (1991)
- 19 25 van Dinther-Janssen, et al., *Annals. Rheumatic Dis.*, 52:672 (1993)
- 20 20 Elices, et al., *J. Clin. Invest.*, 93:405 (1994)
- 21 30 Postigo, et al., *J. Clin. Invest.*, 89:1445 (1991)
- 22 Paul, et al., *Transpl. Proceed.*, 25:813 (1993)
- 23 Okarhara, et al., *Can. Res.*, 54:3233 (1994)
- 35 24 Paavonen, et al., *Int. J. Can.*, 58:298 (1994)
- 25 25 Schadendorf, et al., *J. Path.*, 170:429 (1993)
- 26 40 Bao, et al., *Diff.*, 52:239 (1993)
- 27 Lauri, et al., *British J. Cancer*, 68:862 (1993)
- 28 Kawaguchi, et al., *Japanese J. Cancer Res.*, 83:1304 (1992)



29 Kogan, et al., *U.S. Patent No. 5,510,332*, issued April 23, 1996

30 International Patent Appl. Publication No. WO 96/01644

5

All of the above publications, patents and patent applications are herein incorporated by reference in their entirety to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety.

10

#### State of the Art

VLA-4 (also referred to as  $\alpha_4\beta_1$  integrin and CD49d/CD29), first identified by Hemler and Takada<sup>1</sup> is a member of the  $\beta_1$  integrin family of cell surface receptors, each of which comprises two subunits, an  $\alpha$  chain and a  $\beta$  chain. VLA-4 contains an  $\alpha_4$  chain and a  $\beta_1$  chain. There are at least nine  $\beta_1$  integrins, all sharing the same  $\beta_1$  chain and each having a distinct  $\alpha$  chain. These nine receptors all bind a different complement of the various cell matrix molecules, such as fibronectin, laminin, and collagen. VLA-4, for example, binds to fibronectin. VLA-4 also binds non-matrix molecules that are expressed by endothelial and other cells. These non-matrix molecules include VCAM-1, which is expressed on cytokine-activated human umbilical vein endothelial cells in culture. Distinct epitopes of VLA-4 are responsible for the fibronectin and VCAM-1 binding activities and each activity has been shown to be inhibited independently.<sup>2</sup>

25

Intercellular adhesion mediated by VLA-4 and other cell surface receptors is associated with a number of inflammatory responses. At the site of an injury or other inflammatory stimulus, activated vascular endothelial cells express molecules that are adhesive for leukocytes. The mechanics of leukocyte adhesion to endothelial cells involves, in part, the recognition and binding of cell surface receptors on leukocytes to the corresponding cell surface molecules on endothelial cells. Once bound, the leukocytes migrate

30

-- 4 --

across the blood vessel wall to enter the injured site and release chemical mediators to combat infection. For reviews of adhesion receptors of the immune system, *see*, for example, Springer<sup>3</sup> and Osborn<sup>4</sup>.

5           Inflammatory brain disorders, such as experimental autoimmune encephalomyelitis (EAE), multiple sclerosis (MS) and meningitis, are examples of central nervous system disorders in which the endothelium/leukocyte adhesion mechanism results in destruction to otherwise healthy brain tissue. Large numbers of leukocytes migrate across  
10       the blood brain barrier (BBB) in subjects with these inflammatory diseases. The leukocytes release toxic mediators that cause extensive tissue damage resulting in impaired nerve conduction and paralysis.

          In other organ systems, tissue damage also occurs via an adhesion  
15       mechanism resulting in migration or activation of leukocytes. For example, it has been shown that the initial insult following myocardial ischemia to heart tissue can be further complicated by leukocyte entry to the injured tissue causing still further insult (Vedder et al.<sup>5</sup>). Other inflammatory or medical conditions mediated by an adhesion mechanism include, by way of  
20       example, asthma<sup>6-8</sup>, Alzheimer's disease, atherosclerosis<sup>9-10</sup>, AIDS dementia<sup>11</sup>, diabetes<sup>12-14</sup> (including acute juvenile onset diabetes), inflammatory bowel disease<sup>15</sup> (including ulcerative colitis and Crohn's disease), multiple sclerosis<sup>16-17</sup>, rheumatoid arthritis<sup>18-21</sup>, tissue  
25       transplantation<sup>22</sup>, tumor metastasis<sup>23-28</sup>, meningitis, encephalitis, stroke, and other cerebral traumas, nephritis, retinitis, atopic dermatitis, psoriasis, myocardial ischemia and acute leukocyte-mediated lung injury such as that which occurs in adult respiratory distress syndrome.

          In view of the above, assays for determining the VLA-4 level in a  
30       biological sample containing VLA-4 would be useful, for example, to


-- 5 --

diagnosis VLA-4 mediated conditions. Additionally, despite these advances in the understanding of leukocyte adhesion, the art has only recently addressed the use of inhibitors of adhesion in the treatment of inflammatory brain diseases and other inflammatory conditions<sup>29,30</sup>. The present invention addresses these and other needs.

### SUMMARY OF THE INVENTION

This invention provides compounds which bind to VLA-4. Such compounds can be used, for example, to assay for the presence of VLA-4 in a sample and in pharmaceutical compositions to inhibit cellular adhesion mediated by VLA-4, for example, binding of VCAM-1 to VLA-4. The compounds of this invention have a binding affinity to VLA-4 as expressed by an  $IC_{50}$  of about 15  $\mu$ M or less (as measured using the procedures described in Example A below).

Accordingly, in one of its method aspects, this invention is directed to a method for treating a disease mediated by VLA-4 in a patient, which method comprises administering a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of formula Ia and/or Ib:



wherein, in formula Ia,  $R^1$  and  $R^2$ , together with the carbon atom and W to which they are bound respectively, are joined to form an aryl,

cycloalkenyl, heteroaryl or heterocyclic group having at least five atoms in the aryl, cycloalkenyl, heteroaryl or heterocyclic group and optionally containing or additionally containing in the case of heteroaryl and heterocyclic groups 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heteroaryl or heterocyclic group is mono-cyclic;

in formula Ib,  $R^1$  and  $R^2$ , together with the carbon atom and  $W'$  to which they are bound respectively, are joined to form a cycloalkyl, cycloalkenyl or heterocyclic group having at least five atoms in the cycloalkyl, cycloalkenyl or heterocyclic group and optionally containing or additionally containing in the case of the heterocyclic group 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heterocyclic group is mono-cyclic;

and further wherein said aryl, cycloalkyl, cycloalkenyl, heteroaryl or heterocyclic group of formula Ia or Ib is optionally substituted, on any ring atom capable of substitution, with 1-3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen, hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino,  $-OS(O)_2$ -alkyl,  $-OS(O)_2$ -substituted alkyl,  $-OS(O)_2$ -aryl,  $-OS(O)_2$ -substituted aryl,  $-OS(O)_2$ -heteroaryl,

-- 7 --

-OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted  
 heterocyclic, -OSO<sub>2</sub>-NRR where each R is independently hydrogen or alkyl,  
 -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-  
 substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl,  
 5 -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-  
 alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-  
 substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted  
 heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted  
 heterocyclic where R is hydrogen or alkyl, -N[S(O)<sub>2</sub>-R']<sub>2</sub> and -N[S(O)<sub>2</sub>-  
 10 NR']<sub>2</sub> where each R' is independently selected from the group consisting of  
 alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted  
 heteroaryl, heterocyclic and substituted heterocyclic;

R<sup>3</sup> and R<sup>3'</sup> are independently selected from the group consisting of  
 hydrogen, isopropyl, -CH<sub>2</sub>Z where Z is selected from the group consisting  
 15 of hydrogen, hydroxyl, acylamino, alkyl, alkoxy, aryloxy, aryl, aryloxyaryl,  
 carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl,  
 carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl,  
 carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic,  
 carboxyl-substituted heterocyclic, cycloalkyl, substituted alkyl, substituted  
 20 alkoxy, substituted aryl, substituted aryloxy, substituted aryloxyaryl,  
 substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and  
 substituted heterocyclic, and

where R<sup>3</sup> and R<sup>3'</sup> are joined to form a substituent selected from the  
 group consisting of =CHZ where Z is defined above provided that Z is not  
 25 hydroxyl or thiol, cycloalkyl, substituted cycloalkyl, cycloalkenyl,  
 substituted cycloalkenyl, heterocyclic and substituted heterocyclic;

Q is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)<sub>2</sub>,  
 and -NR<sup>4</sup>-;

R<sup>4</sup> is selected from the group consisting of hydrogen, alkyl,  
 30 substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, substituted

-- 8 --

cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic or, optionally, R<sup>4</sup> and R<sup>1</sup> or R<sup>4</sup> and R<sup>2</sup>, together with the atoms to which they are bound, are joined to form a heteroaryl, a substituted heteroaryl, a heterocyclic or a substituted heterocyclic group;

W is selected from the group consisting of nitrogen and carbon; and

W' is selected from the group consisting of nitrogen, carbon, oxygen, sulfur, S(O), and S(O)<sub>2</sub>;

X is selected from the group consisting of hydroxyl, alkoxy, substituted alkoxy, alkenoxy, substituted alkenoxy, cycloalkoxy, substituted cycloalkoxy, cycloalkenoxo, substituted cycloalkenoxo, aryloxy, substituted aryloxy, heteroaryloxy, substituted heteroaryloxy, heterocycloxy, substituted heterocycloxy and -NR''R'' where each R'' is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;


and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;

and further wherein the compound of formula Ia and/or Ib has a binding affinity to VLA-4 as expressed by an IC<sub>50</sub> of about 15 μM or less.

Preferably, in the above method, R<sup>3</sup> is -(CH<sub>2</sub>)<sub>x</sub>-Ar-R<sup>9</sup>, where Ar is aryl, substituted aryl, heteroaryl and substituted heteroaryl; R<sup>9</sup> is selected from the group consisting of acyl, acylamino, acyloxy, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, oxythiocarbonylamino, thioamidino, thiocarbonylamino, aminosulfonylamino, aminosulfonyloxy, aminosulfonyl, oxysulfonylamino and oxysulfonyl; and x is an integer from 0 to 4. R<sup>3'</sup> is preferably alkyl or hydrogen; more preferably, R<sup>3'</sup> is hydrogen.

- 9 -

More preferably,  $R^3$  is a group of the formula:




wherein  $R^9$  and  $x$  are as defined herein. Preferably,  $R^9$  is in the *para* position of the phenyl ring; and  $x$  is an integer of from 1 to 4, more preferably,  $x$  is 1.


In a preferred embodiment,  $R^9$  is selected from  $-O-Z-NR^{11}R^{11'}$  and  $-O-Z-R^{12}$  wherein  $R^{11}$  and  $R^{11'}$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, and where  $R^{11}$  and  $R^{11'}$  are joined to form a heterocycle or a substituted heterocycle,  $R^{12}$  is selected from the group consisting of heterocycle and substituted heterocycle, and  $Z$  is selected from the group consisting of  $-C(O)-$  and  $-SO_2-$ . More preferably,  $R^9$  is  $-OC(O)NR^{11}R^{11'}$ , wherein  $R^{11}$  and  $R^{11'}$  are as defined herein.

In the above method,  $Z$  is preferably  $-C(O)-$ . Preferably,  $Q$  is  $-NR^4-$ .

In a preferred embodiment, the above method employs a compound of formula IIa or IIb:



-- 10 --



wherein  $R^3$ ,  $R^{3'}$  and X are as defined herein;

ring A and ring B independently form a heteroaryl or substituted heteroaryl group having two nitrogen atoms in the heteroaryl ring;

$R^5$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

$R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $-SO_2R^{10}$  where  $R^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl;

or optionally, one of,  $R^4$  and ring A,  $R^4$  and  $R^5$ ,  $R^4$  and  $R^6$ , or  $R^5$  and  $R^6$ , together with the atoms to which they are bound, can be joined to form a heterocyclic or substituted heterocyclic ring;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof; and provided that ring B does not form a 6-amino or substituted amino pyrimidin-4-yl group.

Preferably, ring A forms a pyridazine, pyrimidine or pyrazine ring; more preferably, a pyrimidine or pyrazine ring; wherein the pyridazine,




-- 11 --

pyrimidine or pyrazine ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.

Preferably, ring B forms a pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or a 1,1-dioxo-1,2,5-thiadiazole ring; more preferably, a pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or a 1,1-dioxo-1,2,5-thiadiazole ring; wherein the pyridazine, pyrimidine or pyrazine ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.


In another preferred embodiment, the method employs a compound of formula IIIa, IIIc, IIId, IIIe or IIIf:

20



IIIa

25




IIIc

30


- 12 -

5




IIIId

10



IIIe

15



IIIf

20

25

wherein  $R^3$ ,  $R^{3'}$  and  $X$  are as defined herein;

$R^{4'}$  is selected from the group consisting of hydrogen and alkyl or, optionally, one of,  $R^{4'}$  and  $R^5$ ,  $R^{4'}$  and  $R^6$ ,  $R^5$  and  $R^6$ ,  $R^5$  and  $R^8$ , or  $R^6$  and  $R^8$ , together with the atoms to which they are bound, are joined to form a heterocyclic, a substituted heterocyclic, a heteroaryl or substituted heteroaryl

30

group optionally containing from 1 to 3 additional hetero ring atoms selected from the group consisting of oxygen, nitrogen and sulfur;

R<sup>4\*</sup> is selected from the group consisting of hydrogen and alkyl;

5 R<sup>5</sup> is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

10 R<sup>6</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and -SO<sub>2</sub>R<sup>10</sup> where R<sup>10</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl;

15 R<sup>7</sup> and R<sup>8</sup> are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

20 R<sup>16</sup> and R<sup>17</sup> are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen; and

25 R<sup>18</sup> is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

R<sup>20</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted

cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;


R<sup>21</sup> is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;

**$b$  is 1 or 2;**

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

10 Preferably, the method employs a compound of formula III<sub>d</sub>, III<sub>e</sub> or  
III<sub>f</sub>.

15 In another of its method aspects, this invention is directed to a method for treating a disease mediated by VLA-4 in a patient, which method comprises administering a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of formula IVa and/or IVb:



25 wherein, in formula IVa, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form an aryl, cycloalkenyl, heteroaryl or heterocyclic group having at least five atoms in the aryl, cycloalkenyl, heteroaryl or heterocyclic group and optionally  
30 containing or additionally containing in the case of heteroaryl and

-- 15 --

heterocyclic groups 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heteroaryl or heterocyclic group is mono-cyclic;

5 in formula IVb, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W' to which they are bound respectively, are joined to form a cycloalkyl, cycloalkenyl or heterocyclic group having at least five atoms in the cycloalkyl, cycloalkenyl or heterocyclic group and optionally containing or additionally containing in the case the heterocyclic group 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and  
10 wherein the heterocyclic group is mono-cyclic;

and further wherein said aryl, cycloalkyl, cycloalkenyl, heteroaryl or heterocyclic group of formula IVa or IVb is optionally substituted, on any ring atom capable of substitution, with 1-3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen, hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino, guanidosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where each R is independently hydrogen or alkyl,  
25

-NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, -N[S(O)<sub>2</sub>-R']<sub>2</sub> and -N[S(O)<sub>2</sub>-NR']<sub>2</sub> where each R' is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

R<sup>13</sup> is selected from the group consisting of hydrogen, C<sub>1-10</sub> alkyl, Cy, and Cy-C<sub>1-10</sub> alkyl, wherein alkyl is optionally substituted with one to four substituents independently selected from R<sup>a</sup>; and Cy is optionally substituted with one to four substituents independently selected from R<sup>b</sup>;

R<sup>14</sup> is selected from the group consisting of hydrogen, C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, Cy, Cy-C<sub>1-10</sub> alkyl, Cy-C<sub>2-10</sub> alkenyl and Cy-C<sub>2-10</sub> alkynyl, wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents selected from phenyl and R<sup>x</sup>, and Cy is optionally substituted with one to four substituents independently selected from R<sup>y</sup>;

or R<sup>13</sup>, R<sup>14</sup> and the atoms to which they are attached together form a mono- or bicyclic ring containing 0-2 additional heteroatoms selected from N, O and S;

R<sup>15</sup> is selected from the group consisting of C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, aryl, aryl-C<sub>1-10</sub> alkyl, heteroaryl, heteroaryl-C<sub>1-10</sub> alkyl, wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents selected from R<sup>x</sup>, and aryl and heteroaryl are optionally substituted with one to four substituents independently selected from R<sup>y</sup>;

or R<sup>14</sup>, R<sup>15</sup> and the carbon to which they are attached form a 3-7 membered mono- or bicyclic ring containing 0-2 heteroatoms selected from

N, O and S;

-- 17 --

$R^a$  is selected from the group consisting of Cy and a group selected from  $R^s$ , wherein Cy is optionally substituted with one to four substituents independently selected from  $R^c$ ;

5  $R^b$  is selected from the group consisting of  $R^a$ ,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl  $C_{1-10}$  alkyl, heteroaryl  $C_{1-10}$  alkyl, wherein alkyl, alkenyl, alkynyl, aryl, heteroaryl are optionally substituted with a group independently selected from  $R^c$ ;

10  $R^c$  is selected from the group consisting of halogen,  $NO_2$ ,  $C(O)OR^f$ ,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy, aryl, aryl  $C_{1-4}$  alkyl, aryloxy, heteroaryl,  $NR^fR^g$ ,  $R^fC(O)R^g$ ,  $NR^fC(O)NR^fR^g$ , and CN;

$R^d$  and  $R^e$  are independently selected from hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, Cy and Cy  $C_{1-10}$  alkyl, wherein alkyl, alkenyl, alkynyl and Cy are optionally substituted with one to four substituents independently selected from  $R^c$ ;

15 or  $R^d$  and  $R^e$  together with the atoms to which they are attached form a heterocyclic ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen;

$R^f$  and  $R^g$  are independently selected from hydrogen,  $C_{1-10}$  alkyl, Cy and Cy- $C_{1-10}$  alkyl wherein Cy is optionally substituted with  $C_{1-10}$  alkyl; or  
20  $R^f$  and  $R^g$  together with the carbon to which they are attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

$R^h$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, cyano, aryl, aryl  $C_{1-10}$  alkyl, heteroaryl, heteroaryl  $C_{1-10}$  alkyl, and  $-SO_2R^i$ ; wherein alkyl, alkenyl, and alkynyl are optionally  
25 substituted with one to four substituents independently selected from  $R^a$ ; and aryl and heteroaryl are each optionally substituted with one to four substituents independently selected from  $R^b$ ;

$R^i$  is selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,

C<sub>2-10</sub> alkynyl, and aryl; wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from R<sup>c</sup>;

5 R<sup>x</sup> is selected from the group consisting of -OR<sup>d</sup>, -NO<sub>2</sub>, halogen, -S(O)<sub>m</sub>R<sup>d</sup>, -SR<sup>d</sup>, -S(O)<sub>2</sub>OR<sup>d</sup>, -S(O)<sub>m</sub>NR<sup>d</sup>R<sup>e</sup>, -NR<sup>d</sup>R<sup>e</sup>, -O(CR<sup>f</sup>R<sup>e</sup>)<sub>n</sub>NR<sup>d</sup>R<sup>e</sup>, -C(O)R<sup>d</sup>, -CO<sub>2</sub>R<sup>d</sup>, -CO<sub>2</sub>(CR<sup>f</sup>R<sup>e</sup>)<sub>n</sub>CONR<sup>d</sup>R<sup>e</sup>, -OC(O)R<sup>d</sup>, -CN, -C(O)NR<sup>d</sup>R<sup>e</sup>, -NR<sup>d</sup>C(O)R<sup>e</sup>, -OC(O)NR<sup>d</sup>R<sup>e</sup>, -NR<sup>d</sup>C(O)OR<sup>e</sup>, -NR<sup>d</sup>C(O)NR<sup>d</sup>R<sup>e</sup>, -CR<sup>d</sup>(N-OR<sup>e</sup>), CF<sub>3</sub>, oxo, NR<sup>d</sup>C(O)NR<sup>d</sup>SO<sub>2</sub>R<sup>i</sup>, NR<sup>d</sup>S(O)<sub>m</sub>R<sup>e</sup>, -OS(O)<sub>2</sub>OR<sup>d</sup>, and -OP(O)(OR<sup>d</sup>)<sub>2</sub>;

10 R<sup>y</sup> is selected from the group consisting of R<sup>x</sup>, C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, aryl C<sub>1-10</sub>alkyl, heteroaryl C<sub>1-10</sub> alkyl, cycloalkyl, heterocyclyl; wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from R<sup>x</sup>;

Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;

15 m is an integer from 1 to 2;

n is an integer from 1 to 10;

W is selected from the group consisting of carbon and nitrogen;

W' is selected from the group consisting of carbon, nitrogen, oxygen, sulfur, S(O) and S(O)<sub>2</sub>;

20 X' is selected from the group consisting of -C(O)OR<sup>d</sup>, -P(O)(OR<sup>d</sup>)(OR<sup>e</sup>), -P(O)(R<sup>d</sup>)(OR<sup>e</sup>), -S(O)<sub>m</sub>OR<sup>d</sup>, -C(O)NR<sup>d</sup>R<sup>h</sup>, and -5-tetrazolyl;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;

25 and further wherein the compound of formula IVa and/or IVb has a binding affinity to VLA-4 as expressed by an IC<sub>50</sub> of about 15 μM or less.

30 Preferably, in the above method, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form a heteroaryl or substituted heteroaryl group having two nitrogen atoms in the





-- 19 --

heteroaryl ring. Optionally, the heteroaryl ring may contain other heteroatoms such as oxygen or sulfur. More preferably,  $R^1$  and  $R^2$ , together with the carbon atom and W to which they are bound respectively, are joined to form a pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring; more preferably, a pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring; wherein the pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.


Preferably,  $X'$  is  $-C(O)OR^d$ .

In a preferred embodiment, the above method employs a compound of formula Va, Vc, Vd, Ve or Vf:




-- 20 --

5



Vd


10



Ve

15

20



Vf

wherein  $R^{13}$ ,  $R^{14}$ ,  $R^{15}$  and  $X'$  are as defined herein;

25

$R^5$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

30

$R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted

cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $-\text{SO}_2\text{R}^{10}$  where  $\text{R}^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl; and

$\text{R}^7$  and  $\text{R}^8$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$\text{R}^{16}$  and  $\text{R}^{17}$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen; and

$\text{R}^{18}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

$\text{R}^{20}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$\text{R}^{21}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;


$b$  is 1 or 2;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

-- 22 --

More preferably, the above method employs a compound of formula Vd, Ve or Vf.

In yet another of its method aspects, this invention is directed to a method for treating a disease mediated by VLA-4 in a patient, which method comprises administering a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of formula VIa and/or VIb:



wherein, in formula VIa, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form an aryl, cycloalkenyl, heteroaryl or heterocyclic group having at least five atoms in the aryl, cycloalkenyl, heteroaryl or heterocyclic group and optionally containing or additionally containing in the case of heteroaryl and heterocyclic groups 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heteroaryl or heterocyclic group is mono-cyclic;

in formula VIb, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W' to which they are bound respectively, are joined to form a cycloalkyl, cycloalkenyl or heterocyclic group having at least five atoms in the cycloalkyl, cycloalkenyl or heterocyclic group and optionally containing or additionally containing in the case of the heterocyclic group 1 to 3

heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heterocyclic group is mono-cyclic;

and further wherein said aryl, cycloalkyl, cycloalkenyl, heteroaryl or heterocyclic group of formula VIa or VIb is optionally substituted, on any  
5 ring atom capable of substitution, with 1-3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy,  
10 substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen, hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted  
15 thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-}$  substituted alkyl,  $-\text{OS}(\text{O})_2\text{-aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  
20  $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{OSO}_2\text{-NRR}$  where each R is independently hydrogen or alkyl,  $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-}$  substituted aryl,  $-\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-}$  alkyl,  $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-}$  substituted aryl,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heterocyclic}$  where R is hydrogen or alkyl,  $-\text{N}[\text{S}(\text{O})_2\text{-R}']_2$  and  $-\text{N}[\text{S}(\text{O})_2\text{-NR}']_2$  where each R' is independently selected from the group consisting of

alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

$R^{23}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl optionally substituted with one to four substituents independently selected from  $R^a$  and Cy optionally substituted with one to four substituents independently selected from  $R^b$ ;

$R^{24}$  is selected from the group consisting of  $Ar^1-Ar^2-C_{1-10}$  alkyl,  $Ar^1-Ar^2-C_{2-10}$  alkenyl,  $Ar^1-Ar^2-C_{2-10}$  alkynyl, wherein  $Ar^1$  and  $Ar^2$  are independently aryl or heteroaryl each of which is optionally substituted with one to four substituents independently selected from  $R^b$ ; alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents independently selected from  $R^a$ ;

$R^{25}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl, aryl  $C_{1-10}$ alkyl, heteroaryl, and heteroaryl  $C_{1-10}$  alkyl, wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents selected from  $R^a$ , and aryl and heteroaryl are optionally substituted with one to four substituents independently selected from  $R^b$ ;

$R^a$  is selected from the group consisting of Cy,  $-OR^d$ ,  $-NO_2$ , halogen  $-S(O)_mR^d$ ,  $-SR^d$ ,  $-S(O)_2OR^d$ ,  $-S(O)_mNR^dR^e$ ,  $-NR^dR^e$ ,  $-O(CR^fR^g)_nNR^dR^e$ ,  $-C(O)R^d$ ,  $-CO_2R^d$ ,  $-CO_2(CR^fR^g)_nCONR^dR^e$ ,  $-OC(O)R^d$ ,  $-CN$ ,  $-C(O)NR^dR^e$ ,  $-NR^dC(O)R^e$ ,  $-OC(O)NR^dR^e$ ,  $-NR^dC(O)OR^e$ ,  $-NR^dC(O)NR^dR^e$ ,  $-CR^d(N-OR^e)$ ,  $CF_3$ , and  $-OCF_3$ ;

wherein Cy is optionally substituted with one to four substituents independently selected from  $R^e$ ;

$R^b$  is selected from the group consisting of  $R^a$ ,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl  $C_{1-10}$  alkyl, heteroaryl  $C_{1-10}$ alkyl,

wherein alkyl, alkenyl, aryl, heteroaryl are optionally substituted with a group independently selected from  $R^e$ ;

$R^e$  is selected from the group consisting of halogen, amino, carboxy,

-- 25 --

C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, aryl, aryl C<sub>1-4</sub> alkyl, hydroxy, CF<sub>3</sub>, and aryloxy;

R<sup>d'</sup> and R<sup>e'</sup> are independently selected from hydrogen, C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, Cy and Cy C<sub>1-10</sub> alkyl, wherein alkyl, alkenyl, alkynyl and Cy are optionally substituted with one to four substituents  
5 independently selected from R<sup>e'</sup>; or R<sup>d'</sup> and R<sup>e'</sup> together with the atoms to which they are attached form a heterocyclic ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen;

R<sup>f'</sup> and R<sup>g'</sup> are independently selected from hydrogen, C<sub>1-10</sub> alkyl, Cy  
10 and Cy-C<sub>1-10</sub> alkyl; or R<sup>f'</sup> and R<sup>g'</sup> together with the carbon to which they are attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

R<sup>h'</sup> is selected from the group consisting of hydrogen, C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, cyano, aryl, aryl C<sub>1-10</sub> alkyl, heteroaryl,  
15 heteroaryl C<sub>1-10</sub> alkyl, or -SO<sub>2</sub>R<sup>i'</sup>;

wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from R<sup>a'</sup>; and aryl and heteroaryl are each optionally substituted with one to four substituents independently selected from R<sup>b'</sup>;

20 R<sup>i'</sup> is selected from the group consisting of C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, and aryl;

wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from R<sup>c'</sup>;

Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;

25 X<sup>n</sup> is selected from the group consisting of -C(O)OR<sup>d'</sup>, -P(O)(OR<sup>d'</sup>)(OR<sup>e'</sup>), -P(O)(R<sup>d'</sup>)(OR<sup>e'</sup>), -S(O)<sub>m</sub>OR<sup>d'</sup>, -C(O)NR<sup>d'</sup>R<sup>h'</sup>, and -5-tetrazolyl;

m is an integer from 1 to 2;

n is an integer from 1 to 10;

-- 26 --

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;

and further wherein the compound of formula VIa and/or VIb has a binding affinity to VLA-4 as expressed by an  $IC_{50}$  of about  $15\mu M$  or less.

5

Preferably, in the above method,  $R^1$  and  $R^2$ , together with the carbon atom and W to which they are bound respectively, are joined to form a heteroaryl or substituted heteroaryl group having two nitrogen atoms in the heteroaryl ring. Optionally, the heteroaryl ring may contain other  
10 heteroatoms such as oxygen or sulfur. More preferably,  $R^1$  and  $R^2$ , together with the carbon atom and W to which they are bound respectively, are joined to form a pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring; more preferably, a pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring; wherein the  
15 pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl,  
20 heterocyclic, substituted heterocyclic and halogen.

Preferably, in the above method,  $X^n$  is  $-C(O)OR^d$ .

25


Preferably,  $R^{24}$  is  $-\text{CH}_2-\text{Ar}^2-\text{Ar}^1$  and  $R^{25}$  is hydrogen.

In a preferred embodiment, the above method employs a compound of formula VIIa, VIIc, VIId, VIIe or VIIf:




-- 27 --

5




VIIa

10



VIIc


15



VIId

20


25



VIIe

30

5



VIIIf

wherein  $R^{23}$ ,  $R^{24}$ ,  $R^{25}$  and  $X''$  are as defined herein;

10

$R^5$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

15

$R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $-SO_2R^{10}$  where  $R^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl; and

20

$R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

25

$R^{16}$  and  $R^{17}$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen; and

$R^{18}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

5  $R^{20}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

10  $R^{21}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;

$b$  is 1 or 2;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

15

Preferably, the compound employed in the above method is selected from formula VIIId, VIIe or VIIf.


20 The compounds and pharmaceutical compositions of this invention are useful for treating disease conditions mediated by VLA-4 or leucocyte adhesion. Such disease conditions include, by way of example, asthma, Alzheimer's disease, atherosclerosis, AIDS dementia, diabetes (including acute juvenile onset diabetes), inflammatory bowel disease (including ulcerative colitis and Crohn's disease), multiple sclerosis, rheumatoid  
25 arthritis, tissue transplantation, tumor metastasis, meningitis, encephalitis, stroke, and other cerebral traumas, nephritis, retinitis, atopic dermatitis, psoriasis, myocardial ischemia and acute leukocyte-mediated lung injury such as that which occurs in adult respiratory distress syndrome.

-- 30 --

Other disease conditions include, but are not limited to, inflammatory conditions such as erythema nodosum, allergic conjunctivitis, optic neuritis, uveitis, allergic rhinitis, Ankylosing spondylitis, psoriatic arthritis, vasculitis, Reiter's syndrome, systemic lupus erythematosus, progressive  
5 systemic sclerosis, polymyositis, dermatomyositis, Wegner's granulomatosis, aortitis, sarcoidosis, lymphocytopenia, temporal arteritis, pericarditis, myocarditis, congestive heart failure, polyarteritis nodosa, hypersensitivity syndromes, allergy, hypereosinophilic syndromes, Churg-Strauss syndrome, chronic obstructive pulmonary disease, hypersensitivity  
10 pneumonitis, chronic active hepatitis, interstitial cystitis, autoimmune endocrine failure, primary biliary cirrhosis, autoimmune aplastic anemia, chronic persistent hepatitis and thyroiditis.

In a preferred embodiment, the disease condition mediated by VLA-4  
15 is an inflammatory disease.

The present invention is also directed to novel compounds useful for treating a disease condition mediated by VLA-4 or leucocyte adhesion. Accordingly, in one of its composition aspects, this invention is directed to a  
20 compound of formula Ia and/or Ib:



wherein, in formula Ia, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and  
30 W to which they are bound respectively, are joined to form an aryl,

-- 31 --

5 cycloalkenyl, heteroaryl or heterocyclic group having at least five atoms in the aryl, cycloalkenyl, heteroaryl or heterocyclic group and optionally containing or additionally containing in the case of heteroaryl and heterocyclic groups 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heteroaryl or heterocyclic group is mono-cyclic;

10 in formula Ib, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W' to which they are bound respectively, are joined to form a cycloalkyl, cycloalkenyl or heterocyclic group having at least five atoms in the cycloalkyl, cycloalkenyl or heterocyclic group and optionally containing or additionally containing in the case of the heterocyclic group 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heterocyclic group is mono-cyclic;

15 and further wherein said aryl, cycloalkyl, cycloalkenyl, heteroaryl or heterocyclic group of formula Ia or Ib is optionally substituted, on any ring atom capable of substitution, with 1-3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen, hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl,

20  
25  
30

-OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where each R is independently hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, -N[S(O)<sub>2</sub>-R']<sub>2</sub> and -N[S(O)<sub>2</sub>-NR']<sub>2</sub> where each R' is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

R<sup>3</sup> is -(CH<sub>2</sub>)<sub>x</sub>-Ar-R<sup>9</sup>, where Ar is aryl, substituted aryl, heteroaryl and substituted heteroaryl; R<sup>9</sup> is selected from the group consisting of acyl, acylamino, acyloxy, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, oxycarbonylamino, oxythiocarbonylamino, thioamidino, thiocarbonylamino, aminosulfonylamino, aminosulfonyloxy, aminosulfonyl, oxysulfonylamino and oxysulfonyl; x is an integer from 0 to 4;

R<sup>3'</sup> is selected from the group consisting of hydrogen, isopropyl, -CH<sub>2</sub>Z where Z is selected from the group consisting of hydrogen, hydroxyl, acylamino, alkyl, alkoxy, aryloxy, aryl, aryloxyaryl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted alkyl, substituted alkoxy, substituted aryl, substituted aryloxy, substituted aryloxyaryl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

-- 33 --

Q is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)<sub>2</sub>, and -NR<sup>4</sup>-;

R<sup>4</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic or, optionally, R<sup>4</sup> and R<sup>1</sup> or R<sup>4</sup> and R<sup>2</sup>, together with the atoms to which they are bound, are joined to form a heteroaryl, a substituted heteroaryl, a heterocyclic or a substituted heterocyclic group;

W is selected from the group consisting of nitrogen and carbon; and

W' is selected from the group consisting of nitrogen, carbon, oxygen, sulfur, S(O), and S(O)<sub>2</sub>;


X is selected from the group consisting of hydroxyl, alkoxy, substituted alkoxy, alkenoxy, substituted alkenoxy, cycloalkoxy, substituted cycloalkoxy, cycloalkenoxo, substituted cycloalkenoxo, aryloxy, substituted aryloxy, heteroaryloxy, substituted heteroaryloxy, heterocycloxy, substituted heterocycloxy and -NR<sup>n</sup>R<sup>n</sup> where each R<sup>n</sup> is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;

and further wherein the compound of formula Ia and/or Ib has a binding affinity to VLA-4 as expressed by an IC<sub>50</sub> of about 15 μM or less.

More preferably, R<sup>3</sup> is a group of the formula:

30



wherein  $R^9$  and  $x$  are as defined herein. Preferably,  $R^9$  is in the *para* position of the phenyl ring; and  $x$  is an integer of from 1 to 4, more preferably,  $x$  is 1.

5

Preferably,  $R^{3'}$  is hydrogen.


In a preferred embodiment,  $R^9$  is selected from the group consisting of  $-O-Z-NR^{11}R^{11'}$  and  $-O-Z-R^{12}$  wherein  $R^{11}$  and  $R^{11'}$  are independently  
10 selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, and where  $R^{11}$  and  $R^{11'}$  are joined to form a heterocycle or a substituted heterocycle,  $R^{12}$  is selected from the group consisting of heterocycle and substituted heterocycle, and  $Z$  is selected  
15 from the group consisting of  $-C(O)-$  and  $-SO_2-$ . More preferably,  $R^9$  is  $-OC(O)NR^{11}R^{11'}$ , wherein  $R^{11}$  and  $R^{11'}$  are as defined herein.

Preferably, in the above compounds,  $Z$  is  $-C(O)-$  and  $Q$  is preferably  $-NR^4-$ .

20

In a preferred embodiment, this invention is directed to compounds of formula IIa or IIb:


25



30



-- 35 --



5

wherein X is as defined herein;

$R^3$  is  $-(CH_2)_x-Ar-R^9$ , where Ar is aryl, substituted aryl, heteroaryl and substituted heteroaryl;  $R^9$  is selected from the group consisting of acyl, acylamino, acyloxy, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, oxycarbonylamino, oxythiocarbonylamino, thioamidino, thiocarbonylamino, aminosulfonylamino, aminosulfonyloxy, aminosulfonyl, oxysulfonylamino and oxysulfonyl; x is an integer from 0 to 4;

$R^{3'}$  is selected from the group consisting of hydrogen, isopropyl,  $-CH_2Z$  where Z is selected from the group consisting of hydrogen, hydroxyl, acylamino, alkyl, alkoxy, aryloxy, aryl, aryloxyaryl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted alkyl, substituted alkoxy, substituted aryl, substituted aryloxy, substituted aryloxyaryl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

ring A and ring B independently form a heteroaryl or substituted heteroaryl group having two nitrogen atoms in the heteroaryl ring;

$R^5$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

30

-- 36 --

R<sup>6</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and -SO<sub>2</sub>R<sup>10</sup> where R<sup>10</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl;

or optionally, one of, R<sup>4</sup> and ring A, R<sup>4</sup> and R<sup>5</sup>, R<sup>4</sup> and R<sup>6</sup>, or R<sup>5</sup> and R<sup>6</sup>, together with the atoms to which they are bound, can be joined to form a heterocyclic or substituted heterocyclic ring;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof; and provided that ring B does not form a 6-amino or substituted amino pyrimidin-4-yl group.

R<sup>3'</sup> is preferably hydrogen. Preferably,  $x$  is an integer from 1 to 4; more preferably,  $x$  is 1.

Preferably, ring A forms a pyridazine, pyrimidine or pyrazine ring; more preferably, a pyrimidine or pyrazine ring; wherein the pyridazine, pyrimidine or pyrazine ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.

Preferably, ring B forms a pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or a 1,1-dioxo-1,2,5-thiadiazole ring; more preferably, a pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or a 1,1-dioxo-1,2,5-thiadiazole ring; wherein the pyridazine, pyrimidine or pyrazine ring is optionally substituted with 1 to 3 substituents selected from the group


-- 37 --

consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.

5

In another preferred embodiment, this invention is directed to compounds of formula IIIa, IIIc, IIId, IIIe or IIIf:


10



IIIa


15

20



IIIc

25




IIIId

30


-- 38 --

5



IIIe

10



IIIf

15

wherein X is as defined herein;

$R^3$  is  $-(CH_2)_x-Ar-R^9$ , where Ar is aryl, substituted aryl, heteroaryl and substituted heteroaryl;  $R^9$  is selected from the group consisting of acyl, acylamino, acyloxy, aminoacyl, aminocarbonylamino, aminothiocabonylamino, aminocarbonyloxy, oxycarbonylamino, oxythiocarbonylamino, thioamidino, thiocarbonylamino, aminosulfonylamino, aminosulfonyloxy, aminosulfonyl, oxysulfonylamino and oxysulfonyl;  $x$  is an integer from 0 to 4;

25

$R^{3'}$  is selected from the group consisting of hydrogen, isopropyl,  $-CH_2Z$  where Z is selected from the group consisting of hydrogen, hydroxyl, acylamino, alkyl, alkoxy, aryloxy, aryl, aryloxyaryl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic,

30

carboxyl-substituted heterocyclic, cycloalkyl, substituted alkyl, substituted alkoxy, substituted aryl, substituted aryloxy, substituted aryloxyaryl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

5           R<sup>4'</sup> is selected from the group consisting of hydrogen and alkyl or, optionally, one of, R<sup>4'</sup> and R<sup>5</sup>, R<sup>4'</sup> and R<sup>6</sup>, R<sup>5</sup> and R<sup>6</sup>, R<sup>5</sup> and R<sup>8</sup>, or R<sup>6</sup> and R<sup>8</sup>, together with the atoms to which they are bound, are joined to form a heterocyclic, a substituted heterocyclic, a heteroaryl or substituted heteroaryl group optionally containing from 1 to 3 additional hetero ring atoms selected  
10       from the group consisting of oxygen, nitrogen and sulfur;

          R<sup>4'</sup> is selected from the group consisting of hydrogen and alkyl;

          R<sup>5</sup> is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted  
15       heterocyclic, heteroaryl and substituted heteroaryl;

          R<sup>6</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and -SO<sub>2</sub>R<sup>10</sup> where R<sup>10</sup> is selected from the  
20       group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl;

          R<sup>7</sup> and R<sup>8</sup> are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted  
25       heterocyclic and halogen;

          R<sup>16</sup> and R<sup>17</sup> are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl,

-- 40 --

heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen; and

5  $R^{18}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

10  $R^{20}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$R^{21}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;


$b$  is 1 or 2;

15 and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

In the above compounds,  $R^3$  is preferably hydrogen. Preferably,  $x$  is an integer from 1 to 4; more preferably,  $x$  is 1.

20 Preferably, the compound is selected from formula IIIId, IIIe or IIIf.

In another of its composition aspects, this invention is directed to a compound of formula IVa:



wherein R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form a heteroaryl group having two nitrogen atoms in the heteroaryl ring;

and further wherein said heteroaryl group is optionally substituted, on  
5 any ring atom capable of substitution, with 1-3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy,  
10 substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen, hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted  
15 thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl,  
20 -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where each R is independently hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-  
25 substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl,

$-N[S(O)_2-R']_2$  and  $-N[S(O)_2-NR']_2$  where each  $R'$  is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

5  $R^{13}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl, Cy, and Cy- $C_{1-10}$  alkyl, wherein alkyl is optionally substituted with one to four substituents independently selected from  $R^a$ ; and Cy is optionally substituted with one to four substituents independently selected from  $R^b$ ;

10  $R^{14}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, Cy, Cy- $C_{1-10}$  alkyl, Cy- $C_{2-10}$  alkenyl and Cy- $C_{2-10}$  alkynyl, wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents selected from phenyl and  $R^x$ , and Cy is optionally substituted with one to four substituents independently selected from  $R^y$ ;

15 or  $R^{13}$ ,  $R^{14}$  and the atoms to which they are attached together form a mono- or bicyclic ring containing 0-2 additional heteratoms selected from N, O and S;

20  $R^{15}$  is selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl, aryl- $C_{1-10}$  alkyl, heteroaryl, heteroaryl- $C_{1-10}$  alkyl, wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents selected from  $R^x$ , and aryl and heteroaryl are optionally substituted with one to four substituents independently selected from  $R^y$ ;

or  $R^{14}$ ,  $R^{15}$  and the carbon to which they are attached form a 3-7 membered mono- or bicyclic ring containing 0-2 heteroatoms selected from N, O and S;

25  $R^a$  is selected from the group consisting of Cy and a group selected from  $R^x$ , wherein Cy is optionally substituted with one to four substituents independently selected from  $R^c$ ;

30  $R^b$  is selected from the group consisting of  $R^a$ ,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl  $C_{1-10}$  alkyl, heteroaryl  $C_{1-10}$  alkyl, wherein alkyl, alkenyl, alkynyl, aryl, heteroaryl are optionally substituted with a group independently selected from  $R^c$ ;



$R^c$  is selected from the group consisting of halogen,  $\text{NO}_2$ ,  $\text{C}(\text{O})\text{OR}^f$ ,  $\text{C}_{1-4}$  alkyl,  $\text{C}_{1-4}$  alkoxy, aryl, aryl  $\text{C}_{1-4}$  alkyl, aryloxy, heteroaryl,  $\text{NR}^f\text{R}^g$ ,  $\text{R}^f\text{C}(\text{O})\text{R}^g$ ,  $\text{NR}^f\text{C}(\text{O})\text{NR}^f\text{R}^g$ , and  $\text{CN}$ ;

$R^d$  and  $R^e$  are independently selected from hydrogen,  $\text{C}_{1-10}$  alkyl,  $\text{C}_{2-10}$  alkenyl,  $\text{C}_{2-10}$  alkynyl, Cy and Cy  $\text{C}_{1-10}$  alkyl, wherein alkyl, alkenyl, alkynyl and Cy are optionally substituted with one to four substituents independently selected from  $R^c$ ;

or  $R^d$  and  $R^e$  together with the atoms to which they are attached form a heterocyclic ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen;

$R^f$  and  $R^g$  are independently selected from hydrogen,  $\text{C}_{1-10}$  alkyl, Cy and Cy- $\text{C}_{1-10}$  alkyl wherein Cy is optionally substituted with  $\text{C}_{1-10}$  alkyl; or  $R^f$  and  $R^g$  together with the carbon to which they are attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

$R^h$  is selected from the group consisting of hydrogen,  $\text{C}_{1-10}$  alkyl,  $\text{C}_{2-10}$  alkenyl,  $\text{C}_{2-10}$  alkynyl, cyano, aryl, aryl  $\text{C}_{1-10}$  alkyl, heteroaryl, heteroaryl  $\text{C}_{1-10}$  alkyl, and  $-\text{SO}_2\text{R}^i$ ; wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from  $R^a$ ; and aryl and heteroaryl are each optionally substituted with one to four substituents independently selected from  $R^b$ ;

$R^i$  is selected from the group consisting of  $\text{C}_{1-10}$  alkyl,  $\text{C}_{2-10}$  alkenyl,  $\text{C}_{2-10}$  alkynyl, and aryl; wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from  $R^c$ ;

$R^x$  is selected from the group consisting of  $-\text{OR}^d$ ,  $-\text{NO}_2$ , halogen,  $-\text{S}(\text{O})_m\text{R}^d$ ,  $-\text{SR}^d$ ,  $-\text{S}(\text{O})_2\text{OR}^d$ ,  $-\text{S}(\text{O})_m\text{NR}^d\text{R}^e$ ,  $-\text{NR}^d\text{R}^e$ ,  $-\text{O}(\text{CR}^f\text{R}^g)_n\text{NR}^d\text{R}^e$ ,  $-\text{C}(\text{O})\text{R}^d$ ,  $-\text{CO}_2\text{R}^d$ ,  $-\text{CO}_2(\text{CR}^f\text{R}^g)_n\text{CONR}^d\text{R}^e$ ,  $-\text{OC}(\text{O})\text{R}^d$ ,  $-\text{CN}$ ,  $-\text{C}(\text{O})\text{NR}^d\text{R}^e$ ,  $-\text{NR}^d\text{C}(\text{O})\text{R}^e$ ,  $-\text{OC}(\text{O})\text{NR}^d\text{R}^e$ ,  $-\text{NR}^d\text{C}(\text{O})\text{OR}^e$ ,  $-\text{NR}^d\text{C}(\text{O})\text{NR}^d\text{R}^e$ ,  $-\text{CR}^d(\text{N}-\text{OR}^e)$ ,  $\text{CF}_3$ , oxo,  $\text{NR}^d\text{C}(\text{O})\text{NR}^d\text{SO}_2\text{R}^i$ ,  $\text{NR}^d\text{S}(\text{O})_m\text{R}^e$ ,  $-\text{OS}(\text{O})_2\text{OR}^d$ , and

-OP(O)(OR<sup>d</sup>)<sub>2</sub>;

R<sup>y</sup> is selected from the group consisting of R<sup>x</sup>, C<sub>1-10</sub> alkyl, C<sub>2-10</sub> alkenyl, C<sub>2-10</sub> alkynyl, aryl C<sub>1-10</sub>alkyl, heteroaryl C<sub>1-10</sub> alkyl, cycloalkyl, heterocyclyl; wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from R<sup>x</sup>;

Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;

m is an integer from 1 to 2;

n is an integer from 1 to 10;

W is selected from the group consisting of carbon and nitrogen;

W' is selected from the group consisting of carbon, nitrogen, oxygen, sulfur, S(O) and S(O)<sub>2</sub>;

X' is selected from the group consisting of -C(O)OR<sup>d</sup>, -P(O)(OR<sup>d</sup>)(OR<sup>e</sup>), -P(O)(R<sup>d</sup>)(OR<sup>e</sup>), -S(O)<sub>m</sub>OR<sup>d</sup>, -C(O)NR<sup>d</sup>R<sup>h</sup>, and -5-tetrazolyl;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;

and further wherein the compound of formula IV has a binding affinity to VLA-4 as expressed by an IC<sub>50</sub> of about 15μM or less;

and provided that when R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form a 2-arylpyrimidin-4-yl group and R<sup>14</sup> is hydrogen, then R<sup>15</sup> is not alkyl of from 1 to 6 carbon atoms optionally substituted with hydroxyl; and when R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form a 5-arylpyrazin-2-yl group and R<sup>14</sup> is hydrogen, then R<sup>15</sup> is not 4-hydroxybenzyl.



In the above compounds, R<sup>1</sup> and R<sup>2</sup> are preferably joined to form an pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring; more preferably, a pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring; wherein the pyridazine,

-- 45 --

pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.


Preferably, in the above compounds, X' is -C(O)OR<sup>d</sup>.

In a preferred embodiment, this invention is directed to compounds of formula Va, Vc, Vd, Ve or Vf:




-- 46 --

5



Vd


10



Ve

15

20



Vf

wherein  $R^{13}$ ,  $R^{14}$ ,  $R^{15}$  and  $X'$  are as defined herein;

25

$R^5$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

$R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted

30

cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $-\text{SO}_2\text{R}^{10}$  where  $\text{R}^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl; and

$\text{R}^7$  and  $\text{R}^8$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$\text{R}^{16}$  and  $\text{R}^{17}$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen; and

$\text{R}^{18}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

$\text{R}^{20}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;


$\text{R}^{21}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;

$b$  is 1 or 2;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

More preferably, the compound is selected from formula Vd, Ve or Vf.

In yet another of its composition aspects, this invention is directed to  
 5 a compound of formula VIa and/or VIb:



15 wherein, in formula VIa, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form an aryl, cycloalkenyl, heteroaryl or heterocyclic group having at least five atoms in the aryl, cycloalkenyl, heteroaryl or heterocyclic group and optionally containing or additionally containing in the case of heteroaryl and heterocyclic groups 1 to 3 heteroatoms selected from the group consisting of  
 20 oxygen, nitrogen and sulfur, and wherein the heteroaryl or heterocyclic group is mono-cyclic;

in formula VIb, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W' to which they are bound respectively, are joined to form a cycloalkyl, cycloalkenyl or heterocyclic group having at least five atoms in the  
 25 cycloalkyl, cycloalkenyl or heterocyclic group and optionally containing or additionally containing in the case of the heterocyclic group 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heterocyclic group is mono-cyclic;

and further wherein said aryl, cycloalkyl, cycloalkenyl, heteroaryl or  
 30 heterocyclic group of formula VIa or VIb is optionally substituted, on any

ring atom capable of substitution, with 1-3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino,

5 aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen, hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl,

10 thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-}$

15 substituted alkyl,  $-\text{OS}(\text{O})_2\text{-aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{OSO}_2\text{-NRR}$  where each R is independently hydrogen or alkyl,  $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-}$

20 substituted aryl,  $-\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-}$

substituted aryl,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heterocyclic}$  where R is hydrogen or alkyl,  $-\text{N}[\text{S}(\text{O})_2\text{-R}']_2$  and  $-\text{N}[\text{S}(\text{O})_2\text{-NR}']_2$  where each R' is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

25

$\text{R}^{23}$  is selected from the group consisting of hydrogen,  $\text{C}_{1-10}$  alkyl optionally substituted with one to four substituents independently selected

from  $R^a$  and Cy optionally substituted with one to four substituents independently selected from  $R^b$ ;

$R^{24}$  is selected from the group consisting of  $Ar^1-Ar^2-C_{1-10}$  alkyl,  $Ar^1-Ar^2-C_{2-10}$  alkenyl,  $Ar^1-Ar^2-C_{2-10}$  alkynyl, wherein  $Ar^1$  and  $Ar^2$  are independently aryl or heteroaryl each of which is optionally substituted with one to four substituents independently selected from  $R^b$ ; alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents independently selected from  $R^a$ ;

$R^{25}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl, aryl  $C_{1-10}$ alkyl, heteroaryl, and heteroaryl  $C_{1-10}$  alkyl, wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents selected from  $R^a$ , and aryl and heteroaryl are optionally substituted with one to four substituents independently selected from  $R^b$ ;

$R^a$  is selected from the group consisting of Cy,  $-OR^d$ ,  $-NO_2$ , halogen  $-S(O)_mR^d$ ,  $-SR^d$ ,  $-S(O)_2OR^d$ ,  $-S(O)_mNR^dR^c$ ,  $-NR^dR^c$ ,  $-O(CR^fR^g)_nNR^dR^c$ ,  $-C(O)R^d$ ,  $-CO_2R^d$ ,  $-CO_2(CR^fR^g)_nCONR^dR^c$ ,  $-OC(O)R^d$ ,  $-CN$ ,  $-C(O)NR^dR^c$ ,  $-NR^dC(O)R^c$ ,  $-OC(O)NR^dR^c$ ,  $-NR^dC(O)OR^c$ ,  $-NR^dC(O)NR^dR^c$ ,  $-CR^d(N-OR^c)$ ,  $CF_3$ , and  $-OCF_3$ ;

wherein Cy is optionally substituted with one to four substituents independently selected from  $R^c$ ;

$R^b$  is selected from the group consisting of  $R^a$ ,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl  $C_{1-10}$  alkyl, heteroaryl  $C_{1-10}$ alkyl,

wherein alkyl, alkenyl, aryl, heteroaryl are optionally substituted with a group independently selected from  $R^c$ ;

$R^c$  is selected from the group consisting of halogen, amino, carboxy,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy, aryl, aryl  $C_{1-4}$ alkyl, hydroxy,  $CF_3$ , and aryloxy;

$R^d$  and  $R^e$  are independently selected from hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, Cy and Cy  $C_{1-10}$ alkyl, wherein alkyl, alkenyl, alkynyl and Cy are optionally substituted with one to four substituents



independently selected from  $R^c$ ; or  $R^d$  and  $R^e$  together with the atoms to which they are attached form a heterocyclic ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen;

5  $R^f$  and  $R^g$  are independently selected from hydrogen,  $C_{1-10}$  alkyl, Cy and Cy- $C_{1-10}$  alkyl; or  $R^f$  and  $R^g$  together with the carbon to which they are attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

10  $R^h$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, cyano, aryl, aryl  $C_{1-10}$  alkyl, heteroaryl, heteroaryl  $C_{1-10}$  alkyl, or  $-SO_2R^i$ ;

wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from  $R^a$ ; and aryl and heteroaryl are each optionally substituted with one to four substituents  
15 independently selected from  $R^b$ ;

$R^i$  is selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, and aryl;

wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from  $R^c$ ;

20 Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;

$X''$  is selected from the group consisting of  $-C(O)OR^d$ ,  $-P(O)(OR^d)(OR^e)$ ,  $-P(O)(R^d)(OR^e)$ ,  $-S(O)_mOR^d$ ,  $-C(O)NR^dR^h$ , and -5-tetrazolyl;

$m$  is an integer from 1 to 2;

25  $n$  is an integer from 1 to 10;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;


and further wherein the compounds of formula VIa and/or VIb have a binding affinity to VLA-4 as expressed by an  $IC_{50}$  of about  $15\mu M$  or less.

In the above compounds, R<sup>1</sup> and R<sup>2</sup> are preferably joined to form a heteroaryl or substituted heteroaryl group having two nitrogen atoms in the heteroaryl ring. More preferably, R<sup>1</sup> and R<sup>2</sup> are joined to form a pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring; more preferably, a pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring; wherein the pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.

Preferably, X" is -C(O)OR<sup>d'</sup>.


In the above compounds, R<sup>24</sup> is preferably -CH<sub>2</sub>-Ar<sup>2</sup>-Ar<sup>1</sup> and R<sup>25</sup> is preferably hydrogen.

In a preferred embodiment, this invention is directed to compounds of formula VIIa, VIIc, VIId, VIIe or VIIf:




VIIa

5




VIIc

10




VIIId

15



VIIe

20



VIIIf

25

wherein  $R^{24}$ ,  $R^{25}$  and  $X''$  are as defined herein;

$R^5$  is selected from the group consisting of alkyl, substituted alkyl,  
30 alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted

cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

$R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $-SO_2R^{10}$  where  $R^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl; and

$R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$R^{16}$  and  $R^{17}$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen; and

$R^{18}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

$R^{20}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$R^{21}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;

$b$  is 1 or 2;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

Preferably, the compound is selected from formula VIId, VIIE or  
5 VIIf.

This invention also provides pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of the compounds defined herein.

10

In the above compounds, when X is other than -OH or pharmaceutical salts thereof, X is preferably a substituent which will convert (e.g., hydrolyze, metabolize, etc.) *in vivo* to a compound where X is -OH or a salt thereof. Accordingly, suitable X groups are any art recognized  
15 pharmaceutically acceptable groups which will hydrolyze or otherwise convert *in vivo* to a hydroxyl group or a salt thereof including, by way of example, esters (X is alkoxy, substituted alkoxy, cycloalkoxy, substituted cycloalkoxy, alkenoxy, substituted alkenoxy, cycloalkenoxo, substituted cycloalkenoxo, aryloxy, substituted aryloxy, heteroaryloxy, substituted  
20 heteroaryloxy, heterocycloxy, substituted heterocycloxy, and the like).

Unless otherwise defined, R<sup>3</sup> and R<sup>15</sup> in the above compounds are preferably selected from all possible isomers arising by substitution with the following groups:

25 4-methylbenzyl,  
4-hydroxybenzyl,  
4-methoxybenzyl,  
4-*t*-butoxybenzyl,  
4-benzyloxybenzyl,  
30 4-[ $\phi$ -CH(CH<sub>3</sub>)O-]benzyl,

- 4-[ $\phi$ -CH(COOH)O-]benzyl,  
4-[BocNHCH<sub>2</sub>C(O)NH-]benzyl,  
4-chlorobenzyl,  
4-[NH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl,  
5 4-carboxybenzyl,  
4-[CbzNHCH<sub>2</sub>CH<sub>2</sub>NH-]benzyl,  
3-hydroxy-4-( $\phi$ -OC(O)NH-)benzyl,  
4-[HOOCCH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl,  
benzyl,  
10 4-[2'-carboxylphenoxy-]benzyl,  
4-[ $\phi$ -C(O)NH-]benzyl,  
3-carboxybenzyl,  
4-iodobenzyl,  
4-hydroxy-3,5-diiodobenzyl,  
15 4-hydroxy-3-iodobenzyl,  
4-[2'-carboxyphenyl-]benzyl,  
 $\phi$ -CH<sub>2</sub>CH<sub>2</sub>-,  
4-nitrobenzyl,  
2-carboxybenzyl,  
20 4-[dibenzylamino]-benzyl,  
4-[(1'-cyclopropyl)piperidin-4'-yl]C(O)NH-]benzyl,  
4-[-NHC(O)CH<sub>2</sub>NHBoc]benzyl,  
4-carboxybenzyl,  
4-hydroxy-3-nitrobenzyl,  
25 4-[-NHC(O)CH(CH<sub>3</sub>)NHBoc]benzyl,  
4-[-NHC(O)CH(CH<sub>2</sub> $\phi$ )NHBoc]benzyl,  
isobutyl,  
methyl,  
4-[CH<sub>3</sub>C(O)NH-]benzyl,  
30 -CH<sub>2</sub>-(3-indolyl),

- n*-butyl,  
*t*-butyl-OC(O)CH<sub>2</sub>-,  
*t*-butyl-OC(O)CH<sub>2</sub>CH<sub>2</sub>-,  
H<sub>2</sub>NC(O)CH<sub>2</sub>-,  
5 H<sub>2</sub>NC(O)CH<sub>2</sub>CH<sub>2</sub>-,  
BocNH-(CH<sub>2</sub>)<sub>4</sub>-,  
*t*-butyl-OC(O)-(CH<sub>2</sub>)<sub>2</sub>-,  
HOOCCH<sub>2</sub>-,  
HOOC(CH<sub>2</sub>)<sub>2</sub>-,  
10 H<sub>2</sub>N(CH<sub>2</sub>)<sub>4</sub>-,  
isopropyl,  
(1-naphthyl)-CH<sub>2</sub>-,  
(2-naphthyl)-CH<sub>2</sub>-,  
(2-thiophenyl)-CH<sub>2</sub>-,  
15 (φ-CH<sub>2</sub>-OC(O)NH-(CH<sub>2</sub>)<sub>4</sub>-,  
cyclohexyl-CH<sub>2</sub>-,  
benzyloxy-CH<sub>2</sub>-,  
HOCH<sub>2</sub>-,  
5-(3-N-benzyl)imidazolyl-CH<sub>2</sub>-,  
20 2-pyridyl-CH<sub>2</sub>-,  
3-pyridyl-CH<sub>2</sub>-,  
4-pyridyl-CH<sub>2</sub>-,  
5-(3-N-methyl)imidazolyl-CH<sub>2</sub>-,  
N-benzylpiperid-4-yl-CH<sub>2</sub>-,  
25 N-Boc-piperidin-4-yl-CH<sub>2</sub>-,  
N-(phenyl-carbonyl)piperidin-4-yl-CH<sub>2</sub>-,  
H<sub>3</sub>CSCH<sub>2</sub>CH<sub>2</sub>-,  
1-N-benzylimidazol-4-yl-CH<sub>2</sub>-,  
*iso*-propyl-C(O)NH-(CH<sub>2</sub>)<sub>4</sub>-,  
30 *iso*-butyl-C(O)NH-(CH<sub>2</sub>)<sub>4</sub>-,

- phenyl-C(O)NH-(CH<sub>2</sub>)<sub>4</sub>-,  
benzyl-C(O)NH-(CH<sub>2</sub>)<sub>4</sub>-,  
allyl-C(O)NH-(CH<sub>2</sub>)<sub>4</sub>-,  
4-(3-N-methylimidazolyl)-CH<sub>2</sub>-,  
5 4-imidazolyl,  
4-[(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-O]-benzyl,  
4-[(benzyl)<sub>2</sub>N]-benzyl,  
4-aminobenzyl,  
allyloxy-C(O)NH(CH<sub>2</sub>)<sub>4</sub>-,  
10 allyloxy-C(O)NH(CH<sub>2</sub>)<sub>3</sub>-,  
allyloxy-C(O)NH(CH<sub>2</sub>)<sub>2</sub>-,  
NH<sub>2</sub>C(O)CH<sub>2</sub>-,  
φ-CH=,  
2-pyridyl-C(O)NH-(CH<sub>2</sub>)<sub>4</sub>-,  
15 4-methylpyrid-3-yl-C(O)NH-(CH<sub>2</sub>)<sub>4</sub>-,  
3-methylthien-2-yl-C(O)NH-(CH<sub>2</sub>)<sub>4</sub>-,  
2-pyrrolyl-C(O)NH-(CH<sub>2</sub>)<sub>4</sub>-,  
2-furanyl-C(O)NH-(CH<sub>2</sub>)<sub>4</sub>-,  
4-methylphenyl-SO<sub>2</sub>-N(CH<sub>3</sub>)CH<sub>2</sub>C(O)NH(CH<sub>2</sub>)<sub>4</sub>-,  
20 4-[cyclopentylacetylenyl]-benzyl,  
4-[-NHC(O)-(N-Boc)-pyrrolidin-2-yl]-benzyl-,  
1-N-methylimidazol-4-yl-CH<sub>2</sub>-,  
1-N-methylimidazol-5-yl-CH<sub>2</sub>-,  
imidazol-5-yl-CH<sub>2</sub>-,  
25 6-methylpyrid-3-yl-C(O)NH-(CH<sub>2</sub>)<sub>4</sub>-,  
4-[2'-carboxymethylphenyl]-benzyl,  
4-[-NHC(O)NHCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-φ]-benzyl,  
4-[-NHC(O)NHCH<sub>2</sub>CH<sub>2</sub>-φ]-benzyl,  
-CH<sub>2</sub>C(O)NH(CH<sub>2</sub>)<sub>4</sub>φ,  
30 4-[φ(CH<sub>2</sub>)<sub>4</sub>O]-benzyl,



- 4-[-C≡C-φ-4'-φ]-benzyl,  
 4-[-C≡C-CH<sub>2</sub>-O-S(O)<sub>2</sub>-4'-CH<sub>3</sub>-φ]-benzyl,  
 4-[-C≡C-CH<sub>2</sub>NHC(O)NH<sub>2</sub>]-benzyl,  
 4-[-C≡C-CH<sub>2</sub>-O-4'-COOCH<sub>2</sub>CH<sub>3</sub>-φ]-benzyl,  
 5 4-[-C≡C-CH(NH<sub>2</sub>)-cyclohexyl]-benzyl,  
 -(CH<sub>2</sub>)<sub>4</sub>NHC(O)CH<sub>2</sub>-3-indolyl,  
 -(CH<sub>2</sub>)<sub>4</sub>NHC(O)CH<sub>2</sub>CH<sub>2</sub>-3-indolyl,  
 -(CH<sub>2</sub>)<sub>4</sub>NHC(O)-3-(5-methoxyindolyl),  
 -(CH<sub>2</sub>)<sub>4</sub>NHC(O)-3-(1-methylindolyl),  
 10 -(CH<sub>2</sub>)<sub>4</sub>NHC(O)-4-(-SO<sub>2</sub>(CH<sub>3</sub>)-φ),  
 -(CH<sub>2</sub>)<sub>4</sub>NHC(O)-4-(C(O)CH<sub>3</sub>)-phenyl,  
 -(CH<sub>2</sub>)<sub>4</sub>NHC(O)-4-fluorophenyl,  
 -(CH<sub>2</sub>)<sub>4</sub>NHC(O)CH<sub>2</sub>O-4-fluorophenyl,  
 4-[-C≡C-(2-pyridyl)]benzyl,  
 15 4-[-C≡C-CH<sub>2</sub>-O-phenyl]benzyl,  
 4-[-C≡C-CH<sub>2</sub>OCH<sub>3</sub>]benzyl,  
 4-[-C≡C-(3-hydroxyphenyl)]benzyl,  
 4-[-C≡C-CH<sub>2</sub>-O-4'-(-C(O)OC<sub>2</sub>H<sub>5</sub>)phenyl]benzyl,  
 4-[-C≡C-CH<sub>2</sub>CH(C(O)OCH<sub>3</sub>)<sub>2</sub>]benzyl,  
 20 4-[-C≡C-CH<sub>2</sub>NH-(4,5-dihydro-4-oxo-5-phenyl-oxazol-2-yl),  
 3-aminobenzyl,  
 4-[-C≡C-CH<sub>2</sub>CH(NHC(O)CH<sub>3</sub>)C(O)OH]-benzyl,  
 -CH<sub>2</sub>C(O)NHCH(CH<sub>3</sub>)φ,  
 -CH<sub>2</sub>C(O)NHCH<sub>2</sub>-(4-dimethylamino)-φ,  
 25 -CH<sub>2</sub>C(O)NHCH<sub>2</sub>-4-nitrophenyl,  
 -CH<sub>2</sub>CH<sub>2</sub>C(O)N(CH<sub>3</sub>)CH<sub>2</sub>-φ,  
 -CH<sub>2</sub>CH<sub>2</sub>C(O)NHCH<sub>2</sub>CH<sub>2</sub>-(N-methyl)-2-pyrrolyl,  
 -CH<sub>2</sub>CH<sub>2</sub>C(O)NHCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>,  
 -CH<sub>2</sub>CH<sub>2</sub>C(O)NHCH<sub>2</sub>CH<sub>2</sub>-3-indolyl,  
 30 -CH<sub>2</sub>C(O)N(CH<sub>3</sub>)CH<sub>2</sub>phenyl,

- CH<sub>2</sub>C(O)NH(CH<sub>2</sub>)<sub>2</sub>-(N-methyl)-2-pyrrolyl,
- CH<sub>2</sub>C(O)NHCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>,
- CH<sub>2</sub>C(O)NHCH<sub>2</sub>CH<sub>2</sub>-3-indolyl,
- 5    -(CH<sub>2</sub>)<sub>2</sub>C(O)NHCH(CH<sub>3</sub>)φ,
- (CH<sub>2</sub>)<sub>2</sub>C(O)NHCH<sub>2</sub>-4-dimethylaminophenyl,
- (CH<sub>2</sub>)<sub>2</sub>C(O)NHCH<sub>2</sub>-4-nitrophenyl,
- CH<sub>2</sub>C(O)NH-4-[-NHC(O)CH<sub>3</sub>-phenyl],
- CH<sub>2</sub>C(O)NH-4-pyridyl,
- CH<sub>2</sub>C(O)NH-4-[dimethylaminophenyl],
- 10    -CH<sub>2</sub>C(O)NH-3-methoxyphenyl,
- CH<sub>2</sub>CH<sub>2</sub>C(O)NH-4-chlorophenyl,
- CH<sub>2</sub>CH<sub>2</sub>C(O)NH-2-pyridyl,
- CH<sub>2</sub>CH<sub>2</sub>C(O)NH-4-methoxyphenyl,
- CH<sub>2</sub>CH<sub>2</sub>C(O)NH-3-pyridyl,
- 15    4-[(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>O-]benzyl,
- (CH<sub>2</sub>)<sub>3</sub>NHC(NH)NH-SO<sub>2</sub>-4-methylphenyl,
- 4-[(CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>O-]benzyl,
- (CH<sub>2</sub>)<sub>4</sub>NHC(O)NHCH<sub>2</sub>CH<sub>3</sub>,
- (CH<sub>2</sub>)<sub>4</sub>NHC(O)NH-phenyl,
- 20    -(CH<sub>2</sub>)<sub>4</sub>NHC(O)NH-4-methoxyphenyl,
- 4-[4'-pyridyl-C(O)NH-]benzyl,
- 4-[3'-pyridyl-C(O)NH-]benzyl,
- 4-[-NHC(O)NH-3'-methylphenyl]benzyl,
- 4-[-NHC(O)CH<sub>2</sub>NHC(O)NH-3'-methylphenyl]benzyl,
- 25    4-[-NHC(O)-(2',3'-dihydroindol-2-yl)]benzyl,
- 4-[-NHC(O)-(2',3'-dihydro-N-Boc-indol-2-yl)]benzyl,
- p-[-OCH<sub>2</sub>CH<sub>2</sub>-1'-(4'-pyrimidinyl)-piperazinyl]benzyl,
- 4-[-OCH<sub>2</sub>CH<sub>2</sub>-(1'-piperidinyl)]benzyl,
- 4-[-OCH<sub>2</sub>CH<sub>2</sub>-(1'-pyrrolidinyl)]benzyl,
- 30    4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(1'-piperidinyl)]benzyl-,

- CH<sub>2</sub>-3-(1,2,4-triazolyl),  
4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-4-(3'-chlorophenyl)-piperazin-1-yl]benzyl,  
4-[-OCH<sub>2</sub>CH<sub>2</sub>N(φ)CH<sub>2</sub>CH<sub>3</sub>]benzyl,  
4-[-OCH<sub>2</sub>-3'-(N-Boc)-piperidiny]benzyl,  
5 4-[di-*n*-pentylamino]benzyl,  
4-[*n*-pentylamino]benzyl,  
4-[di-*iso*-propylamino-CH<sub>2</sub>CH<sub>2</sub>O-]benzyl,  
4-[-OCH<sub>2</sub>CH<sub>2</sub>-(N-morpholinyl)]benzyl,  
4-[-O-(3'-(N-Boc)-piperidiny]benzyl,  
10 4-[-OCH<sub>2</sub>CH(NHBoc)CH<sub>2</sub>cyclohexyl]benzyl,  
*p*-[OCH<sub>2</sub>CH<sub>2</sub>-(N-piperidiny]benzyl,  
4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(4-*m*-chlorophenyl)-piperazin-1-yl]benzyl,  
4-[-OCH<sub>2</sub>CH<sub>2</sub>-(N-homopiperidiny]benzyl,  
4-[-NHC(O)-3'-(N-Boc)-piperidiny]benzyl,  
15 4-[-OCH<sub>2</sub>CH<sub>2</sub>N(benzyl)<sub>2</sub>]benzyl,  
-CH<sub>2</sub>-2-thiazolyl,  
3-hydroxybenzyl,  
4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>]benzyl,  
4-[-NHC(S)NHCH<sub>2</sub>CH<sub>2</sub>-(N-morpholino)]benzyl,  
20 4-[-OCH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]benzyl,  
4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>]benzyl,  
4-[CH<sub>3</sub>(CH<sub>2</sub>)<sub>4</sub>NH-]benzyl,  
4-[N-*n*-butyl,N-*n*-pentylamino-]benzyl,  
4-[-NHC(O)-4'-piperidiny]benzyl,  
25 4-[-NHC(O)CH(NHBoc)(CH<sub>2</sub>)<sub>4</sub>NHCbz]benzyl,  
4-[-NHC(O)-(1',2',3',4'-tetrahydro-N-Boc-isoquinolin-1'-yl)]benzyl,  
*p*-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-1'-(4'-methyl)-piperazinyl]benzyl,  
-(CH<sub>2</sub>)<sub>4</sub>NH-Boc,  
3-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>]benzyl,  
30 4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>]benzyl,

- 3-[-OCH<sub>2</sub>CH<sub>2</sub>-(1'-pyrrolidiny)]benzyl,  
4-[-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)benzyl]benzyl,  
4-[-NHC(S)NHCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-(N-morpholino)]benzyl,  
4-[-OCH<sub>2</sub>CH<sub>2</sub>-(N-morpholino)]benzyl,  
5 4-[-NHCH<sub>2</sub>-(4'-chlorophenyl)]benzyl,  
4-[-NHC(O)NH-(4'-cyanophenyl)]benzyl,  
4-[-OCH<sub>2</sub>COOH]benzyl,  
4-[-OCH<sub>2</sub>COO-*t*-butyl]benzyl,  
4-[-NHC(O)-5'-fluoroindol-2-yl]benzyl,  
10 4-[-NHC(S)NH(CH<sub>2</sub>)<sub>2</sub>-1-piperidiny]]benzyl,  
4-[-N(SO<sub>2</sub>CH<sub>3</sub>)(CH<sub>2</sub>)<sub>3</sub>-N(CH<sub>3</sub>)<sub>2</sub>]benzyl,  
4-[-NHC(O)CH<sub>2</sub>CH(C(O)OCH<sub>2</sub>φ)-NHCbz]benzyl,  
4-[-NHS(O)<sub>2</sub>CF<sub>3</sub>]benzyl,  
3-[-O-(N-methylpiperidin-4'-yl)]benzyl,  
15 4-[-C(=NH)NH<sub>2</sub>]benzyl,  
4-[-NHSO<sub>2</sub>-CH<sub>2</sub>Cl]benzyl,  
4-[-NHC(O)-(1',2',3',4'-tetrahydroisoquinolin-2'-yl)]benzyl,  
4-[-NHC(S)NH(CH<sub>2</sub>)<sub>3</sub>-N-morpholino]benzyl,  
4-[-NHC(O)CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)NHBoc]benzyl,  
20 4-[-C(O)NH<sub>2</sub>]benzyl,  
4-[-NHC(O)NH-3'-methoxyphenyl]benzyl,  
4-[-OCH<sub>2</sub>CH<sub>2</sub>-indol-3'-yl]benzyl,  
4-[-OCH<sub>2</sub>C(O)NH-benzyl]benzyl,  
4-[-OCH<sub>2</sub>C(O)O-benzyl]benzyl,  
25 4-[-OCH<sub>2</sub>C(O)OH]benzyl,  
4-[-OCH<sub>2</sub>-2'-(4',5'-dihydro)imidazolyl]benzyl,  
-CH<sub>2</sub>C(O)NHCH<sub>2</sub>-(4-dimethylamino)phenyl,  
-CH<sub>2</sub>C(O)NHCH<sub>2</sub>-(4-dimethylamino)phenyl,  
4-[-NHC(O)-L-2'-pyrrolidiny-N-SO<sub>2</sub>-4'-methylphenyl]benzyl,  
30 4-[-NHC(O)NHCH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>]benzyl,

- 4-aminobenzyl]benzyl,  
4-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-hydroxy-4-(3-methoxypyrrol-2-yl)-  
piperazinyl]benzyl,  
4-[-O-(N-methylpiperidin-4'-yl)]benzyl,  
5 3-methoxybenzyl,  
4-[-NHC(O)-piperidin-3'-yl]benzyl,  
4-[-NHC(O)-pyridin-2'-yl]benzyl,  
4-[-NHCH<sub>2</sub>-(4'-chlorophenyl)]benzyl,  
4-[-NHC(O)-(N-(4'-CH<sub>3</sub>- $\phi$ -SO<sub>2</sub>)-L-pyrrolidin-2'-yl)]benzyl,  
10 4-[-NHC(O)NHCH<sub>2</sub>CH<sub>2</sub>- $\phi$ ]benzyl,  
4-[-OCH<sub>2</sub>C(O)NH<sub>2</sub>]benzyl,  
4-[-OCH<sub>2</sub>C(O)NH-*t*-butyl]benzyl,  
4-[-OCH<sub>2</sub>CH<sub>2</sub>-1-(4-hydroxy-4-phenyl)-piperidinyl]benzyl,  
4-[-NHSO<sub>2</sub>-CH=CH<sub>2</sub>]benzyl,  
15 4-[-NHSO<sub>2</sub>-CH<sub>2</sub>CH<sub>2</sub>Cl]benzyl,  
-CH<sub>2</sub>C(O)NHCH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>,  
4-[(1'-Cbz-piperidin-4'-yl)C(O)NH-]benzyl,  
4-[(1'-Boc-piperidin-4'-yl)C(O)NH-]benzyl,  
4-[(2'-bromophenyl)C(O)NH-]benzyl,  
20 4-[-NHC(O)-pyridin-4'-yl]benzyl,  
4-[(4'-(CH<sub>3</sub>)<sub>2</sub>NC(O)O-phenyl)-C(O)NH-]benzyl,  
4-[-NHC(O)-1'-methylpiperidin-4'-yl-]benzyl,  
4-(dimethylamino)benzyl,  
4-[-NHC(O)-(1'-N-Boc)-piperidin-2'-yl]benzyl,  
25 3-[-NHC(O)-pyridin-4'-yl]benzyl,  
4-[(*tert*-butyl-O(O)CCH<sub>2</sub>-O-benzyl)-NH-]benzyl,  
[BocNHCH<sub>2</sub>C(O)NH-]butyl,  
4-benzylbenzyl,  
2-hydroxyethyl,  
30 4-[(Et)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NHC(S)NH-]benzyl,

- 4-[(1'-Boc-4'-hydroxypyrrolidin-2'-yl)C(O)NH-]benzyl,  
4-[ $\phi$ CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NHC(S)NH-]benzyl,  
4-[(perhydroindolin-2'-yl)C(O)NH-]benzyl,  
2-[4-hydroxy-4-(3-methoxythien-2-yl)piperidin-1-yl]ethyl,  
5 4-[(1'-Boc-perhydroindolin-2'-yl)-C(O)NH-]benzyl,  
4-[*N*-3-methylbutyl-*N*-trifluoromethanesulfonyl]amino]benzyl,  
4-[*N*-vinylsulfonyl]amino]benzyl,  
4-[2-(2-azabicyclo[3.2.2]octan-2-yl)ethyl-O-]benzyl,  
4-[4'-hydroxypyrrolidin-2'-yl)C(O)NH-]benzyl,  
10 4-( $\phi$ NHC(S)NH)benzyl,  
4-(EtNHC(S)NH)benzyl,  
4-( $\phi$ CH<sub>2</sub>NHC(S)NH)benzyl,  
3-[(1'-Boc-piperidin-2'-yl)C(O)NH-]benzyl,  
3-[piperidin-2'-yl-C(O)NH-]benzyl,  
15 4-[(3'-Boc-thiazolidin-4'-yl)C(O)NH-]benzyl,  
4-(pyridin-3'-yl-NHC(S)NH)benzyl,  
4-(CH<sub>3</sub>-NHC(S)NH)benzyl,  
4-(H<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH)benzyl,  
4-(BocHNCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH)benzyl,  
20 4-(pyridin-4'-yl-CH<sub>2</sub>NH)benzyl,  
4-[(*N,N*-di(4-*N,N*-dimethylamino)benzyl)amino]benzyl,  
4-[(1-Cbz-piperidin-4-yl)C(O)NH-]butyl,  
4-[ $\phi$ CH<sub>2</sub>OCH<sub>2</sub>(BocHN)CHC(O)NH]benzyl,  
4-[(piperidin-4'-yl)C(O)NH-]benzyl,  
25 4-[(pyrrolidin-2'-yl)C(O)NH-]benzyl,  
4-(pyridin-3'-yl-C(O)NH)butyl,  
4-(pyridin-4'-yl-C(O)NH)butyl,  
4-(pyridin-3'-yl-C(O)NH)benzyl,  
4-[CH<sub>3</sub>NHCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl,  
30 4-[CH<sub>3</sub>N(Boc)CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C(O)NH-]benzyl,


- 4-(aminomethyl)benzyl,  
4- $[\phi\text{CH}_2\text{OCH}_2(\text{H}_2\text{N})\text{CHC}(\text{O})\text{NH}]$ benzyl,  
4-[(1',4'-di(Boc)piperazin-2'-yl)-C(O)NH-]benzyl,  
4-[(piperazin-2'-yl)-C(O)NH-]benzyl,  
5 4-[(*N*-toluenesulfonylpyrrolidin-2'-yl)C(O)NH-]butyl,  
4-[-NHC(O)-4'-piperidiny]butyl,  
4-[-NHC(O)-1'-*N*-Boc-piperidin-2'-yl]benzyl,  
4-[-NHC(O)-piperidin-2'-yl]benzyl,  
4-[(1'-*N*-Boc-2',3'-dihydroindolin-2'-yl)-C(O)NH]benzyl,  
10 4-(pyridin-3'-yl-CH<sub>2</sub>NH)benzyl,  
4-[(piperidin-1'-yl)C(O)CH<sub>2</sub>-O-]benzyl,  
4-[(CH<sub>3</sub>)<sub>2</sub>CH]<sub>2</sub>NC(O)CH<sub>2</sub>-O-]benzyl,  
4-[HO(O)C(Cbz-NH)CHCH<sub>2</sub>CH<sub>2</sub>-C(O)NH-]benzyl,  
4- $[\phi\text{CH}_2\text{O}(\text{O})\text{C}(\text{Cbz-NH})\text{CHCH}_2\text{CH}_2\text{-C}(\text{O})\text{NH-}]$ benzyl,  
15 4-[-NHC(O)-2'-methoxyphenyl]benzyl,  
4-[(pyrazin-2'-yl)C(O)NH-]benzyl,  
4-[HO(O)C(NH<sub>2</sub>)CHCH<sub>2</sub>CH<sub>2</sub>-C(O)NH-]benzyl,  
4-(2'-formyl-1',2',3',4'-tetrahydroisoquinolin-3'-yl-CH<sub>2</sub>NH-)benzyl,  
*N*-Cbz-NHCH<sub>2</sub>-,  
20 4-[(4'-methylpiperazin-1'-yl)C(O)O-]benzyl,  
4-[CH<sub>3</sub>(*N*-Boc)NCH<sub>2</sub>C(O)NH-]benzyl,  
4-[-NHC(O)-(1',2',3',4'-tetrahydro-*N*-Boc-isoquinolin-3'-yl)-]benzyl,  
4-[CH<sub>3</sub>NHCH<sub>2</sub>C(O)NH-]benzyl,  
(CH<sub>3</sub>)<sub>2</sub>NC(O)CH<sub>2</sub>-,  
25 4-(*N*-methylacetamido)benzyl,  
4-(1',2',3',4'-tetrahydroisoquinolin-3'-yl-CH<sub>2</sub>NH-)benzyl,  
4-[(CH<sub>3</sub>)<sub>2</sub>NHCH<sub>2</sub>C(O)NH-]benzyl,  
(1-toluenesulfonylimidizol-4-yl)methyl,  
4-[(1'-Boc-piperidin-4'-yl)C(O)NH-]benzyl,  
30 4-trifluoromethylbenzyl,

- 4-[(2'-bromophenyl)C(O)NH-]benzyl,  
4-[(CH<sub>3</sub>)<sub>2</sub>NC(O)NH-]benzyl,  
4-[CH<sub>3</sub>OC(O)NH-]benzyl,  
4-[(CH<sub>3</sub>)<sub>2</sub>NC(O)O-]benzyl,  
5 4-[(CH<sub>3</sub>)<sub>2</sub>NC(O)N(CH<sub>3</sub>)-]benzyl,  
4-[CH<sub>3</sub>OC(O)N(CH<sub>3</sub>)-]benzyl,  
4-(*N*-methyltrifluoroacetamido)benzyl,  
4-[(1'-methoxycarbonylpiperidin-4'-yl)C(O)NH-]benzyl,  
4-[(4'-phenylpiperidin-4'-yl)C(O)NH-]benzyl,  
10 4-[(4'-phenyl-1'-Boc-piperidin-4'-yl)-C(O)NH-]benzyl,  
4-[(piperidin-4'-yl)C(O)O-]benzyl, 4-[(1'-methylpiperidin-4'-yl)-  
O-]benzyl,  
4-[(1'-methylpiperidin-4'-yl)C(O)O-]benzyl,  
4-[(4'-methylpiperazin-1'-yl)C(O)NH-]benzyl,  
15 3-[(CH<sub>3</sub>)<sub>2</sub>NC(O)O-]benzyl,  
4-[(4'-phenyl-1'-Boc-piperidin-4'-yl)-C(O)O-]benzyl,  
4-(*N*-toluenesulfonylamino)benzyl,  
4-[(CH<sub>3</sub>)<sub>3</sub>CC(O)NH-]benzyl,  
4-[(morpholin-4'-yl)C(O)NH-]benzyl,  
20 4-[(CH<sub>3</sub>CH<sub>2</sub>)<sub>2</sub>NC(O)NH-]benzyl,  
4-[-C(O)NH-(4'-piperidiny)]benzyl,  
4-[(2'-trifluoromethylphenyl)C(O)NH-]benzyl,  
4-[(2'-methylphenyl)C(O)NH-]benzyl,  
4-[(CH<sub>3</sub>)<sub>2</sub>NS(O)<sub>2</sub>O-]benzyl,  
25 4-[(pyrrolidin-2'-yl)C(O)NH-]benzyl,  
4-[NHC(O)-piperidin-1'-yl]benzyl,  
4-[(thiomorpholin-4'-yl)C(O)NH-]benzyl,  
4-[(thiomorpholin-4'-yl sulfone)-C(O)NH-]benzyl,  
4-[(morpholin-4'-yl)C(O)O-]benzyl,  
30 3-nitro-4-(CH<sub>3</sub>OC(O)CH<sub>2</sub>O-)benzyl,



- (2-benzoxazolinon-6-yl)methyl-,  
(2*H*-1,4-benzoxazin-3(4*H*)-one-7-yl)methyl-,  
4-[(CH<sub>3</sub>)<sub>2</sub>NS(O)<sub>2</sub>NH-]benzyl,  
4-[(CH<sub>3</sub>)<sub>2</sub>NS(O)<sub>2</sub>N(CH<sub>3</sub>)-]benzyl,  
5 4-[(thiomorpholin-4'-yl)C(O)O-]benzyl,  
4-[(thiomorpholin-4'-yl sulfone)-C(O)O-]benzyl,  
4-[(piperidin-1'-yl)C(O)O-]benzyl,  
4-[(pyrrolidin-1'-yl)C(O)O-]benzyl,  
4-[(4'-methylpiperazin-1'-yl)C(O)O-]benzyl,  
10 4-[(2'-methylpyrrolidin-1'-yl)-,  
(pyridin-4-yl)methyl-,  
4-[(piperazin-4'-yl)-C(O)O-]benzyl,  
4-[(1'-Boc-piperazin-4'-yl)-C(O)O-]benzyl,  
4-[(4'-acetyl)piperazin-1'-yl)C(O)O-]benzyl,  
15 *p*-[(4'-methanesulfonylpiperazin-1'-yl)-benzyl,  
3-nitro-4-[(morpholin-4'-yl)-C(O)O-]benzyl,  
4-[(CH<sub>3</sub>)<sub>2</sub>NC(S)]<sub>2</sub>N-}benzyl,  
*N*-Boc-2-aminoethyl-,  
4-[(1,1-dioxothiomorpholin-4-yl)-C(O)O-]benzyl,  
20 4-[(CH<sub>3</sub>)<sub>2</sub>NS(O)<sub>2</sub>-]benzyl,  
4-(imidazolid-2'-one-1'-yl)benzyl,  
4-[(piperidin-1'-yl)C(O)O-]benzyl,  
1-N-benzyl-imidazol-4-yl-CH<sub>2</sub>-,  
3,4-dioxyethylenebenzyl (i.e., 3,4-ethylenedioxybenzyl),  
25 3,4-dioxymethylenebenzyl (i.e., 3,4-methylenedioxybenzyl),  
4-[N(SO<sub>2</sub>)(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>]benzyl,  
4-(3'-formylimidazolid-2'-one-1'-yl)benzyl,  
4-[NHC(O)CH(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>)NHBoc]benzyl,  
[2'-[4"-hydroxy-4"-(3'''-methoxythien-2'''-yl)piperidin-2"-  
30 yl]ethoxy]benzyl, and

-- 68 --



Preferably,  $\text{R}^5$  in the above compounds is selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl. Even more preferably  $\text{R}^5$  is selected from the group consisting of 4-methylphenyl, methyl, benzyl, *n*-butyl, *n*-hexyl, 4-chlorophenyl, 1-naphthyl, 2-naphthyl, 4-methoxyphenyl, phenyl, 2,4,6-trimethylphenyl, 2-(methoxycarbonyl)phenyl, 2-carboxyphenyl, 3,5-dichlorophenyl, 4-trifluoromethylphenyl, 3,4-dichlorophenyl, 3,4-dimethoxyphenyl, 4-( $\text{CH}_3\text{C}(\text{O})\text{NH-}$ )phenyl, 4-trifluoromethoxyphenyl, 4-cyanophenyl, isopropyl, 3,5-di-(trifluoromethyl)phenyl, 4-*t*-butylphenyl, 4-*t*-butoxyphenyl, 4-nitrophenyl, 2-thienyl, 1-N-methyl-3-methyl-5-chloropyrazol-4-yl, phenethyl, 1-N-methylimidazol-4-yl, 4-bromophenyl, 4-amidinophenyl, 4-methylamidinophenyl, 4-[ $\text{CH}_3\text{SC}(=\text{NH})$ ]phenyl, 5-chloro-2-thienyl, 2,5-dichloro-4-thienyl, 1-N-methyl-4-pyrazolyl, 2-thiazolyl, 5-methyl-1,3,4-thiadiazol-2-yl, 4-[ $\text{H}_2\text{NC}(\text{S})$ ]phenyl, 4-aminophenyl, 4-fluorophenyl, 2-fluorophenyl, 3-fluorophenyl, 3,5-difluorophenyl, pyridin-3-yl, pyrimidin-2-yl, 4-(3'-dimethylamino-*n*-propoxy)-phenyl, and 1-methylpyrazol-4-yl.

Preferably,  $\text{R}^{13}$  in the above compounds is selected from hydrogen or  $\text{C}_{1-6}$  alkyl; more preferably, hydrogen or  $\text{C}_{1-3}$  alkyl; and still more preferably, hydrogen or methyl.

In a preferred embodiment,  $\text{R}^{14}$  in the above compounds is preferably hydrogen and  $\text{R}^{15}$  is preferably  $\text{C}_{1-10}$  alkyl or Cy- $\text{C}_{1-10}$  alkyl, wherein alkyl is optionally substituted with one to four substituents selected from phenyl and  $\text{R}^x$ , and Cy is optionally substituted with one to four substituents independently selected from  $\text{R}^y$ , or  $\text{R}^{14}$  and  $\text{R}^{15}$  and the carbon to which they are attached together from a 3-7 membered mono- or bicyclic carbon only

ring. For the purpose of  $R^{15}$ , Cy is preferably aryl, more preferably phenyl. In a preferred embodiment,  $R^{15}$  is phenyl- $C_{1-3}$  alkyl, wherein phenyl is optionally substituted with one or two groups selected from  $R^y$ . Additional preferred embodiments for  $R^{14}$  and  $R^{15}$  are disclosed in International Patent Application Publication No. WO 98/53814, which application is  
5 incorporated herein by reference in its entirety.

In a preferred embodiment of the above compounds,  $R^{16}$  is substituted amino;  $R^{17}$  and/or  $R^{20}$  are hydrogen; and  $R^{18}$  and/or  $R^{21}$  are alkyl, substituted  
10 alkyl, aryl or substituted aryl.

In a preferred embodiment,  $R^{23}$  in the above compounds is hydrogen. Preferably,  $R^{24}$  in the above compounds is  $Ar^1-Ar^2-C_{1-10}$  alkyl wherein  $Ar^1$  and  $Ar^2$  are optionally substituted with from 1 to 4 groups independently  
15 selected from  $R^b$  and  $R^{25}$  is hydrogen. More preferably,  $R^{24}$  is  $Ar^1-Ar^2-C_{1-3}$  alkyl wherein  $Ar^1$  and  $Ar^2$  are optionally substituted with from 1 to 4 groups independently selected from  $R^b$ ; still more preferably,  $R^{24}$  is  $-CH_2-Ar^2-Ar^1$  and  $R^{25}$  is hydrogen. Additional preferred embodiments are disclosed in International Patent Application Publication No. WO 98/53817, which  
20 application is incorporated herein by reference in its entirety.

Preferably,  $R^3$  and  $R^{3'}$ , or  $R^{14}$  and  $R^{15}$ , or  $R^{24}$  and  $R^{25}$  are derived from L-amino acids or other similarly configured starting materials. Alternatively, racemic mixtures can be used.  
25

Preferably,  $x$  in the above compounds is an integer from 1 to 4.

This invention also provides methods for binding VLA-4 in a biological sample which method comprises contacting the biological sample

-- 70 --


with a compound of this invention under conditions wherein said compound binds to VLA-4.

5       The pharmaceutical compositions may be used to treat disease conditions mediated by VLA-4 or leucocyte adhesion. Such disease conditions include, by way of example, asthma, Alzheimer's disease, atherosclerosis, AIDS dementia, diabetes (including acute juvenile onset diabetes), inflammatory bowel disease (including ulcerative colitis and Crohn's disease), multiple sclerosis, rheumatoid arthritis, tissue  
10       transplantation, tumor metastasis, meningitis, encephalitis, stroke, and other cerebral traumas, nephritis, retinitis, atopic dermatitis, psoriasis, myocardial ischemia and acute leukocyte-mediated lung injury such as that which occurs in adult respiratory distress syndrome.

15       Other disease conditions include, but are not limited to, inflammatory conditions such as erythema nodosum, allergic conjunctivitis, optic neuritis, uveitis, allergic rhinitis, ankylosing spondylitis, psoriatic arthritis, vasculitis, Reiter's syndrome, systemic lupus erythematosus, progressive systemic sclerosis, polymyositis, dermatomyositis, Wegner's granulomatosis, aortitis,  
20       sarcoidosis, lymphocytopenia, temporal arteritis, pericarditis, myocarditis, congestive heart failure, polyarteritis nodosa, hypersensitivity syndromes, allergy, hypereosinophilic syndromes, Churg-Strauss syndrome, chronic obstructive pulmonary disease, hypersensitivity pneumonitis, chronic active hepatitis, interstitial cystitis, autoimmune endocrine failure, primary biliary  
25       cirrhosis, autoimmune aplastic anemia, chronic persistent hepatitis and thyroiditis.

Preferred compounds of this invention include those set forth in the Tables below:

Table I




R <sup>5</sup>	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>9</sup>	X
4-CH <sub>3</sub> -Ph-	H-	H-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OC(CH <sub>3</sub> ) <sub>3</sub>
4-CH <sub>3</sub> -Ph-	H-	H-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
4-CH <sub>3</sub> -Ph-	CH <sub>3</sub> -	H-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OC(CH <sub>3</sub> ) <sub>3</sub>
4-CH <sub>3</sub> -Ph-	CH <sub>3</sub> -	H-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
4-CH <sub>3</sub> -Ph-	4-CH <sub>3</sub> -Ph-	H-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
1-CH <sub>3</sub> - pyrazol-4-yl-	CH <sub>3</sub> -	H-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
4-CH <sub>3</sub> -Ph-	CH <sub>3</sub> -	H-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OCH(CH <sub>3</sub> ) <sub>2</sub>
3-pyridyl-	CH <sub>3</sub> -	H-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OC(CH <sub>3</sub> ) <sub>3</sub>
1-( <i>n</i> -C <sub>4</sub> H <sub>9</sub> )- pyrazol-4-yl-	CH <sub>3</sub> -	H-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OC(CH <sub>3</sub> ) <sub>3</sub>
4-CH <sub>3</sub> -Ph-	CH <sub>3</sub> -	H-	H-	H-	-OH
1-( <i>n</i> -C <sub>4</sub> H <sub>9</sub> )- pyrazol-4-yl-	CH <sub>3</sub> -	H-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
3-pyridyl-	CH <sub>3</sub> -	H-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
4-CH <sub>3</sub> -Ph-	CH <sub>3</sub> -	(CH <sub>3</sub> ) <sub>2</sub> N-	H-	H-	-OH
1-CH <sub>3</sub> - pyrazol-4-yl-	CH <sub>3</sub> -	H-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OCH(CH <sub>3</sub> ) <sub>2</sub>

-- 72 --

R <sup>5</sup>	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>9</sup>	X
3-pyridyl-	CH <sub>3</sub> -	H-	H-	4-(1-CH <sub>3</sub> - piperazin-4- yl)C(O)O-	-OCH(CH <sub>3</sub> ) <sub>2</sub>
3-pyridyl-	CH <sub>3</sub> -	H-	H-	4-(1-CH <sub>3</sub> - piperazin-4- yl)C(O)O-	-OC(CH <sub>3</sub> ) <sub>3</sub>
3-pyridyl-	CH <sub>3</sub> -	H-	H-	4-(1-CH <sub>3</sub> - piperazin-4-yl)- C(O)O-	-OH

Ph = phenyl

Table II



R <sup>16'</sup>	R <sup>20'</sup>	R <sup>18'</sup>	R <sup>19</sup>	X
Cl-	H-	NO <sub>2</sub> -	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	PhCH <sub>2</sub> O-	H-	-OH
H-	H-	PhCH <sub>2</sub> O-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H	H	3-NO <sub>2</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH

R <sup>16'</sup>	R <sup>20'</sup>	R <sup>18'</sup>	R <sup>19</sup>	X
H-	H-	3-pyridyl-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	2-PhCH <sub>2</sub> CH <sub>2</sub> -	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	(CH <sub>3</sub> ) <sub>2</sub> NC(O)- (CH <sub>2</sub> ) <sub>2</sub> -	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	Ph-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	2-CF <sub>3</sub> -Ph-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	2- HOCH <sub>2</sub> Ph-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	CF <sub>3</sub> CH <sub>2</sub> -	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	PhCH <sub>2</sub> -	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OCH(CH <sub>3</sub> ) <sub>2</sub>
H-	H-	2-PhCH <sub>2</sub> CH <sub>2</sub> -	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OCH(CH <sub>3</sub> ) <sub>2</sub>
H-	H-	2-PhCH <sub>2</sub> CH <sub>2</sub> -	H-	-OCH(CH <sub>3</sub> ) <sub>2</sub>
cyclohexyl- (CH <sub>3</sub> )N-	H-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> -	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	2-CH <sub>3</sub> O-Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	2-F-Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
(CH <sub>3</sub> ) <sub>2</sub> CH- (CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
(CH <sub>3</sub> ) <sub>2</sub> CH-NH-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
(CH <sub>3</sub> ) <sub>2</sub> CHCH <sub>2</sub> - (CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> - (CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
(CH <sub>3</sub> ) <sub>2</sub> N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
cyclohexyl- (CH <sub>3</sub> )N-	H-	3-pyridyl-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH

R <sup>16'</sup>	R <sup>20'</sup>	R <sup>18'</sup>	R <sup>19</sup>	X
H-	H-	2-PhCF <sub>2</sub> CH <sub>2</sub> -	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	Cl-	2-PhCF <sub>2</sub> CH <sub>2</sub> -	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
(HOCH <sub>2</sub> CH <sub>2</sub> ) <sub>2</sub> N-	H-	H-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
(HOCH <sub>2</sub> CH <sub>2</sub> ) <sub>2</sub> N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
Ph(CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
(CH <sub>3</sub> ) <sub>2</sub> CHO-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
(CH <sub>3</sub> ) <sub>2</sub> CHCH <sub>2</sub> - CH <sub>2</sub> (CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
CH <sub>3</sub> NH-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
2-CH <sub>3</sub> -Ph-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
HOCH <sub>2</sub> CH <sub>2</sub> - (CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
cyclohexyl-NH-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
1-CH <sub>3</sub> -piperidin- 4-yl-(CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
(CH <sub>3</sub> ) <sub>2</sub> CH- (CH <sub>3</sub> CH <sub>2</sub> -)N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	2,4,6-tri-CH <sub>3</sub> - Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	(CH <sub>3</sub> ) <sub>2</sub> CH-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub> - (CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> - (CH <sub>3</sub> CH <sub>2</sub> -)N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
(CH <sub>3</sub> CH <sub>2</sub> ) <sub>2</sub> N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
CH <sub>3</sub> CH <sub>2</sub> - (CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	cyclohexyl-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
(furan-2-yl)CH <sub>2</sub> - (CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH




R <sup>16'</sup>	R <sup>20'</sup>	R <sup>18'</sup>	R <sup>19</sup>	X
4-Cl-Ph-(CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	thien-3-yl-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	thien-2-yl-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
HOCH <sub>2</sub> CH <sub>2</sub> -(CH <sub>3</sub> )N-	H-	2-F-Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	piperidin-1-yl-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	(CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> ) <sub>2</sub> -CH-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
cyclobutyl-(CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	2-HOCH <sub>2</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	2,6-di-F-Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	2,4-di-CH <sub>3</sub> O-pyrimidin-5-yl	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
cyclohexyl-(CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	2-CF <sub>3</sub> -Ph-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
cyclohexyl-(CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> O-Ph-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
(CH <sub>3</sub> ) <sub>2</sub> CH-(CH <sub>3</sub> )N-	H-	2-F-Ph-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
(CH <sub>3</sub> ) <sub>2</sub> CH-(CH <sub>3</sub> )N-	H-	2-F-Ph-	2-CH <sub>3</sub> O-Ph-	-OH
cyclohexyl-(CH <sub>3</sub> )N-	H-	2,6-di-F-Ph-	2,6-di-F-Ph-	-OH
cyclohexyl-(CH <sub>3</sub> )N-	H-	2-HOCH <sub>2</sub> -Ph-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
(HOCH <sub>2</sub> CH <sub>2</sub> ) <sub>2</sub> N-	H-	2,4,6-tri-CH <sub>3</sub> -Ph-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
cyclohexyl-(CH <sub>3</sub> )N-	H-	2-CF <sub>3</sub> -Ph-	2-NC-Ph-	-OH

R <sup>16'</sup>	R <sup>20'</sup>	R <sup>18'</sup>	R <sup>19</sup>	X
cyclohexyl-(CH <sub>3</sub> )N-	H-	thien-3-yl-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
cyclohexyl-(CH <sub>3</sub> )N-	H-	thien-2-yl-	4-CF <sub>3</sub> -Ph-	-OH
cyclohexyl-(CH <sub>3</sub> )N-	H-	3-pyridyl-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
cyclohexyl-(CH <sub>3</sub> )N-	H-	3-NO <sub>2</sub> -Ph-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
cyclohexyl-(CH <sub>3</sub> )N-	H-	2,6-di-Cl-Ph-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
cyclohexyl-(CH <sub>3</sub> )N-	H-	4-pyridyl-	3-HOCH <sub>2</sub> -Ph-	-OH
(CH <sub>3</sub> ) <sub>2</sub> CH-(CH <sub>3</sub> CH <sub>2</sub> -)N-	H-	2,6-di-CH <sub>3</sub> O-Ph-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
cyclohexyl-(CH <sub>3</sub> )N-	H-	2,3-di-Cl-Ph-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
CH <sub>3</sub> CH <sub>2</sub> -(CH <sub>3</sub> )N-	H-	2,4,6-tri-CH <sub>3</sub> -Ph-	2-NC-Ph-	-OH
(CH <sub>3</sub> ) <sub>2</sub> CH-(CH <sub>3</sub> )N-	H-	2,4,6-tri-CH <sub>3</sub> -Ph-	3-pyridyl-	-OH
(HOCH <sub>2</sub> CH <sub>2</sub> ) <sub>2</sub> N-	H-	2,4,6-tri-CH <sub>3</sub> -Ph-	2-NC-Ph-	-OH
1-CH <sub>3</sub> -piperidin-4-yl-(CH <sub>3</sub> )N-	H-	2-NC-Ph-	2,6-di-F-Ph-	-OH
(CH <sub>3</sub> ) <sub>2</sub> CH-(CH <sub>3</sub> CH <sub>2</sub> -)N-	H-	2,4,6-tri-CH <sub>3</sub> -Ph-	2-CH <sub>3</sub> -Ph-	-OH
4-Cl-Ph-(CH <sub>3</sub> )N-	H-	2,4,6-tri-CH <sub>3</sub> -Ph-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
H-	H-	PhCH <sub>2</sub> CH <sub>2</sub> -(CH <sub>3</sub> )N-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> -(CH <sub>3</sub> )N-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH

R <sup>16'</sup>	R <sup>20'</sup>	R <sup>18'</sup>	R <sup>19</sup>	X
H-	H-	(CH <sub>3</sub> ) <sub>2</sub> CH-(CH <sub>3</sub> )N-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	(CH <sub>3</sub> ) <sub>3</sub> C-(CH <sub>3</sub> )N-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	(CH <sub>3</sub> ) <sub>2</sub> CH-(CH <sub>3</sub> CH <sub>2</sub> -)N-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	4-pyridyl-CH <sub>2</sub> CH <sub>2</sub> -(CH <sub>3</sub> )N-	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	PhCH <sub>2</sub> CH <sub>2</sub> -(CH <sub>3</sub> )N-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
H-	H-	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> -(CH <sub>3</sub> )N-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
H-	H-	(CH <sub>3</sub> ) <sub>2</sub> CH-(CH <sub>3</sub> )N-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
H-	H-	(CH <sub>3</sub> ) <sub>3</sub> C-(CH <sub>3</sub> )N-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
H-	H-	(CH <sub>3</sub> ) <sub>2</sub> CH-(CH <sub>3</sub> CH <sub>2</sub> -)N-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
H-	H-	4-pyridyl-CH <sub>2</sub> CH <sub>2</sub> -(CH <sub>3</sub> )N-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
cyclohexyl-(CH <sub>3</sub> )N-	H-	CH <sub>3</sub> CH <sub>2</sub> -	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
H-	H-	CF <sub>3</sub> CH <sub>2</sub> -	2,6-di-CH <sub>3</sub> O-Ph-	-OH
cyclohexyl-(CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
H-	H-	2-F-Ph-	2,6-di-CH <sub>3</sub> O-Ph-	-OH
CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> -(CH <sub>3</sub> )N-	H-	2-CH <sub>3</sub> -Ph-	2,6-di-CH <sub>3</sub> O-Ph-	-OH

Ph = phenyl


Table III



$R^5$	$R^6$	$R^{7'}$	$R^{8'}$	$R^{9'}$	X
4- $CH_3$ -Ph-	$CH_3$ -	H-	H-	4- $(CH_3)_2NC(O)O$ -	-OH
4- $CH_3$ -Ph-	$CH_3$ -	H-	H-	4- $(CH_3)_2NC(O)O$ -	-OCH $(CH_3)_2$

Ph = phenyl

Table IV



R <sup>5</sup>	R <sup>6</sup>	b	R <sup>9'</sup>	X
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> -	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> -	2	4-HO-	-OH
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> -	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> -	2	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
CH <sub>3</sub> -	CH <sub>3</sub> -	1	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OC(CH <sub>3</sub> ) <sub>3</sub>
3-CH <sub>3</sub> -PhNH- C(O)NH(CH <sub>2</sub> ) <sub>2</sub> -	H-	2	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)O-	-OH
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> -	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> -	2	4-(1-CH <sub>3</sub> - piperazin-4-yl)C(O)O-	-OH

Ph = phenyl

Accordingly, this invention is also directed to each of the following compounds:

- 5            *N*-(2-chloro-5-nitropyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- N*-[5-(*N*-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester,
- 10           *N*-[5-(*N*-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- N*-[5-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester,
- 15           *N*-[5-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- N*-[5-(*N,N*-di-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- 20           *N*-[5-[*N*-(1-*N'*-methylpyrazol-4-ylsulfonyl)-*N*-methylamino]pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- N*-[5-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine isopropyl ester,
- 25           *N*-[5-(*N*-methyl-*N*-3-pyridylsulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester,
- 30           *N*-(5-(*N*-methyl-*N*-(1-butylpyrazol-4-yl)sulfonylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- N*-(5-(2,4-dimethoxypyrimidin-5-yl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- 35           *N*-(5-(2,6-difluorophenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- N*-(5-(2-hydroxymethylphenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- 40           *N*-(2-(*N*-cyclohexylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- 45

- N*-(2-(*N*-methyl-*N*-(1-methylpiperidin-4-yl)amino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 5 *N*-(2-(*N*-ethyl-*N*-isopropylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 10 *N*-(5-isopropylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-butylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 15 *N*-(2-(*N*-ethyl-*N*-propylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N,N*-diethylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 20 *N*-(2-(*N*-methyl-*N*-ethylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-benzyloxypyrimidin-4-yl)-L-phenylalanine,
- 25 *N*-(5-benzyloxypyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl)-L-phenylalanine,
- 30 *N*-(5-(*N*-methyl-*N*-3-pyridinesulfonylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 35 *N*-(5-phenylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(3-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrazin-2-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 40 *N*-(5-(2,2,2-trifluoroethyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,

- N*-(5-(*N*-methyl-*N*-3-pyridinesulfonylamino)pyrimidin-4-yl)-*L*-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine isopropyl ester,
- 5 *N*-(5-benzylpyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(*N*-methyl-*N*-3-pyridinesulfonylamino)pyrimidin-4-yl)-*L*-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine tert-butyl ester,
- 10 *N*-(5-(2-trifluoromethylphenyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(2-*N,N*-dimethylcarbamylethyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 15 *N*-(5-(*N*-methyl-*N*-3-(1-methylpyrazole)sulfonylamino)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine isopropyl ester,
- 20 *N*-(6-phenylpyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(6-(2-trifluoromethylphenyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 25 *N*-(6-(2-hydroxymethylphenyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-cyclohexylpyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 30 *N*-(2-(*N*-methyl-*N*-2-furanmethylamino)-5-(2-tolyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 35 *N*-(2-(*N*-methyl-*N*-4-chlorophenylamino)-5-(2-tolyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(3-thienyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 40 *N*-(5-(2-thienyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-2-hydroxyethylamino)-5-(2-fluorophenyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 45



- N*-(5-(piperidin-1-yl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 5 *N*-(5-(1-propylbutyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-cyclobutylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 10 *N*-(2-(*N,N*-bis-(2-hydroxyethyl)amino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 15 *N*-(2-(*N,N*-bis-(2-hydroxyethyl)amino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-phenylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 20 *N*-(2-(isopropoxy)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-3-methylbutylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 25 *N*-(2-(*N*-methylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(2-tolyl)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 30 *N*-(2-(*N*-methyl-*N*-2-hydroxyethylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-2-methylpropylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 35 *N*-(2-(*N*-methyl-*N*-propylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 40 *N*-(2-(*N,N*-dimethylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(3-pyridyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 45

- N*-(5-(2-phenyl-2,2-difluoroethyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 5 *N*-(5-(2-phenyl-2,2-difluoroethyl)-6-chloropyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(2-phenylethyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 10 *N*-(2-(*N*-methyl-*N*-cyclohexylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-propylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 15 *N*-(5-(2-methoxyphenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(2-fluorophenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 20 *N*-(2-(*N*-Methyl-*N*-isopropylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N*-isopropylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 25 *N*-(5-(2-phenylethyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine isopropyl ester,
- 30 *N*-(3-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrazin-2-yl)-L-phenylalanine isopropyl ester,
- N*-(5-(2-phenylethyl)pyrimidin-4-yl)-L-phenylalanine isopropyl ester,
- 35 *N*-(5-(*N*-methyl-*N*-3-pyridinesulfonylamino)pyrimidin-4-yl)-L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 40 *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-ethylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,

- N*-(5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine isopropyl ester,
- 5 *N*-(5-(3-nitrophenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(3-pyridyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 10 *N*-(5-(2-phenylethyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-*N,N*-dimethylamino-5-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl)-L-phenylalanine,
- 15 *N*-(5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2-methoxyphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- 20 *N*-(2-(*N*-methyl-*N*-isopropylamino)-5-(2-fluorophenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- 25 *N*-(2-(*N*-methyl-*N*-isopropylamino)-5-(2-fluorophenyl)pyrimidin-4-yl)-L-4-(2-methoxyphenyl)phenylalanine,
- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2,6-difluorophenyl)pyrimidin-4-yl)-L-4-(2,6-difluorophenyl)phenylalanine,
- 30 *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2-hydroxymethylphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- 35 *N*-(2-(*N,N*-bis-(2-hydroxyethyl)amino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2-trifluoromethylphenyl)pyrimidin-4-yl)-L-4-(2-cyanophenyl)phenylalanine,
- 40 *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(3-thienyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- 45

- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2-thienyl)pyrimidin-4-yl)-L-4-(4-trifluoromethylphenyl)phenylalanine,
- 5 *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(3-pyridyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(3-nitrophenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- 10 *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2,6-dichlorophenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(4-pyridyl)pyrimidin-4-yl)-L-4-(3-hydroxymethylphenyl)phenylalanine,
- 15 *N*-(2-(*N*-ethyl-*N*-isopropylamino)-5-(2,6-dimethoxyphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2,3-dichlorophenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- 20 *N*-(2-(*N*-methyl-*N*-ethylamino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(2-cyanophenyl)phenylalanine,
- N*-(2-(*N*-methyl-*N*-isopropylamino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(3-pyridyl)phenylalanine,
- 25 *N*-(2-(*N,N*-bis-(2-hydroxyethyl)amino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(2-cyanophenyl)phenylalanine,
- 30 *N*-(2-(*N*-methyl-*N*-(1-methylpiperidin-4-yl)amino)-5-(2-cyanophenyl)pyrimidin-4-yl)-L-4-(2,6-difluorophenyl)phenylalanine,
- N*-(2-(*N*-ethyl-*N*-isopropylamino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(*o*-tolyl)phenylalanine,
- 35 *N*-(2-(*N*-methyl-*N*-4-chlorophenylamino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- 40 *N*-(5-(*N*-methyl-*N*-2-(phenyl)ethylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(*N*-methyl-*N*-hexylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 45

- N*-(5-(*N*-methyl-*N*-isopropylamino)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 5 *N*-(5-(*N*-methyl-*N-tert*-butylamino)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(*N*-ethyl-*N*-isopropylamino)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 10 *N*-(5-(*N*-methyl-*N*-2-(4-pyridyl)ethyl-pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(*N*-methyl-*N*-2-(phenyl)ethylamino)pyrimidin-4-yl)-*L*-4-(4-(2,6-dimethoxyphenyl)phenylalanine,
- 15 *N*-(5-(*N*-methyl-*N*-hexylamino)pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,
- N*-(5-(*N*-methyl-*N*-isopropylamino)pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,
- 20 *N*-(5-(*N*-methyl-*N-tert*-butylamino)pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,
- N*-(5-(*N*-ethyl-*N*-isopropylamino)pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,
- 25 *N*-(5-(*N*-methyl-*N*-2-(4-pyridyl)ethyl-pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,
- 30 *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-ethylpyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(4-(*N,N*-di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-*L*-tyrosine,
- 35 *N*-(4-(*N,N*-di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(4-(*N,N*-dimethylamino)-1-oxo-1,2,5-thiadiazol-3-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester,
- 40 *N*-[4-(2-(3-methylphenylaminocarbonylamino)ethyl-ylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl]-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine
- 45

-- 88 --

N-(4-(N,N-di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine,

5 N-(5-(2,2,2-trifluoroethyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,

N-(2-(N-cyclohexyl-N-methyl)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,

10 N-(5-(2-fluorophenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,

15 N-(2-(N-methyl-N-propyl)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,

N-(3-chloropyrazin-2-yl)-L-4-[1-(*tert*-butoxycarbonyl)piperidin-4-ylcarbonylamino]phenylalanine ethyl ester,

20 and pharmaceutically acceptable salts thereof.

### DETAILED DESCRIPTION OF THE INVENTION

As above, this invention relates to compounds which inhibit leukocyte adhesion and, in particular, leukocyte adhesion mediated by VLA-4.

25 However, prior to describing this invention in further detail, the following terms will first be defined.

#### Definitions

30 As used herein, "alkyl" refers to alkyl groups preferably having from 1 to 10 carbon atoms and more preferably 1 to 6 carbon atoms. This term is exemplified by groups such as methyl, *t*-butyl, *n*-heptyl, octyl and the like.

35 "Substituted alkyl" refers to an alkyl group, preferably of from 1 to 10 carbon atoms, having from 1 to 5 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl,

substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted  
aryloxyaryl, cyano, halogen, hydroxyl, nitro, carboxyl, carboxylalkyl,  
carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted  
cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl,  
5 carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted  
heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone,  
thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl,  
thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted  
thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl,  
10 substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy,  
substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy,  
heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino,  
oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-  
aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted  
15 heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-  
NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted  
alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl,  
-NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-  
substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl,  
20 -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl,  
-NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-  
NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-  
alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino,  
mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono-  
25 and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono-  
and di-substituted heterocyclic amino, unsymmetric di-substituted amines  
having different substituents selected from alkyl, substituted alkyl, aryl,  
substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and  
substituted heterocyclic and substituted alkyl groups having amino groups  
30 blocked by conventional blocking groups such as Boc, Cbz, formyl, and the

-- 90 --

like or alkyl/substituted alkyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl.

"Alkoxy" refers to the group "alkyl-O-" which includes, by way of example, methoxy, ethoxy, *n*-propoxy, *iso*-propoxy, *n*-butoxy, *tert*-butoxy, *sec*-butoxy, *n*-pentoxy, *n*-hexoxy, 1,2-dimethylbutoxy, and the like.

10

"Substituted alkoxy" refers to the group "substituted alkyl-O-".

"Alkenoxy" refers to the group "alkenyl-O-".

15

"Substituted alkenoxy" refers to the group "substituted alkenyl-O-".

"Acyl" refers to the groups H-C(O)-, alkyl-C(O)-, substituted alkyl-C(O)-, alkenyl-C(O)-, substituted alkenyl-C(O)-, alkynyl-C(O)-, substituted alkynyl-C(O)-, cycloalkyl-C(O)-, substituted cycloalkyl-C(O)-, aryl-C(O)-, substituted aryl-C(O)-, heteroaryl-C(O)-, substituted heteroaryl-C(O)-, heterocyclic-C(O)-, and substituted heterocyclic-C(O)- wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

20

25

"Acylamino" refers to the group -C(O)NRR where each R is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl,

30



substituted heteroaryl, heterocyclic, substituted heterocyclic and where each R is joined to form together with the nitrogen atom a heterocyclic or substituted heterocyclic ring wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Thiocarbonylamino" refers to the group -C(S)NRR where each R is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and where each R is joined to form, together with the nitrogen atom a heterocyclic or substituted heterocyclic ring wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Acyloxy" refers to the groups alkyl-C(O)O-, substituted alkyl-C(O)O-, alkenyl-C(O)O-, substituted alkenyl-C(O)O-, alkynyl-C(O)O-, substituted alkynyl-C(O)O-, aryl-C(O)O-, substituted aryl-C(O)O-, cycloalkyl-C(O)O-, substituted cycloalkyl-C(O)O-, heteroaryl-C(O)O-, substituted heteroaryl-C(O)O-, heterocyclic-C(O)O-, and substituted heterocyclic-C(O)O- wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Oxysulfonyl" refers to the groups alkyl-SO<sub>2</sub>O-, substituted alkyl-SO<sub>2</sub>O-, alkenyl-SO<sub>2</sub>O-, substituted alkenyl-SO<sub>2</sub>O-, alkynyl-SO<sub>2</sub>O-, substituted

-- 92 --

alkynyl-SO<sub>2</sub>O-, aryl-SO<sub>2</sub>O-, substituted aryl-SO<sub>2</sub>O-, cycloalkyl-SO<sub>2</sub>O-, substituted cycloalkyl-SO<sub>2</sub>O-, heteroaryl-SO<sub>2</sub>O-, substituted heteroaryl-SO<sub>2</sub>O-, heterocyclic-SO<sub>2</sub>O-, and substituted heterocyclic-SO<sub>2</sub>O- wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

10 "Alkenyl" refers to alkenyl group preferably having from 2 to 10 carbon atoms and more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-2 sites of alkenyl unsaturation.

15 "Substituted alkenyl" refers to alkenyl groups having from 1 to 5 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro, 20 carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, 25 substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, 30 oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-

substituted alkyl,  $-\text{OS}(\text{O})_2\text{-aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  
 $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted}$   
heterocyclic,  $-\text{OSO}_2\text{-NRR}$  where R is hydrogen or alkyl,  $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  
 $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted aryl}$ ,  
5  $-\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-}$   
heterocyclic,  $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-alkyl}$ ,  
 $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted}$   
aryl,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heteroaryl}$ ,  
 $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heterocyclic}$  where R is  
10 hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted  
alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino,  
mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino,  
mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic  
amino, unsymmetric di-substituted amines having different substituents  
15 selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl,  
substituted heteroaryl, heterocyclic and substituted heterocyclic and  
substituted alkenyl groups having amino groups blocked by conventional  
blocking groups such as Boc, Cbz, formyl, and the like or alkenyl/substituted  
alkenyl groups substituted with  $-\text{SO}_2\text{-alkyl}$ ,  $-\text{SO}_2\text{-substituted alkyl}$ ,  $-\text{SO}_2\text{-}$   
20 alkenyl,  $-\text{SO}_2\text{-substituted alkenyl}$ ,  $-\text{SO}_2\text{-cycloalkyl}$ ,  $-\text{SO}_2\text{-substituted}$   
cycloalkyl,  $-\text{SO}_2\text{-aryl}$ ,  $-\text{SO}_2\text{-substituted aryl}$ ,  $-\text{SO}_2\text{-heteroaryl}$ ,  $-\text{SO}_2\text{-substituted}$   
heteroaryl,  $-\text{SO}_2\text{-heterocyclic}$ ,  $-\text{SO}_2\text{-substituted heterocyclic}$  and  $-\text{SO}_2\text{NRR}$   
where R is hydrogen or alkyl.

25 "Alkynyl" refers to alkynyl group preferably having from 2 to 10  
carbon atoms and more preferably 3 to 6 carbon atoms and having at least 1  
and preferably from 1-2 sites of alkynyl unsaturation.

30 "Substituted alkynyl" refers to alkynyl groups having from 1 to 5  
substituents selected from the group consisting of alkoxy, substituted alkoxy,

acyl, acylamino, thiocarbonylamino, acyloxy, amino, amidino, alkylamidino,  
 thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino,  
 aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy,  
 aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro,  
 5 carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl,  
 carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl,  
 carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic,  
 carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl,  
 guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl,  
 10 substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl,  
 substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic,  
 heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic,  
 cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted  
 heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy,  
 15 oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-}$   
 substituted alkyl,  $-\text{OS}(\text{O})_2\text{-aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  
 $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted}$   
 heterocyclic,  $-\text{OSO}_2\text{-NRR}$  where R is hydrogen or alkyl,  $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  
 $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted aryl}$ , -  
 20  $\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heterocyclic}$ ,  
 $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-alkyl}$ ,  
 $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  
 $-\text{NRS}(\text{O})_2\text{-NR-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-}$   
 substituted heteroaryl,  $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted}$   
 25 heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono-  
 and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-  
 substituted arylamino, mono- and di-heteroarylamino, mono- and di-  
 substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-  
 substituted heterocyclic amino, unsymmetric di-substituted amines having  
 30 different substituents selected from alkyl, substituted alkyl, aryl, substituted

aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and substituted alkynyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkynyl/substituted alkynyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl.

10 "Amidino" refers to the group H<sub>2</sub>NC(=NH)- and the term "alkylamidino" refers to compounds having 1 to 3 alkyl groups (e.g., alkylHNC(=NH)-).

15 "Thioamidino" refers to the group RSC(=NH)- where R is hydrogen or alkyl.

"Amino" refers to the group -NH<sub>2</sub>.

20 "Substituted amino" refers to the group -NRR, where each R group is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic, provided that both R groups are not hydrogen; or the R groups can be joined together with the nitrogen atom to form a heterocyclic or substituted heterocyclic ring.

"Aminoacyl" refers to the groups -NRC(O)alkyl, -NRC(O)substituted alkyl, -NRC(O)cycloalkyl, -NRC(O)substituted cycloalkyl, -NRC(O)alkenyl, -NRC(O)substituted alkenyl, -NRC(O)alkynyl, -NRC(O)substituted alkynyl, -NRC(O)aryl, -NRC(O)substituted aryl, -NRC(O)heteroaryl, -NRC(O)substituted heteroaryl, -NRC(O)heterocyclic, and -NRC(O)substituted heterocyclic where R is hydrogen or alkyl and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminosulfonyl" refers to the groups -NRSO<sub>2</sub>alkyl, -NRSO<sub>2</sub>substituted alkyl, -NRSO<sub>2</sub>cycloalkyl, -NRSO<sub>2</sub>substituted cycloalkyl, -NRSO<sub>2</sub>alkenyl, -NRSO<sub>2</sub>substituted alkenyl, -NRSO<sub>2</sub>alkynyl, -NRSO<sub>2</sub>substituted alkynyl, -NRSO<sub>2</sub>aryl, -NRSO<sub>2</sub>substituted aryl, -NRSO<sub>2</sub>heteroaryl, -NRSO<sub>2</sub>substituted heteroaryl, -NRSO<sub>2</sub>heterocyclic, and -NRSO<sub>2</sub>substituted heterocyclic where R is hydrogen or alkyl and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminocarbonyloxy" refers to the groups -NRC(O)O-alkyl, -NRC(O)O-substituted alkyl, -NRC(O)O-alkenyl, -NRC(O)O-substituted alkenyl, -NRC(O)O-alkynyl, -NRC(O)O-substituted alkynyl, -NRC(O)O-cycloalkyl, -NRC(O)O-substituted cycloalkyl, -NRC(O)O-aryl, -NRC(O)O-substituted aryl, -NRC(O)O-heteroaryl, -NRC(O)O-substituted heteroaryl, -NRC(O)O-heterocyclic, and -NRC(O)O-substituted heterocyclic where R is hydrogen or alkyl and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl,

substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminosulfonyloxy" refers to the groups -NRSO<sub>2</sub>O-alkyl,  
5 -NRSO<sub>2</sub>O-substituted alkyl, -NRSO<sub>2</sub>O-alkenyl, -NRSO<sub>2</sub>O-substituted alkenyl,  
-NRSO<sub>2</sub>O-alkynyl, -NRSO<sub>2</sub>O-substituted alkynyl, -NRSO<sub>2</sub>O-cycloalkyl,  
-NRSO<sub>2</sub>O-substituted cycloalkyl, -NRSO<sub>2</sub>O-aryl, -NRSO<sub>2</sub>O-substituted aryl,  
-NRSO<sub>2</sub>O-heteroaryl, -NRSO<sub>2</sub>O-substituted heteroaryl,  
-NRSO<sub>2</sub>O-heterocyclic, and -NRSO<sub>2</sub>O-substituted heterocyclic where R is  
10 hydrogen or alkyl and wherein alkyl, substituted alkyl, alkenyl, substituted  
alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl,  
substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and  
substituted heterocyclic are as defined herein.

15 "Oxycarbonylamino" refers to the groups -OC(O)NH<sub>2</sub>, -OC(O)NRR,  
-OC(O)NR-alkyl, -OC(O)NR-substituted alkyl, -OC(O)NR-alkenyl,  
-OC(O)NR-substituted alkenyl, -OC(O)NR-alkynyl, -OC(O)NR-substituted  
alkynyl, -OC(O)NR-cycloalkyl, -OC(O)NR-substituted cycloalkyl,  
-OC(O)NR-aryl, -OC(O)NR-substituted aryl, -OC(O)NR-heteroaryl,  
20 -OC(O)NR-substituted heteroaryl, -OC(O)NR-heterocyclic, and  
-OC(O)NR-substituted heterocyclic where R is hydrogen, alkyl or where each  
R is joined to form, together with the nitrogen atom a heterocyclic or  
substituted heterocyclic ring and wherein alkyl, substituted alkyl, alkenyl,  
substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted  
25 cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl,  
heterocyclic and substituted heterocyclic are as defined herein.

"Oxythiocarbonylamino" refers to the groups -OC(S)NH<sub>2</sub>,  
-OC(S)NRR, -OC(S)NR-alkyl, -OC(S)NR-substituted alkyl, -OC(S)NR-  
30 alkenyl, -OC(S)NR-substituted alkenyl, -OC(S)NR-alkynyl, -OC(S)NR-

substituted alkynyl, -OC(S)NR-cycloalkyl, -OC(S)NR-substituted cycloalkyl,  
-OC(S)NR-aryl, -OC(S)NR-substituted aryl, -OC(S)NR-heteroaryl,  
-OC(S)NR-substituted heteroaryl, -OC(S)NR-heterocyclic, and  
-OC(S)NR-substituted heterocyclic where R is hydrogen, alkyl or where each  
5 R is joined to form together with the nitrogen atom a heterocyclic or  
substituted heterocyclic ring and wherein alkyl, substituted alkyl, alkenyl,  
substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted  
cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl,  
heterocyclic and substituted heterocyclic are as defined herein.

10  
"Oxysulfonylamino" refers to the groups -OSO<sub>2</sub>NH<sub>2</sub>, -OSO<sub>2</sub>NRR,  
-OSO<sub>2</sub>NR-alkyl, -OSO<sub>2</sub>NR-substituted alkyl, -OSO<sub>2</sub>NR-alkenyl,  
-OSO<sub>2</sub>NR-substituted alkenyl, -OSO<sub>2</sub>NR-alkynyl, -OSO<sub>2</sub>NR-substituted  
alkynyl, -OSO<sub>2</sub>NR-cycloalkyl, -OSO<sub>2</sub>NR-substituted cycloalkyl,  
15 -OSO<sub>2</sub>NR-aryl, -OSO<sub>2</sub>NR-substituted aryl, -OSO<sub>2</sub>NR-heteroaryl,  
-OSO<sub>2</sub>NR-substituted heteroaryl, -OSO<sub>2</sub>NR-heterocyclic, and  
-OSO<sub>2</sub>NR-substituted heterocyclic where R is hydrogen, alkyl or where each  
R is joined to form, together with the nitrogen atom a heterocyclic or  
substituted heterocyclic ring and wherein alkyl, substituted alkyl, alkenyl,  
20 substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted  
cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl,  
heterocyclic and substituted heterocyclic are as defined herein.

25 "Aminocarbonylamino" refers to the groups -NRC(O)NRR,  
-NRC(O)NR-alkyl, -NRC(O)NR-substituted alkyl, -NRC(O)NR-alkenyl,  
-NRC(O)NR-substituted alkenyl, -NRC(O)NR-alkynyl,  
-NRC(O)NR-substituted alkynyl, -NRC(O)NR-aryl, -NRC(O)NR-substituted  
aryl, -NRC(O)NR-cycloalkyl, -NRC(O)NR-substituted cycloalkyl,  
30 -NRC(O)NR-heteroaryl, and -NRC(O)NR-substituted heteroaryl,



-NRC(O)NR-heterocyclic, and -NRC(O)NR-substituted heterocyclic where each R is independently hydrogen, alkyl or where each R is joined to form together with the nitrogen atom a heterocyclic or substituted heterocyclic ring as well as where one of the amino groups is blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

10 "Aminothiocabonylamino" refers to the groups -NRC(S)NRR, -NRC(S)NR-alkyl, -NRC(S)NR-substituted alkyl, -NRC(S)NR-alkenyl, -NRC(S)NR-substituted alkenyl, -NRC(S)NR-alkynyl, -NRC(S)NR-substituted alkynyl, -NRC(S)NR-aryl, -NRC(S)NR-substituted aryl, -NRC(S)NR-cycloalkyl, -NRC(S)NR-substituted cycloalkyl, -NRC(S)NR-heteroaryl, and -NRC(S)NR-substituted heteroaryl, -NRC(S)NR-heterocyclic, and -NRC(S)NR-substituted heterocyclic where each R is independently hydrogen, alkyl or where each R is joined to form together with the nitrogen atom a heterocyclic or substituted heterocyclic ring as well as where one of the amino groups is blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

25 "Aminosulfonylamino" refers to the groups -NRSO<sub>2</sub>NRR, -NRSO<sub>2</sub>NR-alkyl, -NRSO<sub>2</sub>NR-substituted alkyl, -NRSO<sub>2</sub>NR-alkenyl, -NRSO<sub>2</sub>NR-substituted alkenyl, -NRSO<sub>2</sub>NR-alkynyl, -NRSO<sub>2</sub>NR-substituted alkynyl, -NRSO<sub>2</sub>NR-aryl, -NRSO<sub>2</sub>NR-substituted aryl, -NRSO<sub>2</sub>NR-cycloalkyl, -NRSO<sub>2</sub>NR-substituted cycloalkyl,

-NRSO<sub>2</sub>NR-heteroaryl, and -NRSO<sub>2</sub>NR-substituted heteroaryl,  
-NRSO<sub>2</sub>NR-heterocyclic, and -NRSO<sub>2</sub>NR-substituted heterocyclic, where  
each R is independently hydrogen, alkyl or where each R is joined to form  
together with the nitrogen atom a heterocyclic or substituted heterocyclic ring  
5 as well as where one of the amino groups is blocked by conventional blocking  
groups such as Boc, Cbz, formyl, and the like and wherein alkyl, substituted  
alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl,  
substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted  
heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

10

"Aryl" or "Ar" refers to an unsaturated aromatic carbocyclic group of  
from 6 to 14 carbon atoms having a single ring (e.g., phenyl) or multiple  
condensed rings (e.g., naphthyl or anthryl) which condensed rings may or  
may not be aromatic (e.g., 2-benzoxazolinone, 2H-1,4-benzoxazin-3(4H)-  
15 one-7yl, and the like). Preferred aryls include phenyl and naphthyl.

Substituted aryl refers to aryl groups which are substituted with from  
1 to 3 substituents selected from the group consisting of hydroxy, acyl,  
acylamino, thiocarbonylamino, acyloxy, alkyl, substituted alkyl, alkoxy,  
20 substituted alkoxy, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl,  
amidino, alkylamidino, thioamidino, amino, aminoacyl, aminocarbonyloxy,  
aminocarbonylamino, aminothiocarbonylamino, aryl, substituted aryl,  
aryloxy, substituted aryloxy, cycloalkoxy, substituted cycloalkoxy,  
heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted  
25 heterocyclyloxy, carboxyl, carboxylalkyl, carboxyl-substituted alkyl,  
carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-  
substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl,  
carboxylheterocyclic, carboxyl-substituted heterocyclic, carboxylamido,  
cyano, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl,  
30 thioheteroaryl, substituted thioheteroaryl, thiocycloalkyl, substituted

thiocycloalkyl, thioheterocyclic, substituted thioheterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, halo, nitro, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino,  $-S(O)_2$ -alkyl,  $-S(O)_2$ -substituted alkyl,  $-S(O)_2$ -cycloalkyl,  $-S(O)_2$ -substituted cycloalkyl,  $-S(O)_2$ -alkenyl,  $-S(O)_2$ -substituted alkenyl,  $-S(O)_2$ -aryl,  $-S(O)_2$ -substituted aryl,  $-S(O)_2$ -heteroaryl,  $-S(O)_2$ -substituted heteroaryl,  $-S(O)_2$ -heterocyclic,  $-S(O)_2$ -substituted heterocyclic,  $-OS(O)_2$ -alkyl,  $-OS(O)_2$ -substituted alkyl,  $-OS(O)_2$ -aryl,  $-OS(O)_2$ -substituted aryl,  $-OS(O)_2$ -heteroaryl,  $-OS(O)_2$ -substituted heteroaryl,  $-OS(O)_2$ -heterocyclic,  $-OS(O)_2$ -substituted heterocyclic,  $-OSO_2$ -NRR where R is hydrogen or alkyl,  $-NRS(O)_2$ -alkyl,  $-NRS(O)_2$ -substituted alkyl,  $-NRS(O)_2$ -aryl,  $-NRS(O)_2$ -substituted aryl,  $-NRS(O)_2$ -heteroaryl,  $-NRS(O)_2$ -substituted heteroaryl,  $-NRS(O)_2$ -heterocyclic,  $-NRS(O)_2$ -substituted heterocyclic,  $-NRS(O)_2$ -NR-alkyl,  $-NRS(O)_2$ -NR-substituted alkyl,  $-NRS(O)_2$ -NR-aryl,  $-NRS(O)_2$ -NR-substituted aryl,  $-NRS(O)_2$ -NR-heteroaryl,  $-NRS(O)_2$ -NR-substituted heteroaryl,  $-NRS(O)_2$ -NR-heterocyclic,  $-NRS(O)_2$ -NR-substituted heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and amino groups on the substituted aryl blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or substituted with  $-SO_2NRR$  where R is hydrogen or alkyl.

"Aryloxy" refers to the group aryl-O- which includes, by way of example, phenoxy, naphthoxy, and the like.

"Substituted aryloxy" refers to substituted aryl-O- groups.

5

"Aryloxyaryl" refers to the group -aryl-O-aryl.

"Substituted aryloxyaryl" refers to aryloxyaryl groups substituted with from 1 to 3 substituents on either or both aryl rings selected from the group consisting of hydroxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkyl, substituted alkyl, alkoxy, substituted alkoxy, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, amidino, alkylamidino, thioamidino, amino, aminoacyl, aminocarbonyloxy, aminocarbonylamino, aminothiocabonylamino, aryl, substituted aryl, aryloxy, substituted aryloxy, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, carboxylamido, cyano, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thioheteroaryl, substituted thioheteroaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheterocyclic, substituted thioheterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, halo, nitro, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -S(O)<sub>2</sub>-alkyl, -S(O)<sub>2</sub>-substituted alkyl, -S(O)<sub>2</sub>-cycloalkyl, -S(O)<sub>2</sub>-substituted cycloalkyl, -S(O)<sub>2</sub>-alkenyl, -S(O)<sub>2</sub>-substituted alkenyl, -S(O)<sub>2</sub>-aryl, -S(O)<sub>2</sub>-substituted aryl, -S(O)<sub>2</sub>-heteroaryl, -S(O)<sub>2</sub>-substituted heteroaryl, -S(O)<sub>2</sub>-heterocyclic, -S(O)<sub>2</sub>-

30

substituted heterocyclic,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{OSO}_2\text{-NRR}$  where R is hydrogen or alkyl,  $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  
5  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heterocyclic}$  where R is hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino, mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic amino, unsymmetric di-substituted amines  
10 having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and amino groups on the substituted aryl blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or substituted with  $-\text{SO}_2\text{NRR}$  where R is hydrogen or alkyl.

20

"Cycloalkyl" refers to cyclic alkyl groups of from 3 to 8 carbon atoms having a single cyclic ring including, by way of example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclooctyl and the like. Excluded from this definition are multi-ring alkyl groups such as adamantanyl, etc.

25

"Cycloalkenyl" refers to cyclic alkenyl groups of from 3 to 8 carbon atoms having single or multiple unsaturation but which are not aromatic.

"Substituted cycloalkyl" and "substituted cycloalkenyl" refer to a  
30 cycloalkyl and cycloalkenyl groups, preferably of from 3 to 8 carbon atoms,

having from 1 to 5 substituents selected from the group consisting of oxo  
 (=O), thioxo (=S), alkoxy, substituted alkoxy, acyl, acylamino,  
 thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino,  
 aminoacyl, aminocarbonylamino, aminothiocabonylamino,  
 5 aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy,  
 aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro,  
 carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl,  
 carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl,  
 carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic,  
 10 carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl,  
 guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl,  
 substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl,  
 substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic,  
 heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic,  
 15 cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted  
 heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy,  
 oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-  
 substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl,  
 -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted  
 20 heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl,  
 -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-  
 substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl,  
 -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic,  
 -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl,  
 25 -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-  
 substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted  
 heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono-  
 and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-  
 substituted arylamino, mono- and di-heteroarylamino, mono- and di-  
 30 substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-

-- 105 --

substituted heterocyclic amino, unsymmetric di-substituted amines having different substituents selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic and substituted alkynyl groups having amino groups blocked by  
5 conventional blocking groups such as Boc, Cbz, formyl, and the like or alkynyl/substituted alkynyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic  
10 and -SO<sub>2</sub>NRR where R is hydrogen or alkyl.

"Cycloalkoxy" refers to -O-cycloalkyl groups.

15 "Substituted cycloalkoxy" refers to -O-substituted cycloalkyl groups.

"Cycloalkenoxy" refers to -O-cycloalkenyl groups.

20 "Substituted cycloalkenoxy" refers to -O-substituted cycloalkenyl groups.

25 "Guanidino" refers to the groups -NRC(=NR)NRR, -NRC(=NR)NR-alkyl, -NRC(=NR)NR-substituted alkyl, -NRC(=NR)NR-alkenyl, -NRC(=NR)NR-substituted alkenyl, -NRC(=NR)NR-alkynyl, -NRC(=NR)NR-substituted alkynyl, -NRC(=NR)NR-aryl, -NRC(=NR)NR-substituted aryl, -NRC(=NR)NR-cycloalkyl, -NRC(=NR)NR-heteroaryl, -NRC(=NR)NR-substituted heteroaryl, -NRC(=NR)NR-heterocyclic, and -NRC(=NR)NR-substituted heterocyclic where each R is independently hydrogen and alkyl as well as where one of the amino groups is blocked by conventional blocking groups such as Boc,  
30 Cbz, formyl, and the like and wherein alkyl, substituted alkyl, alkenyl,

substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

5           "Guanidinosulfone" refers to the groups -NRC(=NR)NRSO<sub>2</sub>-alkyl, -NRC(=NR)NRSO<sub>2</sub>-substituted alkyl, -NRC(=NR)NRSO<sub>2</sub>-alkenyl, -NRC(=NR)NRSO<sub>2</sub>-substituted alkenyl, -NRC(=NR)NRSO<sub>2</sub>-alkynyl, -NRC(=NR)NRSO<sub>2</sub>-substituted alkynyl, -NRC(=NR)NRSO<sub>2</sub>-aryl, -NRC(=NR)NRSO<sub>2</sub>-substituted aryl, -NRC(=NR)NRSO<sub>2</sub>-cycloalkyl, -NRC(=NR)NRSO<sub>2</sub>-substituted cycloalkyl, -NRC(=NR)NRSO<sub>2</sub>-heteroaryl, 10           and -NRC(=NR)NRSO<sub>2</sub>-substituted heteroaryl, -NRC(=NR)NRSO<sub>2</sub>-heterocyclic, and -NRC(=NR)NRSO<sub>2</sub>-substituted heterocyclic where each R is independently hydrogen and alkyl and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted 15           heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Halo" or "halogen" refers to fluoro, chloro, bromo and iodo and preferably is either chloro or bromo.

20

"Heteroaryl" refers to an aromatic carbocyclic group of from 2 to 10 carbon atoms and 1 to 4 heteroatoms selected from oxygen, nitrogen and sulfur within the ring or oxides thereof. Such heteroaryl groups can have a single ring (e.g., pyridyl or furyl) or multiple condensed rings (e.g., 25           indoliziny or benzothienyl). Additionally, the heteroatoms of the heteroaryl group may be oxidized, i.e., to form pyridine N-oxides or 1,1-dioxo-1,2,5-thiadiazoles and the like. Preferred heteroaryls include pyridyl, pyrrolyl, indolyl, furyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1-oxo-1,2,5-thiadiazolyl and 1,1-dioxo-1,2,5-thiadiazolyl. The term "heteroaryl having two nitrogen 30           atoms in the heteroaryl ring" refers to a heteroaryl group having two, and



only two, nitrogen atoms in the heteroaryl ring and optionally containing 1 or 2 other heteroatoms in the heteroaryl ring, such as oxygen or sulfur

"Substituted heteroaryl" refers to heteroaryl groups which are

5 substituted with from 1 to 3 substituents selected from the group consisting of hydroxy, acyl, acylamino, thiocarbonylamino, acyloxy, alkyl, substituted alkyl, alkoxy, substituted alkoxy, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, amidino, alkylamidino, thioamidino, amino, aminoacyl, aminocarbonyloxy, aminocarbonylamino, aminothiocabonylamino, aryl,

10 substituted aryl, aryloxy, substituted aryloxy, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted

15 heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, carboxylamido, cyano, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thioheteroaryl, substituted thioheteroaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheterocyclic, substituted thioheterocyclic, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, halo, nitro,

20 heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -S(O)<sub>2</sub>-alkyl, -S(O)<sub>2</sub>-substituted alkyl, -S(O)<sub>2</sub>-cycloalkyl, -S(O)<sub>2</sub>-substituted cycloalkyl,

25 -S(O)<sub>2</sub>-alkenyl, -S(O)<sub>2</sub>-substituted alkenyl, -S(O)<sub>2</sub>-aryl, -S(O)<sub>2</sub>-substituted aryl, -S(O)<sub>2</sub>-heteroaryl, -S(O)<sub>2</sub>-substituted heteroaryl, -S(O)<sub>2</sub>-heterocyclic, -S(O)<sub>2</sub>-substituted heterocyclic, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted

30 heterocyclic, -OSO<sub>2</sub>-NRR where R is hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl,

-NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl,  
-NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-  
heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl,  
-NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted  
5 aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl,  
-NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is  
hydrogen or alkyl, mono- and di-alkylamino, mono- and di-(substituted  
alkyl)amino, mono- and di-arylamino, mono- and di-substituted arylamino,  
mono- and di-heteroarylamino, mono- and di-substituted heteroarylamino,  
10 mono- and di-heterocyclic amino, mono- and di-substituted heterocyclic  
amino, unsymmetric di-substituted amines having different substituents  
selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl,  
substituted heteroaryl, heterocyclic and substituted heterocyclic and amino  
groups on the substituted aryl blocked by conventional blocking groups such  
15 as Boc, Cbz, formyl, and the like or substituted with -SO<sub>2</sub>NRR where R is  
hydrogen or alkyl.

"Heteroaryloxy" refers to the group -O-heteroaryl and "substituted  
heteroaryloxy" refers to the group -O-substituted heteroaryl.

20  
"Heterocycle" or "heterocyclic" refers to a saturated or unsaturated  
group having a single ring or multiple condensed rings, from 1 to 10 carbon  
atoms and from 1 to 4 hetero atoms selected from nitrogen, sulfur or oxygen  
within the ring wherein, in fused ring systems, one or more of the rings can  
25 be aryl or heteroaryl.

"Substituted heterocyclic" refers to heterocycle groups which are  
substituted with from 1 to 3 substituents selected from the group consisting  
of oxo (=O), thioxo (=S), alkoxy, substituted alkoxy, acyl, acylamino,  
30 thiocarbonylamino, acyloxy, amino, amidino, alkylamidino, thioamidino,

aminoacyl, aminocarbonylamino, aminothiocarbonylamino,  
aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy,  
aryloxyaryl, substituted aryloxyaryl, halogen, hydroxyl, cyano, nitro,  
carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl,  
5 carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl,  
carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic,  
carboxyl-substituted heterocyclic, cycloalkyl, substituted cycloalkyl,  
guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl,  
substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl,  
10 substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic,  
heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic,  
cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted  
heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy,  
oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-}$   
15 substituted alkyl,  $-\text{OS}(\text{O})_2\text{-aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  
 $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted}$   
heterocyclic,  $-\text{OSO}_2\text{-NRR}$  where R is hydrogen or alkyl,  
 $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-}$   
substituted aryl,  $-\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  
20  $-\text{NRS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  
 $-\text{NRS}(\text{O})_2\text{-NR-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  
 $-\text{NRS}(\text{O})_2\text{-NR-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-}$   
substituted heteroaryl,  $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted}$   
heterocyclic where R is hydrogen or alkyl, mono- and di-alkylamino, mono-  
25 and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-  
substituted arylamino, mono- and di-heteroarylamino, mono- and di-  
substituted heteroarylamino, mono- and di-heterocyclic amino, mono- and di-  
substituted heterocyclic amino, unsymmetric di-substituted amines having  
different substituents selected from alkyl, substituted alkyl, aryl, substituted  
30 aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted

-- 110 --

heterocyclic and substituted alkynyl groups having amino groups blocked by conventional blocking groups such as Boc, Cbz, formyl, and the like or alkynyl/substituted alkynyl groups substituted with -SO<sub>2</sub>-alkyl, -SO<sub>2</sub>-substituted alkyl, -SO<sub>2</sub>-alkenyl, -SO<sub>2</sub>-substituted alkenyl, -SO<sub>2</sub>-cycloalkyl, -SO<sub>2</sub>-substituted cycloalkyl, -SO<sub>2</sub>-aryl, -SO<sub>2</sub>-substituted aryl, -SO<sub>2</sub>-heteroaryl, -SO<sub>2</sub>-substituted heteroaryl, -SO<sub>2</sub>-heterocyclic, -SO<sub>2</sub>-substituted heterocyclic and -SO<sub>2</sub>NRR where R is hydrogen or alkyl.

Examples of heterocycles and heteroaryls include, but are not limited to, azetidine, pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, dihydroindole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine, naphthylpyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, phenanthroline, isothiazole, phenazine, isoxazole, phenoxazine, phenothiazine, imidazolidine, imidazoline, piperidine, piperazine, indoline, phthalimide, 1,2,3,4-tetrahydroisoquinoline, 4,5,6,7-tetrahydrobenzo[b]thiophene, thiazole, thiazolidine, thiophene, benzo[b]thiophene, morpholino, thiomorpholino, piperidiny, pyrrolidine, tetrahydrofuranyl, and the like.

"Heterocycloxy" refers to the group -O-heterocyclic and "substituted heterocycloxy" refers to the group -O-substituted heterocyclic.

"Thiol" refers to the group -SH.

"Thioalkyl" refers to the groups -S-alkyl

"Substituted thioalkyl" refers to the group -S-substituted alkyl.

"Thiocycloalkyl" refers to the groups -S-cycloalkyl.

-- 111 --

"Substituted thiocycloalkyl" refers to the group -S-substituted cycloalkyl.

5 "Thioaryl" refers to the group -S-aryl and "substituted thioaryl" refers to the group -S-substituted aryl.

"Thioheteroaryl" refers to the group -S-heteroaryl and "substituted thioheteroaryl" refers to the group -S-substituted heteroaryl.

10 "Thioheterocyclic" refers to the group -S-heterocyclic and "substituted thioheterocyclic" refers to the group -S-substituted heterocyclic.

"Pharmaceutically acceptable salt" refers to pharmaceutically acceptable salts of a compound of Formula I which salts are derived from a variety of organic and inorganic counter ions well known in the art and include, by way of example only, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium, and the like; and when the molecule contains a basic functionality, salts of organic or inorganic acids, such as hydrochloride, hydrobromide, tartrate, mesylate, acetate, maleate, oxalate and the like.

15  
20

#### Compound Preparation

The compounds of this invention can be prepared from readily available starting materials using the following general methods and procedures. It will be appreciated that where typical or preferred process conditions (i.e., reaction temperatures, times, mole ratios of reactants, solvents, pressures, etc.) are given, other process conditions can also be used unless otherwise stated. Optimum reaction conditions may vary with the particular reactants or solvent used, but such conditions can be determined by one skilled in the art by routine optimization procedures.


25  
30

-- 112 --

Additionally, as will be apparent to those skilled in the art, conventional protecting groups may be necessary to prevent certain functional groups from undergoing undesired reactions. Suitable protecting groups for various functional groups as well as suitable conditions for protecting and deprotecting particular functional groups are well known in the art. For example, numerous protecting groups are described in T. W. Greene and G. M. Wuts, *Protecting Groups in Organic Synthesis*, Second Edition, Wiley, New York, 1991, and references cited therein.

Furthermore, the compounds of this invention will typically contain one or more chiral centers. Accordingly, if desired, such compounds can be prepared or isolated as pure stereoisomers, i.e., as individual enantiomers or diastereomers, or as stereoisomer-enriched mixtures. All such stereoisomers (and enriched mixtures) are included within the scope of this invention, unless otherwise indicated. Pure stereoisomers (or enriched mixtures) may be prepared using, for example, optically active starting materials or stereoselective reagents well-known in the art. Alternatively, racemic mixtures of such compounds can be separated using, for example, chiral column chromatography, chiral resolving agents and the like.


In a preferred method of synthesis, the compounds of this invention are prepared by coupling an amino acid derivative of the formula:



where R<sup>3</sup> and R<sup>3'</sup> are as defined herein and P<sup>1</sup> is a carboxylic acid protecting group (such as an alkyl group, i.e. methyl, ethyl and the like), with a suitably

functionalized heteroaryl or heterocyclic intermediate. For example, such coupling reactions may be performed by displacing a leaving group, such as chloro, bromo, iodo, tosyl and the like, from the heteroaryl or heterocyclic intermediate with the amino group of the amino acid derivative; or by  
5 reductive alkylation of the amino group of amino acid derivative with a carbonyl-functionalized intermediate. Such coupling reactions are well-known to those skilled in the art.

By way of illustration, the synthesis of a representative compound of  
10 formula I is shown in Scheme 1.

Scheme 1

As shown in Scheme 1, 5-nitrouracil, **1**, (commercially available from Aldrich Chemical Company, Milwaukee, Wisconsin USA) is treated with phosphorus oxychloride and *N,N*-dimethylaniline according to the procedure described in Whittaker, *J. Chem. Soc.* **1951**, 1565 to give 1,3-dichloro-4-nitropyrimidine, **2**.

1,3-Dichloro-4-nitropyrimidine, **2**, is then reacted with about one molar equivalent of an amino acid derivative of the formula:  
10  $H_2N-CH(R^3)C(O)X$  where  $R^3$  and  $X$  are as defined herein or  $X$  is  $-OP^1$  where  $P^1$  is a carboxylic acid protecting group, in the presence of a trialkylamine, such as diisopropylethylamine (DIEA). Typically, this reaction is conducted in an inert diluent, such as dichloromethane, at a temperature ranging from about 0°C to about 10°C for about 5 min. to about 6 hours to afford  
15 intermediate **3**.

The nitro group of intermediate **3** is then reduced using a conventional reducing agent, such as hydrogen and a palladium on carbon catalyst. When hydrogen and palladium on carbon are employed as the reducing agent, the  
20 chloro group of intermediate **3** is also removed. This reaction is typically conducted by contacting **3** with a Degussa-type palladium on carbon catalyst (typically 20%) and excess sodium bicarbonate in an inert diluent, such as methanol, under hydrogen (typically about 55 psi) for about 12 to 36 hours at ambient temperature to afford amino intermediate **4**.

25 Amino intermediate **4** is then reacted with a sulfonyl chloride of the formula:  $R^5-S(O)_2-Cl$ , where  $R^5$  is as defined herein, to provide sulfonamide intermediate **5**. This reaction is typically conducted by reacting the amino intermediate **4** with at least one equivalent, preferably about 1.1 to about 2  
30 equivalents, of the sulfonyl chloride in an inert diluent such as



dichloromethane and the like. Generally, the reaction is conducted at a temperature ranging from about -70°C to about 40°C for about 1 to about 24 hours. Preferably, this reaction is conducted in the presence of a suitable base to scavenge the acid generated during the reaction. Suitable bases  
5 include, by way of example, tertiary amines, such as triethylamine, diisopropylethylamine, *N*-methylmorpholine and the like. Alternatively, the reaction can be conducted under Schotten-Baumann-type conditions using aqueous alkali, such as sodium hydroxide and the like, as the base. Upon completion of the reaction, the resulting sulfonamide **5** is recovered by  
10 conventional methods including neutralization, extraction, precipitation, chromatography, filtration, and the like.

Other heteroaryl intermediates may also be employed in the above described reactions including, but not limited to, 2-chloro-3-nitropyrazine (*J. Med. Chem.* **1984**, 27, 1634); 4-chloro-5-nitroimidazole (*J. Chem. Soc.* **1930**, 268); and the like.

The amino acid derivatives employed in the above reactions are either known compounds or compounds that can be prepared from known  
20 compounds by conventional synthetic procedures. For example, amino acid derivatives can be prepared by *C*-alkylating commercially available diethyl 2-acetamidomalonate (Aldrich, Milwaukee, Wisconsin, USA) with an alkyl or substituted alkyl halide. This reaction is typically conducted by treating the diethyl 2-acetamidomalonate with at least one equivalent of sodium ethoxide  
25 and at least one equivalent of an alkyl or substituted alkyl halide in refluxing ethanol for about 6 to about 12 hours. The resulting *C*-alkylated malonate is then deacetylated, hydrolyzed and decarboxylated by heating in aqueous hydrochloric acid at reflux for about 6 to about 12 hours to provide the amino acid, typically as the hydrochloride salt.

Examples of amino acid derivatives suitable for use in the above reactions include, but are not limited to, L-alanine methyl ester, L-isoleucine methyl ester, L-leucine methyl ester, L-valine methyl ester,  $\beta$ -*tert*-butyl-L-aspartic acid methyl ester, L-asparagine *tert*-butyl ester,  $\epsilon$ -Boc-L-lysine methyl ester,  $\epsilon$ -Cbz-L-lysine methyl ester,  $\gamma$ -*tert*-butyl-L-glutamic acid methyl ester, L-glutamine *tert*-butyl ester, L-(*N*-methyl)histidine methyl ester, L-(*N*-benzyl)histidine methyl ester, L-methionine methyl ester, L-(*O*-benzyl)serine methyl ester, L-tryptophan methyl ester, L-phenylalanine methyl ester, L-phenylalanine isopropyl ester, L-phenylalanine benzyl ester, L-phenylalaninamide, *N*-methyl-L-phenylalanine benzyl ester, 3-carboxy-D,L-phenylalanine methyl ester, 4-carboxy-D,L-phenylalanine methyl ester, L-4-chlorophenylalanine methyl ester, L-4-(3-dimethylaminopropoxy)-phenylalanine methyl ester, L-4-iodophenylalanine methyl ester, L-3,4-methylenedioxyphenylalanine methyl ester, L-3,4-ethylenedioxyphenylalanine methyl ester, L-4-nitrophenylalanine methyl ester, L-tyrosine methyl ester, D,L-homophenylalanine methyl ester, L-(*O*-methyl)tyrosine methyl ester, L-(*O*-*tert*-butyl)tyrosine methyl ester, L-(*O*-benzyl)tyrosine methyl ester, L-3,5-diiodotyrosine methyl ester, L-3-iodotyrosine methyl ester,  $\beta$ -(1-naphthyl)-L-alanine methyl ester,  $\beta$ -(2-naphthyl)-L-alanine methyl ester,  $\beta$ -(2-thienyl)-L-alanine methyl ester,  $\beta$ -cyclohexyl-L-alanine methyl ester,  $\beta$ -(2-pyridyl)-L-alanine methyl ester,  $\beta$ -(3-pyridyl)-L-alanine methyl ester,  $\beta$ -(4-pyridyl)-L-alanine methyl ester,  $\beta$ -(2-thiazolyl)-D,L-alanine methyl ester,  $\beta$ -(1,2,4-triazol-3-yl)-D,L-alanine methyl ester, and the like. If desired, of course, other esters or amides of the above-described compounds may also be employed.

Additionally,  $\alpha$ -hydroxy and  $\alpha$ -thio carboxylic acids may also be employed in the above-described reactions. Such compounds are well-known in the art and are either commercially available or may be prepared from

-- 117 --

commercially available starting materials using conventional reagents and reaction conditions.

5       The sulfonyl chlorides employed in the above reaction are also either known compounds or compounds that can be prepared from known compounds by conventional synthetic procedures. Such compounds are typically prepared from the corresponding sulfonic acid, i.e., from compounds of the formula  $R^5-SO_3H$  where  $R^5$  is as defined above, using phosphorous trichloride and phosphorous pentachloride. This reaction is  
10       generally conducted by contacting the sulfonic acid with about 2 to 5 molar equivalents of phosphorous trichloride and phosphorous pentachloride, either neat or in an inert solvent, such as dichloromethane, at temperature in the range of about 0°C to about 80°C for about 1 to about 48 hours to afford the sulfonyl chloride. Alternatively, the sulfonyl chloride can be prepared from  
15       the corresponding thiol compound, i.e., from compounds of the formula  $R^5-SH$  where  $R^5$  is as defined herein, by treating the thiol with chlorine ( $Cl_2$ ) and water under conventional reaction conditions.

Examples of sulfonyl chlorides suitable for use in this invention  
20       include, but are not limited to, methanesulfonyl chloride, 2-propanesulfonyl chloride, 1-butesulfonyl chloride, benzenesulfonyl chloride, 1-naphthalenesulfonyl chloride, 2-naphthalenesulfonyl chloride, *p*-toluenesulfonyl chloride,  $\alpha$ -toluenesulfonyl chloride, 4-acetamidobenzenesulfonyl chloride, 4-amidinobenzenesulfonyl chloride, 4-  
25       *tert*-butylbenzenesulfonyl chloride, 4-bromobenzenesulfonyl chloride, 2-carboxybenzenesulfonyl chloride, 4-cyanobenzenesulfonyl chloride, 3,4-dichlorobenzenesulfonyl chloride, 3,5-dichlorobenzenesulfonyl chloride, 3,4-dimethoxybenzenesulfonyl chloride, 3,5-ditrifluoromethylbenzenesulfonyl chloride, 4-fluorobenzenesulfonyl chloride, 4-methoxybenzenesulfonyl  
30       chloride, 2-methoxycarbonylbenzenesulfonyl chloride, 4-

-- 118 --


5 methylamidobenzenesulfonyl chloride, 4-nitrobenzenesulfonyl chloride, 4-thioamidobenzenesulfonyl chloride, 4-trifluoromethylbenzenesulfonyl chloride, 4-trifluoromethoxybenzenesulfonyl chloride, 2,4,6-trimethylbenzenesulfonyl chloride, 2-phenylethanesulfonyl chloride, 2-thiophenesulfonyl chloride, 5-chloro-2-thiophenesulfonyl chloride, 2,5-dichloro-4-thiophenesulfonyl chloride, 2-thiazolesulfonyl chloride, 2-methyl-4-thiazolesulfonyl chloride, 1-methyl-4-imidazolesulfonyl chloride, 1-methyl-4-pyrazolesulfonyl chloride, 5-chloro-1,3-dimethyl-4-pyrazolesulfonyl chloride, 3-pyridinesulfonyl chloride, 2-pyrimidinesulfonyl chloride and the  
10 like. If desired, a sulfonyl fluoride, sulfonyl bromide or sulfonic acid anhydride may be used in place of the sulfonyl chloride in the above reaction to form the sulfonamide intermediate 5.

15 If desired, sulfonamide intermediate 5 can be alkylated at the sulfonamide nitrogen atom to provide compound 6. For example, 5 can be contacted with excess diazomethane (generated, for example, using 1-methyl-3-nitro-1-nitrosoguanidine and sodium hydroxide) to afford 6 where R<sup>6</sup> is methyl. Other conventional alkylation procedures and reagents may also be employed to prepare various compounds of this invention.

20

In another preferred embodiment, compounds of this invention may be prepared by displacement of a leaving group as shown in Scheme 2:

Scheme 2



where  $R^3$ , Q and X are as defined herein; A' is heteroaryl, substituted heteroaryl, heterocyclic or substituted heterocyclic containing two nitrogen atoms in the heteroaryl or heterocyclic ring; and  $L^1$  is a leaving group, such as chloro, bromo, iodo, sulfonate ester and the like.

5


Typically, this reaction is conducted by combining approximately stoichiometric equivalents of 7 and 8 in a suitable inert diluent such as water, dimethylsulfoxide (DMSO) and the like, with an excess of a suitable base such as sodium bicarbonate, sodium hydroxide, etc. to scavenge the acid generated by the reaction. The reaction is preferably conducted at from about 25°C to about 100°C until reaction completion which typically occurs within 1 to about 24 hours. This reaction is further described in U.S. Patent No. 3,598,859, which is incorporated herein by reference in its entirety. Upon reaction completion, the product 9 is recovered by conventional methods including precipitation, chromatography, filtration and the like.

10  
15

In still another alternative embodiment, compounds of this invention in which Q is  $NR^4$  can be prepared by reductive amination of a suitable 2-oxocarboxylic acid ester, 10, such as a pyruvate ester, as shown in Scheme 3:

20

Scheme 3




where A',  $R^3$  and X are as defined herein.

-- 120 --

Generally, this reaction is conducted by combining equamolar amounts of **10** and **11** in an inert diluent such as methanol, ethanol and the like under conditions which provide for imine formation (not shown). The imine formed is then reduced under conventional conditions by a suitable  
5 reducing agent such as sodium cyanoborohydride,  $H_2$ /palladium on carbon and the like to form the product **12**. In a particularly preferred embodiment, the reducing agent is  $H_2$ /palladium on carbon which is incorporated into the initial reaction medium thereby permitting imine reduction *in situ* in a one pot  
10 procedure to provide **12**. The reaction is preferably conducted at from about  $20^\circ C$  to about  $80^\circ C$  at a pressure of from 1 to 10 atmospheres until reaction completion which typically occurs within 1 to about 24 hours. Upon reaction completion, the product **12** is recovered by conventional methods including chromatography, filtration and the like.

15 Alternatively, certain compounds of this invention can be prepared via a rhodium-catalyzed insertion reaction as shown in Scheme 4:

Scheme 4




where  $A''$  is heteroaryl or substituted heteroaryl containing two nitrogen atoms in the heteroaryl ring, and  $R^3$  and  $X$  (preferably alkoxy) are as defined  
20 herein. Typically, this reaction is conducted using rhodium acetate dimer,  $Rh_2(OAc)_4$ , in an inert diluent such as toluene at a temperature ranging from about  $25^\circ C$  to about  $80^\circ C$  for about 1 to 12 hours to afford **15**. This

-- 121 --

reaction is described further in B. R. Henke et. al., *J. Med. Chem.* **1998**, *41*, 5020-5036 and references cited therein.

Similarly, certain compounds of this invention can be prepared by the  
5 copper-catalyzed coupling reaction shown in Scheme 5:

Scheme 5



where A'' is as defined herein, X<sup>3</sup> is halogen, such as chloro, bromo or iodo (preferably iodo), and R<sup>3</sup> and X (preferably alkoxy) are as defined herein. Typically, this reaction is conducted using copper iodide (CuI) and potassium carbonate in an inert diluent such as *N,N*-dimethyl acetamide (DMA) at a  
10 temperature ranging from about 60°C to about 120°C for about 12 to 36 hours to afford 15. This reaction is described further in D. Ma et. al., *J. Am. Chem. Soc.* **1998**, *120*, 12459-12467 and references cited therein.

15 For ease of synthesis, the compounds of this invention are typically prepared as an ester, i.e., where X is an alkoxy or substituted alkoxy group and the like. If desired, the ester group can be hydrolysed using conventional conditions and reagents to provide the corresponding carboxylic acid. Typically, this reaction is conducted by treating the ester with at least one  
20 equivalent of an alkali metal hydroxide, such as lithium, sodium or potassium hydroxide, in an inert diluent, such as methanol or mixtures of methanol and water, at a temperature ranging about 0°C to about 24°C for about 1 to about 12 hours. Alternatively, benzyl esters may be removed by hydrogenolysis

-- 122 --

using a palladium catalyst, such as palladium on carbon, and *tert*-butyl esters can be removed using formic acid to afford the corresponding carboxylic acid.

5           As will be apparent to those skilled in the art, other functional groups present on any of the substituents of the compounds of formulas I-VII can be readily modified or derivatized either before or after the above-described synthetic reactions using well-known synthetic procedures. For example, a  
10       nitro group present on a substituent of a compound of formula I-VII or an intermediate thereof may be readily reduced by hydrogenation in the presence of a palladium catalyst, such as palladium on carbon, to provide the corresponding amino group. This reaction is typically conducted at a temperature of from about 20°C to about 50°C for about 6 to about 24 hours in an inert diluent, such as methanol. Compounds having a nitro group on  
15       the R<sup>3</sup> and/or R<sup>3'</sup> substituent can be prepared, for example, by using a 4-nitrophenylalanine derivative and the like in the above-described coupling reactions.

20           Similarly, a pyridyl group can be hydrogenated in the presence of a platinum catalyst, such as platinum oxide, in an acidic diluent to provide the corresponding piperidinyl analogue. Generally, this reaction is conducted by treating the pyridine compound with hydrogen at a pressure ranging from about 20 psi to about 60 psi, preferably about 40 psi, in the presence of the catalyst at a temperature of about 20°C to about 50°C for about 2 to about 24  
25       hours in an acidic diluent, such as a mixture of methanol and aqueous hydrochloric acid.

30           Additionally, when the R<sup>3</sup> and/or R<sup>3'</sup> substituent of a compound of formula I-VII or an intermediate thereof contains a primary or secondary amino group, such amino groups can be further derivatized either before or



after the above coupling reactions to provide, by way of example, amides, sulfonamides, ureas, thioureas, carbamates, secondary or tertiary amines and the like. Compounds having a primary amino group on the R<sup>3</sup> and/or R<sup>3'</sup> substituent may be prepared, for example, by reduction of the corresponding  
5 nitro compound as described above.

By way of illustration, a compound of formula I-VII or an intermediate thereof having a substituent containing a primary or secondary amino group, such as where R<sup>3</sup> is a (4-aminophenyl)methyl group, can be  
10 readily *N*-acylated using conventional acylating reagents and conditions to provide the corresponding amide. This acylation reaction is typically conducted by treating the amino compound with at least one equivalent, preferably about 1.1 to about 1.2 equivalents, of a carboxylic acid in the presence of a coupling reagent such as a carbodiimide, BOP reagent  
15 (benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate) and the like, in an inert diluent, such as dichloromethane, chloroform, acetonitrile, tetrahydrofuran, *N,N*-dimethylformamide and the like, at a temperature ranging from about 0°C to about 37°C for about 4 to about 24 hours. Preferably, a promoter, such as  
20 *N*-hydroxysuccinimide, 1-hydroxy-benzotriazole and the like, is used to facilitate the acylation reaction. Examples of carboxylic acids suitable for use in this reaction include, but are not limited to, *N*-*tert*-butyloxycarbonylglycine, *N*-*tert*-butyloxycarbonyl-L-phenylalanine, *N*-*tert*-butyloxycarbonyl-L-aspartic acid benzyl ester, benzoic acid, *N*-*tert*-  
25 butyloxycarbonylisonipecotic acid, *N*-methylisonipecotic acid, *N*-*tert*-butyloxycarbonylnipecotic acid, *N*-*tert*-butyloxycarbonyl-L-tetrahydroisoquinoline-3-carboxylic acid, *N*-(toluene-4-sulfonyl)-L-proline and the like.

-- 124 --

Alternatively, a compound of formula I-VII or an intermediate thereof containing a primary or secondary amino group can be *N*-acylated using an acyl halide or a carboxylic acid anhydride to form the corresponding amide. This reaction is typically conducted by contacting the amino compound with  
5 at least one equivalent, preferably about 1.1 to about 1.2 equivalents, of the acyl halide or carboxylic acid anhydride in an inert diluent, such as dichloromethane, at a temperature ranging from about -70°C to about 40°C for about 1 to about 24 hours. If desired, an acylation catalyst such as 4-(*N,N*-dimethylamino)pyridine may be used to promote the acylation reaction.  
10 The acylation reaction is preferably conducted in the presence of a suitable base to scavenge the acid generated during the reaction. Suitable bases include, by way of example, tertiary amines, such as triethylamine, diisopropylethylamine, *N*-methylemorpholine and the like. Alternatively, the reaction can be conducted under Schotten-Baumann-type conditions using  
15 aqueous alkali, such as sodium hydroxide and the like.

Examples of acyl halides and carboxylic acid anhydrides suitable for use in this reaction include, but are not limited to, 2-methylpropionyl chloride, trimethylacetyl chloride, phenylacetyl chloride, benzoyl chloride, 2-  
20 bromobenzoyl chloride, 2-methylbenzoyl chloride, 2-trifluoro-methylbenzoyl chloride, isonicotinoyl chloride, nicotinoyl chloride, picolinoyl chloride, acetic anhydride, succinic anhydride, and the like. Carbamyl chlorides, such as *N,N*-dimethylcarbamyl chloride, *N,N*-diethylcarbamyl chloride and the like, can also be used in this reaction to provide ureas. Similarly,  
25 dicarbonates, such as di-*tert*-butyl dicarbonate, may be employed to provide carbamates.

In a similar manner, a compound of formula I-VII or an intermediate thereof containing a primary or secondary amino group may be *N*-sulfonated  
30 to form a sulfonamide using a sulfonyl halide or a sulfonic acid anhydride.

-- 125 --

Sulfonyl halides and sulfonic acid anhydrides suitable for use in this reaction include, but are not limited to, methanesulfonyl chloride, chloromethanesulfonyl chloride, *p*-toluenesulfonyl chloride, trifluoromethanesulfonic anhydride, and the like. Similarly, sulfamoyl chlorides, such as dimethylsulfamoyl chloride, can be used to provide sulfamides (e.g.,  $>N-SO_2-N<$ ).

Additionally, a primary and secondary amino group present on a substituent of a compound of formula I-VII or an intermediate thereof can be reacted with an isocyanate or a thioisocyanate to give a urea or thiourea, respectively. This reaction is typically conducted by contacting the amino compound with at least one equivalent, preferably about 1.1 to about 1.2 equivalents, of the isocyanate or thioisocyanate in an inert diluent, such as toluene and the like, at a temperature ranging from about 24°C to about 37°C for about 12 to about 24 hours. The isocyanates and thioisocyanates used in this reaction are commercially available or can be prepared from commercially available compounds using well-known synthetic procedures. For example, isocyanates and thioisocyanates are readily prepared by reacting the appropriate amine with phosgene or thiophosgene. Examples of isocyanates and thioisocyanates suitable for use in this reaction include, but are not limited to, ethyl isocyanate, *n*-propyl isocyanate, 4-cyanophenyl isocyanate, 3-methoxyphenyl isocyanate, 2-phenylethyl isocyanate, methyl thioisocyanate, ethyl thioisocyanate, 2-phenylethyl thioisocyanate, 3-phenylpropyl thioisocyanate, 3-(*N,N*-diethylamino)propyl thioisocyanate, phenyl thioisocyanate, benzyl thioisocyanate, 3-pyridyl thioisocyanate, fluorescein isothiocyanate (isomer I) and the like.

Furthermore, when a compound of formula I-VII or an intermediate thereof contains a primary or secondary amino group, the amino group can be reductively alkylated using aldehydes or ketones to form a secondary or

-- 126 --

tertiary amino group. This reaction is typically conducted by contacting the amino compound with at least one equivalent, preferably about 1.1 to about 1.5 equivalents, of an aldehyde or ketone and at least one equivalent based on the amino compound of a metal hydride reducing agent, such as sodium cyanoborohydride, in an inert diluent, such as methanol, tetrahydrofuran, mixtures thereof and the like, at a temperature ranging from about 0°C to about 50°C for about 1 to about 72 hours. Aldehydes and ketones suitable for use in this reaction include, by way of example, benzaldehyde, 4-chlorobenzaldehyde, valeraldehyde and the like.

10

In a similar manner, when a compound of formula I-VII or an intermediate thereof has a substituent containing a hydroxyl group, the hydroxyl group can be further modified or derivatized either before or after the above coupling reactions to provide, by way of example, ethers, carbamates and the like. Compounds having a hydroxyl group on the R<sup>3</sup> substituent, for example, can be prepared using an amino acid derivative derived from tyrosine and the like in the above-described reactions.

By way of example, a compound of formula I-VII or an intermediate thereof having a substituent containing a hydroxyl group, such as where R<sup>3</sup> is a (4-hydroxyphenyl)methyl group, can be readily *O*-alkylated to form ethers. This *O*-alkylation reaction is typically conducted by contacting the hydroxy compound with a suitable alkali or alkaline earth metal base, such as potassium carbonate, in an inert diluent, such as acetone, 2-butanone and the like, to form the alkali or alkaline earth metal salt of the hydroxyl group. This salt is generally not isolated, but is reacted *in situ* with at least one equivalent of an alkyl or substituted alkyl halide or sulfonate, such as an alkyl chloride, bromide, iodide, mesylate or tosylate, to afford the ether. Generally, this reaction is conducted at a temperature ranging from about 60°C to about 150°C for about 24 to about 72 hours. Preferably, a catalytic

30

amount of sodium or potassium iodide is added to the reaction mixture when an alkyl chloride or bromide is employed in the reaction.

Examples of alkyl or substituted alkyl halides and sulfonates suitable for use in this reaction include, but are not limited to, *tert*-butyl bromoacetate, *N-tert*-butyl chloroacetamide, 1-bromoethylbenzene, ethyl  $\alpha$ -bromophenylacetate, 2-(*N*-ethyl-*N*-phenylamino)ethyl chloride, 2-(*N,N*-ethylamino)ethyl chloride, 2-(*N,N*-diisopropylamino)ethyl chloride, 2-(*N,N*-dibenzylamino)ethyl chloride, 3-(*N,N*-ethylamino)propyl chloride, 3-(*N*-benzyl-*N*-methylamino)propyl chloride, *N*-(2-chloroethyl)morpholine, 2-(hexamethyleneimino)ethyl chloride, 3-(*N*-methylpiperazine)propyl chloride, 1-(3-chlorophenyl)-4-(3-chloropropyl)piperazine, 2-(4-hydroxy-4-phenylpiperidine)ethyl chloride, *N-tert*-butyloxycarbonyl-3-piperidinemethyl tosylate, and the like.

Alternatively, a hydroxyl group present on a substituent of a compound of formula I-VII or an intermediate thereof can be *O*-alkylating using the Mitsunobu reaction. In this reaction, an alcohol, such as 3-(*N,N*-dimethylamino)-1-propanol and the like, is reacted with about 1.0 to about 1.3 equivalents of triphenylphosphine and about 1.0 to about 1.3 equivalents of diethyl azodicarboxylate in an inert diluent, such as tetrahydrofuran, at a temperature ranging from about -10°C to about 5°C for about 0.25 to about 1 hour. About 1.0 to about 1.3 equivalents of a hydroxy compound, such as *N-tert*-butyltyrosine methyl ester, is then added and the reaction mixture is stirred at a temperature of about 0°C to about 30°C for about 2 to about 48 hours to provide the *O*-alkylated product.

In a similar manner, a compound of formula I-VII or an intermediate thereof containing an aryl hydroxy group can be reacted with an aryl iodide to provide a diaryl ether. Generally, this reaction is conducted by forming

-- 128 --

the alkali metal salt of the hydroxyl group using a suitable base, such as sodium hydride, in an inert diluent such as xylenes at a temperature of about -25°C to about 10°C. The salt is then treated with about 1.1 to about 1.5 equivalents of cuprous bromide dimethyl sulfide complex at a temperature ranging from about 10°C to about 30°C for about 0.5 to about 2.0 hours, followed by about 1.1 to about 1.5 equivalents of an aryl iodide, such as sodium 2-iodobenzoate and the like. The reaction is then heated to about 70°C to about 150°C for about 2 to about 24 hours to provide the diaryl ether.

10

Additionally, a hydroxy-containing compound can also be readily derivatized to form a carbamate. In one method for preparing such carbamates, a hydroxy compound of formula I-VII or an intermediate thereof is contacted with about 1.0 to about 1.2 equivalents of 4-nitrophenyl chloroformate in an inert diluent, such as dichloromethane, at a temperature ranging from about -25°C to about 0°C for about 0.5 to about 2.0 hours. Treatment of the resulting carbonate with an excess, preferably about 2 to about 5 equivalents, of a trialkylamine, such as triethylamine, for about 0.5 to 2 hours, followed by about 1.0 to about 1.5 equivalents of a primary or secondary amine provides the carbamate. Examples of amines suitable for using in this reaction include, but are not limited to, piperazine, 1-methylpiperazine, 1-acetylpiperazine, morpholine, thiomorpholine, pyrrolidine, piperidine and the like.

15

20

Alternatively, in another method for preparing carbamates, a hydroxy-containing compound is contacted with about 1.0 to about 1.5 equivalents of a carbamyl chloride in an inert diluent, such as dichloromethane, at a temperature ranging from about 25°C to about 70°C for about 2 to about 72 hours. Typically, this reaction is conducted in the presence of a suitable base to scavenge the acid generated during the reaction. Suitable bases include, by

25

30

-- 129 --

way of example, tertiary amines, such as triethylamine, diisopropylethylamine, *N*-methylmorpholine and the like. Additionally, at least one equivalent (based on the hydroxy compound) of 4-(*N,N*-dimethylamino)pyridine is preferably added to the reaction mixture to  
5 facilitate the reaction. Examples of carbamyl chlorides suitable for use in this reaction include, by way of example, dimethylcarbamyl chloride, diethylcarbamyl chloride and the like.

Likewise, when a compound of formula I-VII or an intermediate  
10 thereof contains a primary or secondary hydroxyl group, such hydroxyl groups can be readily converted into a leaving group and displaced to form, for example, amines, sulfides and fluorides. Generally, when a chiral compound is employed in these reactions, the stereochemistry at the carbon atom attached to the derivatized hydroxyl group is typically inverted.

15 These reactions are typically conducted by first converting the hydroxyl group into a leaving group, such as a tosylate, by treatment of the hydroxy compound with at least one equivalent of a sulfonyl halide, such as *p*-toluenesulfonyl chloride and the like, in pyridine. This reaction is  
20 generally conducted at a temperature of from about 0°C to about 70°C for about 1 to about 48 hours. The resulting tosylate can then be readily displaced with sodium azide, for example, by contacting the tosylate with at least one equivalent of sodium azide in an inert diluent, such as a mixture of *N,N*-dimethylformamide and water, at a temperature ranging from about 0°C  
25 to about 37°C for about 1 to about 12 hours to provide the corresponding azido compound. The azido group can then be reduced by, for example, hydrogenation using a palladium on carbon catalyst to provide the amino (-NH<sub>2</sub>) compound.

-- 130 --

Similarly, a tosylate group can be readily displaced by a thiol to form a sulfide. This reaction is typically conducted by contacting the tosylate with at least one equivalent of a thiol, such as thiophenol, in the presence of a suitable base, such as 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), in an inert  
5 diluent, such as *N,N*-dimethylformamide, at a temperature of from about 0°C to about 37°C for about 1 to about 12 hours to provide the sulfide. Additionally, treatment of a tosylate with morpholinosulfur trifluoride in an inert diluent, such as dichloromethane, at a temperature ranging from about 0°C to about 37°C for about 12 to about 24 hours affords the corresponding  
10 fluoro compound.

Furthermore, a compound of formula I-VII or an intermediate thereof having a substituent containing an iodoaryl group, for example, when R<sup>3</sup> is a (4-iodophenyl)methyl group, can be readily converted either before or after  
15 the above coupling reactions into a biaryl compound. Typically, this reaction is conducted by treating the iodoaryl compound with about 1.1 to about 2 equivalents of an arylzinc iodide, such as 2-(methoxycarbonyl)phenylzinc iodide, in the presence of a palladium catalyst, such as palladium tetra(triphenylphosphine), in an inert diluent, such as tetrahydrofuran, at a  
20 temperature ranging from about 24°C to about 30°C until reaction completion. This reaction is further described, for example, in Rieke, *J. Org. Chem.* **1991**, 56, 1445. Additional methods for preparing biaryl derivatives are disclosed in International Publication Number WO 98/53817, published December 3, 1998, the disclosure of which is incorporated herein  
25 by reference in its entirety.

In some cases, the compounds of formula I-VII or intermediates thereof may contain substituents having one or more sulfur atoms. When present, such sulfur atoms can be oxidized either before or after the above  
30 coupling reactions to provide a sulfoxide or sulfone compound using



conventional reagents and reaction conditions. Suitable reagents for oxidizing a sulfide compound to a sulfoxide include, by way of example, hydrogen peroxide, 3-chloroperoxybenzoic acid (MCPBA), sodium periodate and the like. The oxidation reaction is typically conducted by contacting the sulfide compound with about 0.95 to about 1.1 equivalents of the oxidizing reagent in an inert diluent, such as dichloromethane, at a temperature ranging from about -50°C to about 75°C for about 1 to about 24 hours. The resulting sulfoxide can then be further oxidized to the corresponding sulfone by contacting the sulfoxide with at least one additional equivalent of an oxidizing reagent, such as hydrogen peroxide, MCPBA, potassium permanganate and the like. Alternatively, the sulfone can be prepared directly by contacting the sulfide with at least two equivalents, and preferably an excess, of the oxidizing reagent. Such reactions are described further in March, "Advanced Organic Chemistry", 4th Ed., pp. 1201-1202, Wiley Publisher, 1992.

Other procedures and reaction conditions for preparing the compounds of this invention are described in the examples set forth below. Additionally, other procedures for preparing compounds useful in certain aspects of this invention are disclosed in U.S. Serial No. \_\_/\_\_\_\_, filed on even date herewith, entitled "Compounds Which Inhibit Leucocyte Adhesion Mediated by VLA-4" (Attorney Docket No. 002010-525); the disclosure of which is incorporated herein by reference in its entirety.

#### Pharmaceutical Formulations

When employed as pharmaceuticals, the compounds of this invention are usually administered in the form of pharmaceutical compositions. These compounds can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, and intranasal. These compounds are effective as both injectable and oral compositions. Such

-- 132 --

compositions are prepared in a manner well known in the pharmaceutical art and comprise at least one active compound.

5        This invention also includes pharmaceutical compositions which  
contain, as the active ingredient, one or more of the compounds of formula I-  
VII above associated with pharmaceutically acceptable carriers. In making  
the compositions of this invention, the active ingredient is usually mixed with  
an excipient, diluted by an excipient or enclosed within such a carrier which  
can be in the form of a capsule, sachet, paper or other container. The  
10        excipient employed is typically an excipient suitable for administration to  
human subjects or other mammals. When the excipient serves as a diluent, it  
can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier  
or medium for the active ingredient. Thus, the compositions can be in the  
form of tablets, pills, powders, lozenges, sachets, cachets, elixirs,  
15        suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid  
medium), ointments containing, for example, up to 10% by weight of the  
active compound, soft and hard gelatin capsules, suppositories, sterile  
injectable solutions, and sterile packaged powders.

20        In preparing a formulation, it may be necessary to mill the active  
compound to provide the appropriate particle size prior to combining with the  
other ingredients. If the active compound is substantially insoluble, it  
ordinarily is milled to a particle size of less than 200 mesh. If the active  
compound is substantially water soluble, the particle size is normally adjusted  
25        by milling to provide a substantially uniform distribution in the formulation,  
e.g. about 40 mesh.

30        Some examples of suitable excipients include lactose, dextrose,  
sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate,  
alginate, tragacanth, gelatin, calcium silicate, microcrystalline cellulose,

polyvinylpyrrolidone, cellulose, water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxy-  
5 benzoates; sweetening agents; and flavoring agents. The compositions of the invention can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art.

10 The compositions are preferably formulated in a unit dosage form, each dosage containing from about 5 to about 100 mg, more usually about 10 to about 30 mg, of the active ingredient. The term "unit dosage forms" refers to physically discrete units suitable as unitary dosages for human  
15 subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

The active compound is effective over a wide dosage range and is generally administered in a pharmaceutically effective amount. It, will be  
20 understood, however, that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

25 For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical excipient to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention. When referring to these preformulation  
30 compositions as homogeneous, it is meant that the active ingredient is

-- 134 --

dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation is then subdivided into unit dosage forms of the type described above containing from, for example, 0.1  
5 to about 500 mg of the active ingredient of the present invention.

The tablets or pills of the present invention may be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner  
10 dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permit the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or  
15 coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

The liquid forms in which the novel compositions of the present  
20 invention may be incorporated for administration orally or by injection include aqueous solutions suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil, or peanut oil, as well as elixirs and similar  
pharmaceutical vehicles.

25

Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described *supra*.  
30 Preferably the compositions are administered by the oral or nasal respiratory

-- 135 --

route for local or systemic effect. Compositions in preferably pharmaceutically acceptable solvents may be nebulized by use of inert gases. Nebulized solutions may be breathed directly from the nebulizing device or the nebulizing device may be attached to a face masks tent, or intermittent  
5 positive pressure breathing machine. Solution, suspension, or powder compositions may be administered, preferably orally or nasally, from devices which deliver the formulation in an appropriate manner.

The following formulation examples illustrate the pharmaceutical  
10 compositions of the present invention.

Formulation Example 1

Hard gelatin capsules containing the following ingredients are prepared:

15	<u>Ingredient</u>	<u>Quantity (mg/capsule)</u>
	Active Ingredient	30.0
	Starch	305.0
20	Magnesium stearate	5.0

The above ingredients are mixed and filled into hard gelatin capsules in 340 mg quantities.

25 Formulation Example 2

A tablet formula is prepared using the ingredients below:

	<u>Ingredient</u>	<u>Quantity (mg/tablet)</u>
30	Active Ingredient	25.0
	Cellulose, microcrystalline	200.0
	Colloidal silicon dioxide	10.0
	Stearic acid	5.0

-- 136 --

The components are blended and compressed to form tablets, each weighing 240 mg.

Formulation Example 3

5 A dry powder inhaler formulation is prepared containing the following components:

	<u>Ingredient</u>	<u>Weight %</u>
10	Active Ingredient	5
	Lactose	95

The active mixture is mixed with the lactose and the mixture is added  
15 to a dry powder inhaling appliance.

Formulation Example 4

Tablets, each containing 30 mg of active ingredient, are prepared as follows:

	<u>Ingredient</u>	<u>Quantity (mg/tablet)</u>
20	Active Ingredient	30.0 mg
	Starch	45.0 mg
25	Microcrystalline cellulose	35.0 mg
	Polyvinylpyrrolidone (as 10% solution in water)	4.0 mg
	Sodium carboxymethyl starch	4.5 mg
30	Magnesium stearate	0.5 mg
	Talc	<u>1.0 mg</u>
	Total	120 mg

35 The active ingredient, starch and cellulose are passed through a No. 20 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders, which are then passed through a 16 mesh U.S. sieve. The granules so produced are dried at 50° to

-- 137 --

60°C and passed through a 16 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate, and talc, previously passed through a No. 30 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

5

Formulation Example 5

Capsules, each containing 40 mg of medicament are made as follows:

10	<u>Ingredient</u>	<u>Quantity</u> <u>(mg/capsule)</u>
	Active Ingredient	40.0 mg
	Starch	109.0 mg
	Magnesium stearate	<u>1.0 mg</u>
15	Total	150.0 mg

The active ingredient, cellulose, starch, an magnesium stearate are blended, passed through a No. 20 mesh U.S. sieve, and filled into hard gelatin capsules in 150 mg quantities.

20

Formulation Example 6

Suppositories, each containing 25 mg of active ingredient are made as follows:

25

	<u>Ingredient</u>	<u>Amount</u>
30	Active Ingredient	25 mg
	Saturated fatty acid glycerides to	2,000 mg

The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2.0 g capacity and allowed to cool.

35

-- 138 --

Formulation Example 7

Suspensions, each containing 50 mg of medicament per 5.0 ml dose are made as follows:

5	<u>Ingredient</u>	<u>Amount</u>
	Active Ingredient	50.0 mg
	Xanthan gum	4.0 mg
	Sodium carboxymethyl cellulose (11%)	
10	Microcrystalline cellulose (89%)	50.0 mg
	Sucrose	1.75 g
	Sodium benzoate	10.0 mg
	Flavor and Color	q.v.
15	Purified water to	5.0 ml

The medicament, sucrose and xanthan gum are blended, passed through a No. 10 mesh U.S. sieve, and then mixed with a previously made solution of the microcrystalline cellulose and sodium carboxymethyl cellulose in water.

20 The sodium benzoate, flavor, and color are diluted with some of the water and added with stirring. Sufficient water is then added to produce the required volume.

Formulation Example 8

25	<u>Ingredient</u>	<u>Quantity</u> <u>(mg/capsule)</u>
	Active Ingredient	15.0 mg
	Starch	407.0 mg
30	Magnesium stearate	<u>3.0 mg</u>
	Total	425.0 mg

35 The active ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 20 mesh U.S. sieve, and filled into hard gelatin capsules in 560 mg quantities.



-- 139 --

Formulation Example 9

An intravenous formulation may be prepared as follows:

5	<u>Ingredient</u>	<u>Quantity</u>
	Active Ingredient	250.0 mg
	Isotonic saline	1000 ml

Formulation Example 10

10 A topical formulation may be prepared as follows:

	<u>Ingredient</u>	<u>Quantity</u>
	Active Ingredient	1-10 g
15	Emulsifying Wax	30 g
	Liquid Paraffin	20 g
	White Soft Paraffin	to 100 g

20 The white soft paraffin is heated until molten. The liquid paraffin and emulsifying wax are incorporated and stirred until dissolved. The active ingredient is added and stirring is continued until dispersed. The mixture is then cooled until solid.

25 Another preferred formulation employed in the methods of the present invention employs transdermal delivery devices ("patches"). Such transdermal patches may be used to provide continuous or discontinuous infusion of the compounds of the present invention in controlled amounts. The construction and use of transdermal patches for the delivery of  
30 pharmaceutical agents is well known in the art. See, e.g., U.S. Patent 5,023,252, issued June 11, 1991, herein incorporated by reference. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

Direct or indirect placement techniques may be used when it is desirable or necessary to introduce the pharmaceutical composition to the brain.

Direct techniques usually involve placement of a drug delivery catheter into the host's ventricular system to bypass the blood-brain barrier. One such  
5 implantable delivery system used for the transport of biological factors to specific anatomical regions of the body is described in U.S. Patent 5,011,472 which is herein incorporated by reference.

Indirect techniques, which are generally preferred, usually involve  
10 formulating the compositions to provide for drug latentiation by the conversion of hydrophilic drugs into lipid-soluble drugs. Latentiation is generally achieved through blocking of the hydroxy, carbonyl, sulfate, and primary amine groups present on the drug to render the drug more lipid soluble and amenable to transportation across the blood-brain barrier.  
15 Alternatively, the delivery of hydrophilic drugs may be enhanced by intra-arterial infusion of hypertonic solutions which can transiently open the blood-brain barrier.

#### Utility

20 The compounds of this invention can be employed to bind VLA-4 ( $\alpha_4\beta_1$  integrin) in biological samples, i.e., the compounds bind VLA-4 with an  $IC_{50}$  of 15  $\mu M$  or less in a competitive binding assay as described herein. Accordingly, these compounds have utility in, for example, assaying such samples for VLA-4. In such assays, the compounds can be bound to a solid  
25 support and the VLA-4 sample added thereto. The amount of VLA-4 in the sample can be determined by conventional methods such as use of a sandwich ELISA assay. Alternatively, labeled VLA-4 can be used in a competitive assay to measure for the presence of VLA-4 in the sample. Other suitable assays are well known in the art.

In addition, certain of the compounds of this invention inhibit, *in vivo*, adhesion of leukocytes to endothelial cells mediated by VLA-4 by competitive binding to VLA-4. Accordingly, the compounds of this invention can be used in the treatment of diseases mediated by VLA-4 or leukocyte adhesion.

5 Such diseases include inflammatory diseases in mammalian patients such as asthma, Alzheimer's disease, atherosclerosis, AIDS dementia, diabetes (including acute juvenile onset diabetes), inflammatory bowel disease (including ulcerative colitis and Crohn's disease), multiple sclerosis, rheumatoid arthritis, tissue transplantation, tumor metastasis, meningitis,

10 encephalitis, stroke, and other cerebral traumas, nephritis, retinitis, atopic dermatitis, psoriasis, myocardial ischemia and acute leukocyte-mediated lung injury such as that which occurs in adult respiratory distress syndrome.

The biological activity of the compounds identified above may be

15 assayed in a variety of systems. For example, a compound can be immobilized on a solid surface and adhesion of cells expressing VLA-4 can be measured. Using such formats, large numbers of compounds can be screened. Cells suitable for this assay include any leukocytes known to express VLA-4 such as T cells, B cells, monocytes, eosinophils, and

20 basophils. A number of leukocyte cell lines can also be used, examples include Jurkat and U937.

The test compounds can also be tested for the ability to competitively inhibit binding between VLA-4 and VCAM-1, or between VLA-4 and a

25 labeled compound known to bind VLA-4 such as a compound of this invention or antibodies to VLA-4. In these assays, the VCAM-1 can be immobilized on a solid surface. VCAM-1 may also be expressed as a recombinant fusion protein having an Ig tail (e.g., IgG) so that binding to VLA-4 may be detected in an immunoassay. Alternatively, VCAM-1

30 expressing cells, such as activated endothelial cells or VCAM-1 transfected

fibroblasts can be used. For assays to measure the ability to block adhesion to brain endothelial cells, the assays described in International Patent Application Publication No. WO 91/05038 are particularly preferred. This application is incorporated herein by reference in its entirety.

5

Many assay formats employ labelled assay components. The labelling systems can be in a variety of forms. The label may be coupled directly or indirectly to the desired component of the assay according to methods well known in the art. A wide variety of labels may be used. The component  
10 may be labelled by any one of several methods. The most common method of detection is the use of autoradiography with  $^3\text{H}$ ,  $^{125}\text{I}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ , or  $^{32}\text{P}$  labelled compounds or the like. Non-radioactive labels include ligands which bind to labelled antibodies, fluorophores, chemiluminescent agents, enzymes and antibodies which can serve as specific binding pair members for a  
15 labelled ligand. The choice of label depends on sensitivity required, ease of conjugation with the compound, stability requirements, and available instrumentation.

Appropriate *in vivo* models for demonstrating efficacy in treating  
20 inflammatory responses include EAE (experimental autoimmune encephalomyelitis) in mice, rats, guinea pigs or primates, as well as other inflammatory models dependent upon  $\alpha 4$  integrins.

Compounds having the desired biological activity may be modified as  
25 necessary to provide desired properties such as improved pharmacological properties (e.g., *in vivo* stability, bio-availability), or the ability to be detected in diagnostic applications. Stability can be assayed in a variety of ways such as by measuring the half-life of the proteins during incubation with peptidases or human plasma or serum. A number of such protein stability

assays have been described (see, e.g., Verhoef et al., Eur. J. Drug Metab. Pharmacokinet., 1990, 15(2):83-93).

5 For diagnostic purposes, a wide variety of labels may be linked to the compounds, which may provide, directly or indirectly, a detectable signal. Thus, the compounds of the subject invention may be modified in a variety of ways for a variety of end purposes while still retaining biological activity. In addition, various reactive sites may be introduced at the terminus for linking to particles, solid substrates, macromolecules, or the like.

10 Labeled compounds can be used in a variety of *in vivo* or *in vitro* applications. A wide variety of labels may be employed, such as radionuclides (e.g., gamma-emitting radioisotopes such as technetium-99 or indium-111), fluorescers (e.g., fluorescein), enzymes, enzyme substrates, 15 enzyme cofactors, enzyme inhibitors, chemiluminescent compounds, bioluminescent compounds, and the like. Those of ordinary skill in the art will know of other suitable labels for binding to the complexes, or will be able to ascertain such using routine experimentation. The binding of these labels is achieved using standard techniques common to those of ordinary 20 skill in the art.

*In vitro* uses include diagnostic applications such as monitoring inflammatory responses by detecting the presence of leukocytes expressing VLA-4. The compounds of this invention can also be used for isolating or 25 labeling such cells. In addition, as mentioned above, the compounds of the invention can be used to assay for potential inhibitors of VLA-4/VCAM-I interactions.

30 For *in vivo* diagnostic imaging to identify, e.g., sites of inflammation, radioisotopes are typically used in accordance with well known techniques.

-- 144 --

The radioisotopes may be bound to the peptide either directly or indirectly using intermediate functional groups. For instance, chelating agents such as diethylenetriaminepentacetic acid (DTPA) and ethylenediaminetetraacetic acid (EDTA) and similar molecules have been used to bind proteins to metallic ion radioisotopes.

The complexes can also be labeled with a paramagnetic isotope for purposes of *in vivo* diagnosis, as in magnetic resonance imaging (MRI) or electron spin resonance (ESR), both of which are well known. In general, any conventional method for visualizing diagnostic imaging can be used. Usually gamma- and positron-emitting radioisotopes are used for camera imaging and paramagnetic isotopes are used for MRI. Thus, the compounds can be used to monitor the course of amelioration of an inflammatory response in an individual. By measuring the increase or decrease in lymphocytes expressing VLA-4 it is possible to determine whether a particular therapeutic regimen aimed at ameliorating the disease is effective.

The pharmaceutical compositions of the present invention can be used to block or inhibit cellular adhesion associated with a number of diseases and disorders. For instance, a number of inflammatory disorders are associated with integrins or leukocytes. Treatable disorders include, e.g., transplantation rejection (e.g., allograft rejection), Alzheimer's disease, atherosclerosis, AIDS dementia, diabetes (including acute juvenile onset diabetes), retinitis, cancer metastases, rheumatoid arthritis, acute leukocyte-mediated lung injury (e.g., adult respiratory distress syndrome), asthma, nephritis, and acute and chronic inflammation, including atopic dermatitis, psoriasis, myocardial ischemia, and inflammatory bowel disease (including Crohn's disease and ulcerative colitis). In preferred embodiments the pharmaceutical compositions are used to treat inflammatory brain disorders, such as multiple sclerosis (MS), viral meningitis and encephalitis.

Inflammatory bowel disease is a collective term for two similar diseases referred to as Crohn's disease and ulcerative colitis. Crohn's disease is an idiopathic, chronic ulceroinflammatory disease characterized by sharply delimited and typically transmural involvement of all layers of the  
5 bowel wall by a granulomatous inflammatory reaction. Any segment of the gastrointestinal tract, from the mouth to the anus, may be involved, although the disease most commonly affects the terminal ileum and/or colon. Ulcerative colitis is an inflammatory response limited largely to the colonic mucosa and submucosa. Lymphocytes and macrophages are numerous in  
10 lesions of inflammatory bowel disease and may contribute to inflammatory injury.

Asthma is a disease characterized by increased responsiveness of the tracheobronchial tree to various stimuli potentiating paroxysmal constriction  
15 of the bronchial airways. The stimuli cause release of various mediators of inflammation from IgE-coated mast cells including histamine, eosinophilic and neutrophilic chemotactic factors, leukotrienes, prostaglandin and platelet activating factor. Release of these factors recruits basophils, eosinophils and neutrophils, which cause inflammatory injury.

20 Atherosclerosis is a disease of arteries (e.g., coronary, carotid, aorta and iliac). The basic lesion, the atheroma, consists of a raised focal plaque within the intima, having a core of lipid and a covering fibrous cap. Atheromas compromise arterial blood flow and weaken affected arteries. Myocardial  
25 and cerebral infarcts are a major consequence of this disease. Macrophages and leukocytes are recruited to atheromas and contribute to inflammatory injury.

30 Rheumatoid arthritis is a chronic, relapsing inflammatory disease that primarily causes impairment and destruction of joints. Rheumatoid arthritis

usually first affects the small joints of the hands and feet but then may involve the wrists, elbows, ankles and knees. The arthritis results from interaction of synovial cells with leukocytes that infiltrate from the circulation into the synovial lining of the joints. See *e.g.*, Paul, *Immunology* (3d ed.,  
5 Raven Press, 1993).

Another indication for the compounds of this invention is in treatment of organ or graft rejection mediated by VLA-4. Over recent years there has been a considerable improvement in the efficiency of surgical techniques for  
10 transplanting tissues and organs such as skin, kidney, liver, heart, lung, pancreas and bone marrow. Perhaps the principal outstanding problem is the lack of satisfactory agents for inducing immunotolerance in the recipient to the transplanted allograft or organ. When allogeneic cells or organs are transplanted into a host (i.e., the donor and donee are different individuals  
15 from the same species), the host immune system is likely to mount an immune response to foreign antigens in the transplant (host-versus-graft disease) leading to destruction of the transplanted tissue. CD8<sup>+</sup> cells, CD4 cells and monocytes are all involved in the rejection of transplant tissues. Compounds of this invention which bind to alpha-4 integrin are useful, *inter*  
20 *alia*, to block alloantigen-induced immune responses in the donee thereby preventing such cells from participating in the destruction of the transplanted tissue or organ. See, *e.g.*, Paul et al., *Transplant International* 9, 420-425 (1996); Georczynski et al., *Immunology* 87, 573-580 (1996); Georczynski et al., *Transplant. Immunol.* 3, 55-61 (1995); Yang et al., *Transplantation* 60,  
25 71-76 (1995); Anderson et al., *APMIS* 102, 23-27 (1994).

A related use for compounds of this invention which bind to VLA-4 is in modulating the immune response involved in "graft versus host" disease (GVHD). See *e.g.*, Schlegel et al., *J. Immunol.* 155, 3856-3865 (1995).  
30 GVHD is a potentially fatal disease that occurs when immunologically



competent cells are transferred to an allogeneic recipient. In this situation, the donor's immunocompetent cells may attack tissues in the recipient. Tissues of the skin, gut epithelia and liver are frequent targets and may be destroyed during the course of GVHD. The disease presents an especially  
5 severe problem when immune tissue is being transplanted, such as in bone marrow transplantation; but less severe GVHD has also been reported in other cases as well, including heart and liver transplants. The therapeutic agents of the present invention are used, *inter alia*, to block activation of the donor T-cells thereby interfering with their ability to lyse target cells in the  
10 host.

A further use of the compounds of this invention is inhibiting tumor metastasis. Several tumor cells have been reported to express VLA-4 and compounds which bind VLA-4 block adhesion of such cells to endothelial  
15 cells. Steinback et al., *Urol. Res.* 23, 175-83 (1995); Orosz et al., *Int. J. Cancer* 60, 867-71 (1995); Freedman et al., *Leuk. Lymphoma* 13, 47-52 (1994); Okahara et al., *Cancer Res.* 54, 3233-6 (1994).

A further use of the compounds of this invention is in treating multiple  
20 sclerosis. Multiple sclerosis is a progressive neurological autoimmune disease that affects an estimated 250,000 to 350,000 people in the United States. Multiple sclerosis is thought to be the result of a specific autoimmune reaction in which certain leukocytes attack and initiate the destruction of myelin, the insulating sheath covering nerve fibers. In an animal model for  
25 multiple sclerosis, murine monoclonal antibodies directed against VLA-4 have been shown to block the adhesion of leukocytes to the endothelium, and thus prevent inflammation of the central nervous system and subsequent paralysis in the animals<sup>16</sup>.

Pharmaceutical compositions of the invention are suitable for use in a variety of drug delivery systems. Suitable formulations for use in the present invention are found in *Remington's Pharmaceutical Sciences*, Mace Publishing Company, Philadelphia, PA, 17th ed. (1985).

5

In order to enhance serum half-life, the compounds may be encapsulated, introduced into the lumen of liposomes, prepared as a colloid, or other conventional techniques may be employed which provide an extended serum half-life of the compounds. A variety of methods are available for preparing liposomes, as described in, e.g., Szoka, et al., U.S. Patent Nos. 4,235,871, 4,501,728 and 4,837,028 each of which is incorporated herein by reference.

10

The amount administered to the patient will vary depending upon what is being administered, the purpose of the administration, such as prophylaxis or therapy, the state of the patient, the manner of administration, and the like. In therapeutic applications, compositions are administered to a patient already suffering from a disease in an amount sufficient to cure or at least partially arrest the symptoms of the disease and its complications. An amount adequate to accomplish this is defined as "therapeutically effective dose." Amounts effective for this use will depend on the disease condition being treated as well as by the judgment of the attending clinician depending upon factors such as the severity of the inflammation, the age, weight and general condition of the patient, and the like.

15

20

25

The compositions administered to a patient are in the form of pharmaceutical compositions described above. These compositions may be sterilized by conventional sterilization techniques, or may be sterile filtered. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous carrier prior to administration. The pH of the compound

30

preparations typically will be between 3 and 11, more preferably from 5 to 9 and most preferably from 7 to 8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of pharmaceutical salts.

5

The therapeutic dosage of the compounds of the present invention will vary according to, for example, the particular use for which the treatment is made, the manner of administration of the compound, the health and condition of the patient, and the judgment of the prescribing physician. For example, for intravenous administration, the dose will typically be in the range of about 20  $\mu\text{g}$  to about 500  $\mu\text{g}$  per kilogram body weight, preferably about 100  $\mu\text{g}$  to about 300  $\mu\text{g}$  per kilogram body weight. Suitable dosage ranges for intranasal administration are generally about 0.1  $\mu\text{g}$  to 1  $\text{mg}$  per kilogram body weight. Effective doses can be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

15

Compounds of this invention are also capable of binding or antagonizing the actions of  $\alpha_6\beta_1$ ,  $\alpha_9\beta_1$ ,  $\alpha_4\beta_7$ ,  $\alpha_d\beta_2$ ,  $\alpha_e\beta_7$  integrins (although  $\alpha_4\beta_1$  and  $\alpha_9\beta_1$  are preferred in this invention). Accordingly, compounds of this invention are also useful for preventing or reversing the symptoms, disorders or diseases induced by the binding of these integrins to their respective ligands.

20

For example, International Publication Number WO 98/53817, published December 3, 1998 (the disclosure of which is incorporated herein by reference in its entirety) and references cited therein describe disorders mediated by  $\alpha_4\beta_7$ . This reference also describes an assay for determining antagonism of  $\alpha_4\beta_7$  dependent binding to VCAM-Ig fusion protein.

25

Additionally, compounds that bind  $\alpha_d\beta_2$  and  $\alpha_e\beta_7$  integrins are particularly useful for the treatment of asthma and related lung diseases. See,

30

-- 150 --

for example, M. H. Grayson et al., *J. Exp. Med.* **1998**, 188(11) 2187-2191. Compounds that bind  $\alpha_c\beta_7$  integrin are also useful for the treatment of systemic lupus erythematosus (see, for example, M. Pang et al., *Arthritis Rheum.* **1998**, 41(8), 1456-1463); Crohn's disease, ulcerative colitis and  
5 inflammatory bowel disease (IBD) (see, for example, D. Elewaut et al., *Scand J. Gastroenterol* **1998**, 33(7) 743-748); Sjogren's syndrome (see, for example, U. Kroneld et al., *Scand J. Gastroenterol* **1998**, 27(3), 215-218); and rheumatoid arthritis (see, for example, *Scand J. Gastroenterol* **1996**, 44(3), 293-298). And compounds that bind  $\alpha_c\beta_1$  may be useful in preventing  
10 fertilization (see, for example, H. Chen et al., *Chem. Biol.* **1999**, 6, 1-10).

Certain of the compounds within the generic formulas described herein are also useful as synthetic intermediates for other compounds of this invention as illustrated in the examples herein.

15

The following synthetic and biological examples are offered to illustrate this invention and are not to be construed in any way as limiting the scope of this invention. Unless otherwise stated, all temperatures are in degrees Celsius.

20

### EXAMPLES

In the examples below, the following abbreviations have the following meanings. If an abbreviation is not defined, it has its generally accepted meaning.

25	aq or aq.	=	aqueous
	AcOH	=	acetic acid
	bd	=	broad doublet
	bm	=	broad multiplet
	bs	=	broad singlet
30	Bn	=	benzyl
	Boc	=	<i>N</i> - <i>tert</i> -butoxycarbonyl
	Boc <sub>2</sub> O	=	di- <i>tert</i> -butyl dicarbonate

-- 151 --

	BOP	=	benzotriazol-1-yloxy- tris(dimethylamino)phosphonium hexafluorophosphate
	Cbz	=	carbobenzyloxy
5	CHCl <sub>3</sub>	=	chloroform
	CH <sub>2</sub> Cl <sub>2</sub>	=	dichloromethane
	(COCl) <sub>2</sub>	=	oxalyl chloride
	d	=	doublet
	dd	=	doublet of doublets
10	dt	=	doublet of triplets
	DBU	=	1,8-diazabicyclo[5.4.0]undec-7-ene
	DCC	=	1,3-dicyclohexylcarbodiimide
	DMAP	=	4- <i>N,N</i> -dimethylaminopyridine
	DME	=	ethylene glycol dimethyl ether
15	DMF	=	<i>N,N</i> -dimethylformamide
	DMSO	=	dimethylsulfoxide
	EDC	=	1-(3-dimethylaminopropyl)-3- ethylcarbodiimide hydrochloride
	Et <sub>3</sub> N	=	triethylamine
20	Et <sub>2</sub> O	=	diethyl ether
	EtOAc	=	ethyl acetate
	EtOH	=	ethanol
	eq or eq.	=	equivalent
	Fmoc	=	<i>N</i> -(9-fluorenylmethoxycarbonyl)
25	FmocONSu	=	<i>N</i> -(9-fluorenylmethoxycarbonyl)- succinimide
	g	=	grams
	h	=	hour
	H <sub>2</sub> O	=	water
30	HBr	=	hydrobromic acid
	HCl	=	hydrochloric acid
	HOBT	=	1-hydroxybenzotriazole hydrate
	hr	=	hour
	K <sub>2</sub> CO <sub>3</sub>	=	potassium carbonate
35	L	=	liter
	m	=	multiplet
	MeOH	=	methanol
	mg	=	milligram
	MgSO <sub>4</sub>	=	magnesium sulfate
40	mL	=	milliliter
	mm	=	millimeter
	mM	=	millimolar
	mmol	=	millimol
	mp	=	melting point
45	N	=	normal

-- 152 --

	NaCl	=	sodium chloride
	Na <sub>2</sub> CO <sub>3</sub>	=	sodium carbonate
	NaHCO <sub>3</sub>	=	sodium bicarbonate
	NaOEt	=	sodium ethoxide
5	NaOH	=	sodium hydroxide
	NH <sub>4</sub> Cl	=	ammonium chloride
	NMM	=	<i>N</i> -methylmorpholine
	Phe	=	L-phenylalanine
	Pro	=	L-proline
10	psi	=	pounds per square inch
	PtO <sub>2</sub>	=	platinum oxide
	q	=	quartet
	quint.	=	quintet
	rt	=	room temperature
15	s	=	singlet
	sat	=	saturated
	t	=	triplet
	<i>t</i> -BuOH	=	<i>tert</i> -butanol
	TFA	=	trifluoroacetic acid
20	THF	=	tetrahydrofuran
	TLC or tlc	=	thin layer chromatography
	Ts	=	tosyl
	TsCl	=	tosyl chloride
	TsOH	=	tosylate
25	μL	=	microliter

The following Methods may be used to prepare the compounds of this invention.

30

#### Method A

##### Methyl Ester Preparation Procedure

Amino acid methyl esters can be prepared using the method of Brenner and Huber *Helv. Chim. Acta* 1953, 36, 1109.

35

-- 153 --

## Method B

BOP Coupling Procedure

The desired dipeptide ester was prepared by the reaction of a carboxylic acid (1 equivalent) with the appropriate amino acid ester or amino acid ester hydrochloride (1 equivalent), benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate [BOP] (2.0 equivalent), triethylamine (1.1 equivalent), and DMF. The reaction mixture was stirred at room temperature overnight. The crude product is purified flash chromatography to afford the dipeptide ester.

10

## Method C

Hydrogenation Procedure I

Hydrogenation was performed using 10% palladium on carbon (10% by weight) in methanol at 30 psi overnight. The mixture was filtered through a pad of Celite and the filtrate concentrated to yield the desired compound.

15

## Method D

Hydrolysis Procedure I

To a chilled (0°C) THF/H<sub>2</sub>O solution (2:1, 5 - 10 mL) of the appropriate ester was added LiOH (or NaOH) (0.95 equivalents). The temperature was maintained at 0°C and the reaction was complete in 1-3 hours. The reaction mixture was extracted with ethyl acetate and the aqueous phase was lyophilized resulting in the desired carboxylate salt.

20

25

## Method E

Ester Hydrolysis Procedure II

To a chilled (0°C) THF/H<sub>2</sub>O solution (2:1, 5 - 10 mL) of the appropriate ester was added LiOH (1.1 equivalents). The temperature was maintained at 0°C and the reaction was complete in 1-3 hours. The reaction mixture was concentrated and the residue was taken up into H<sub>2</sub>O and the pH adjusted to 2-

30

-- 154 --

3 with aqueous HCl. The product was extracted with ethyl acetate and the combined organic phase was washed with brine, dried over  $\text{MgSO}_4$ , filtered and concentrated to yield the desired acid.

5

#### Method F

##### Ester Hydrolysis Procedure III

The appropriate ester was dissolved in dioxane/ $\text{H}_2\text{O}$  (1:1) and 0.9 equivalents of 0.5 N NaOH was added. The reaction was stirred for 3-16 hours and then concentrated. The resulting residue was dissolved in  $\text{H}_2\text{O}$  and  
10 extracted with ethyl acetate. The aqueous phase was lyophilized to yield the desired carboxylate sodium salt.

#### Method G

##### BOC Removal Procedure

15 Anhydrous hydrochloride (HCl) gas was bubbled through a methanolic solution of the appropriate Boc-amino acid ester at  $0^\circ\text{C}$  for 15 minutes and the reaction mixture was stirred for three hours. The solution was concentrated to a syrup and dissolved in  $\text{Et}_2\text{O}$  and reconcentrated. This procedure was repeated and the resulting solid was placed under high vacuum  
20 overnight.

#### Method H

##### *tert*-Butyl Ester Hydrolysis Procedure I

25 The *tert*-butyl ester was dissolved in  $\text{CH}_2\text{Cl}_2$  and treated with TFA. The reaction was complete in 1-3 hr at which time the reaction mixture was concentrated and the residue dissolved in  $\text{H}_2\text{O}$  and lyophilized to yield the desired acid.

30



-- 155 --

## Method I

EDC Coupling Procedure I

To a  $\text{CH}_2\text{Cl}_2$  solution (5-20 mL) of a carboxylic acid (1 equivalent), the appropriate amino acid ester hydrochloride (1 equivalent), N-methylmorpholine (1.1-2.2 equivalents) and 1-hydroxybenzotriazole (2 equivalents) were mixed, placed in an ice bath and 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide (1.1 equivalents) added. The reaction was allowed to rise to room temperature and stirred overnight. The reaction mixture was poured into  $\text{H}_2\text{O}$  and the organic phase was washed with sat.  $\text{NaHCO}_3$ , brine, dried ( $\text{MgSO}_4$  or  $\text{Na}_2\text{SO}_4$ ), filtered and concentrated. The crude product was purified by column chromatography.

## Method J

EDC Coupling Procedure II

To a DMF solution (5-20 mL) of a carboxylic acid (1 equivalent), the appropriated amino acid ester hydrochloride (1 equivalent),  $\text{Et}_3\text{N}$  (1.1 equivalents) and 1-hydroxybenzotriazole (2 equivalents) were mixed, placed in an ice bath and 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide (1.1 equivalents) added. The reaction was allowed to rise to room temperature and stirred overnight. The reaction mixture was partitioned between  $\text{EtOAc}$  and  $\text{H}_2\text{O}$  and the organic phase washed with 0.2 N citric acid,  $\text{H}_2\text{O}$ , sat.  $\text{NaHCO}_3$ , brine, dried ( $\text{MgSO}_4$  or  $\text{Na}_2\text{SO}_4$ ), filtered and concentrated. The crude product was purified by column chromatography or preparative TLC.

## Method K

*tert*-Butyl Ester Hydrolysis Procedure II

The *tert*-butyl ester was dissolved in  $\text{CH}_2\text{Cl}_2$  (5 mL) and treated with TFA (5 mL). The reaction was complete in 1-3 hours at which time the reaction mixture was concentrated and the residue dissolved in  $\text{H}_2\text{O}$  and

-- 156 --

concentrated. The residue was redissolved in H<sub>2</sub>O and lyophilized to yield the desired product.

#### Method L

5

##### Carbamate Formation Procedure I

Into a reaction vial were combined 15.2 mmol, 1.0 eq. of the starting hydroxy compound (typically a tyrosine derivative) and 1.86 g (15.2 mmol, 1.0 eq) DMAP. Methylene chloride (50 mL), triethylamine (2.12 mL, 1.54 g, 15.2 mmol, 1.0 eq), and dimethylcarbamyl chloride (1.68 mL, 1.96 g, 18.2 mmol, 1.2 eq) were then added. The vial was capped tightly, and the reaction solution swirled to obtain a homogeneous solution. The reaction solution was then heated to 40°C. After 48 h, TLC of the resulting colorless solution indicated complete conversion. The work-up of the reaction solution was as follows: 50 mL EtOAc and 50 mL hexanes was added to the reaction mixture, and the resulting mixture was washed with 0.5 M citric acid (3 x 50 mL), water (2 x 50 mL), 10% K<sub>2</sub>CO<sub>3</sub> (2 x 50 mL), and sat. NaCl (1 x 50 mL); dried with MgSO<sub>4</sub>, filtered and evaporated to afford the desired compound.

20

#### Method M

##### Carbamate Formation Procedure II

Into a reaction vial were combined 84.34 mmol (1.0 eq) of the starting hydroxy compound (typically a tyrosine derivative) and 17.0 g (84.34 mmol, 1.0 eq) 4-nitrophenyl chloroformate. Methylene chloride (700 mL) was added and the vial was capped with a septum. A nitrogen line was attached and the vial was immersed in a 4:1 water/ethanol dry ice slurry with stirring to cool to -15°C. Triethylamine (29.38 mL, 21.33 g, 210.81 mmol, 2.5 eq) was added over five minutes with stirring and the stirring was continued at -10 to -15°C for 1 h. *N*-Methyl piperazine (9.35 mL, 8.45 g, 84.34 mmol, 1.0 eq) was added over three minutes with stirring and stirring was continued

-- 157 --

overnight while warming to room temperature. The reaction mixture was diluted with 700 mL hexanes and the resulting mixture was washed repeatedly with 10%  $K_2CO_3$ , until no yellow color (from 4-nitrophenol) is observed in the aqueous layer. The mixture was then washed with sat. NaCl, dried over anhydrous  $MgSO_4$ , filtered and evaporated. The residue was dissolved in 500 mL of ethanol and evaporated to remove triethylamine. The residue was again dissolved in 500 mL of ethanol and evaporated to remove triethylamine. The residue was then dissolved in 400 mL of ethanol and 600 mL of water was added with stirring to precipitate a solid or oil. If an oil formed, the oil is stirred vigorously to induce it to solidify. The solid is then isolated by filtration. Dissolution, precipitation, and filtration are repeated once and the resulting solid is rinsed with water to remove traces of yellow color. The solid is then subjected to high vacuum until the mass remains constant thereby affording the desired carbamoyloxy compound.

15

#### Method N

##### Preparation of 5-Iodo-4(3H)-pyrimidinone

The procedure of Sakamoto *et. al.* (Chem. Pharm. Bull. 1986, 34(7), 2719-2724) was used to convert 4(3H)-pyrimidinone into 5-iodo-4(3H)-pyrimidinone, which was of sufficient purity for conversion to 4-chloro-5-iodopyrimidine.

20

#### Method O

##### Preparation of 4-Chloro-5-iodopyrimidine

5-Iodo-4(3H)-pyrimidinone (1 eq.) was suspended in toluene to which was added  $POCl_3$  (2.0 eq.). The reaction mixture was heated to reflux for 3 hours, and then cooled and concentrated. The residue was suspended in water, adjusted to pH=7 by addition of 4N sodium hydroxide, and extracted with ethyl acetate. The organic extracts were washed with brine, dried (MgSO<sub>4</sub>), filtered and stripped to give a red oil. The crude product was

25

30

-- 158 --

dissolved in methanol and silica gel was added. Following concentration, the coated silica gel was loaded onto a plug of silica gel and elution with ethyl acetate/hexanes yielded the title compound.

5

#### Method P

##### Preparation of

##### N-(5-Iodopyrimidin-4-yl)-L-4-(N,N-dimethylcarbamyloxy)phenylalanine tert-Butyl Ester

A solution 4-chloro-5-iodopyrimidine (1.0 eq.), L-4-(N,N-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester (1.0 eq), and N,N-diisopropylethyl amine (2.0 eq) in tetrahydrofuran was heated at reflux for 16 hours. The reaction mixture was then cooled and diluted with water and ethyl acetate. The organic phase was washed with 0.2 N citric acid, water, saturated NaHCO<sub>3</sub>, brine, dried (MgSO<sub>4</sub>), filtered and concentrated. The residue was purified by silica gel chromatography using ethyl acetate/hexanes to afford the title compound.

#### Method Q

##### Suzuki Coupling Procedure I

To an ethyleneglycol dimethyl ether solution of tetrakis(triphenylphosphine)palladium (0.04 eq) was added N-(5-iodopyrimidin-4-yl)-L-4-(N,N-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester (1.0 eq.). After stirring for approximately ten minutes a boronic acid or ester (1.2 eq) and 2M Na<sub>2</sub>CO<sub>3</sub> (2.0 eq) were added, and the reaction flask was evacuated and then flushed with nitrogen gas. The reaction was heated at reflux from three to sixteen hours. The reaction mixture was then cooled, diluted with water and ethyl acetate, and the organic phase was washed with 0.2 N citric acid, water, saturated NaHCO<sub>3</sub>, brine, dried (MgSO<sub>4</sub>), filtered and concentrated. Alternatively, the cooled reaction mixture was diluted with ethyl acetate and washed with water, saturated NaHCO<sub>3</sub>, dried (MgSO<sub>4</sub>),

-- 159 --

filtered and concentrated. Either column chromatography or preparative thin layer chromatography on silica gel using ethyl acetate/hexanes afforded the desired product.

5

## Method R

Suzuki Coupling Procedure II

To a dimethylformamide solution of tetrakis(triphenylphosphine)-palladium (0.02 - 0.05 eq) was added *N*-(5-iodopyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester (1.0 eq.). After stirring  
10 for approximately ten minutes, the boronic acid (1.1 - 4.0 eq) and  $K_3PO_4$  (1.5 - 2.0 eq) were added, and the reaction was heated at 100°C for three to sixteen hours. The reaction mixture was then cooled, diluted with water and ethyl acetate, and the organic phase was washed with 0.2 N citric acid, water, saturated  $NaHCO_3$ , brine, dried ( $MgSO_4$ ), filtered and concentrated. Either  
15 column chromatography or preparative thin layer chromatography on silica gel using ethyl acetate/hexanes afforded the desired product.

## Method S

Suzuki Coupling Procedure III

20 An ethyleneglycol dimethyl ether/2M  $Na_2CO_3$  (1:1 by volume) solution of tetrakis(triphenylphosphine)palladium (0.04 eq), *N*-(5-iodopyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester (1.0 eq.), the boronic acid (1.1 eq) and lithium chloride (3.0 eq ) was heated to reflux for approximately six hours. The cooled reaction mixture was diluted with ethyl  
25 acetate and washed with water, brine, dried ( $MgSO_4$ ), filtered and concentrated. The residue was purified by silica gel column chromatography using ethyl acetate/hexanes to afford the desired product.

30

-- 160 --

## Method T

Suzuki Coupling Procedure IV

An ethyleneglycol dimethyl ether/2M Na<sub>2</sub>CO<sub>3</sub>, (1:1 by volume) solution of tetrakis(triphenylphosphine)palladium (0.05 eq), *N*-(5-iodopyrimidin-4-yl)-  
5 L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester (1.0 eq.), the boronic acid (1.5 eq) and tri-*o*-tolylphosphine (0.1 eq) was heated to reflux for approximately three hours. The cooled reaction mixture was diluted with ethyl acetate and water and washed with water, brine, dried (MgSO<sub>4</sub>), filtered and concentrated. The residue was purified by preparative thin layer  
10 chromatography on silica gel using ethyl acetate/hexanes to afford the desired product.

## Method U

Heck Reaction Procedure I

15 A dimethylformamide solution of *N*-(5-iodopyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester (1.0 eq.), *N,N*-dimethylacrylamide (2.0 eq), and triethylamine (6.0 eq) was degassed with nitrogen and then dichlorobis-(triphenylphosphine)palladium was added. The reaction was warmed to 90°C under a stream of nitrogen for 16 hours. The  
20 cooled reaction mixture was diluted with ethyl acetate and water and washed with water, brine, dried (MgSO<sub>4</sub>), filtered and concentrated. The residue was purified by column chromatography on silica gel using ethyl acetate/hexanes followed by preparative thin layer chromatography on silica gel using ethyl acetate/hexanes to afford the desired product.

25

## Method V

Hydrogenation Procedure II

*N*-(5-(2-*N,N*-dimethylcarbamylethyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester was dissolved in ethanol  
30 to which was added 10 % palladium on carbon. The reaction mixture was

-- 161 --

hydrogenated at 35 psi hydrogen for approximately five hours. The reaction mixture was filtered through a pad of Celite, and the filtrate was concentrated. The residue was purified by preparative thin layer chromatography on silica gel using methanol/dichloromethane to afford the  
5 desired product.

#### Method W

##### Heck Reaction Procedure II

To a tetrahydrofuran solution of *N*-(5-iodopyrimidin-4-yl)-*L*-4-(*N,N*-  
10 dimethylcarbamyloxy)phenylalanine *tert*-butyl ester (1.0 eq)  
dichlorobis(triphenylphosphine)palladium, triethylamine (0.05 eq) and  
triphenylphosphine (0.025 eq) was added phenylacetylene (1.5 eq) and  
triethylamine (1.5 eq). After twenty minutes, copper (I) iodide (0.012 eq)  
was added, and the resulting mixture was stirred overnight at room  
15 temperature. The reaction mixture was then diluted with ethyl acetate and  
water and washed with 0.2 N citric acid, water, saturated NaHCO<sub>3</sub>, brine,  
dried (MgSO<sub>4</sub>), filtered and concentrated. The residue was chromatographed  
on a silica gel column using ethyl acetate/hexanes. <sup>1</sup>H NMR analysis showed  
that the desired product to be contaminated with the iodopyrimidine starting  
20 material. However, the product was used without further purification.

#### Method X

##### Hydrogenation Procedure III

Crude *N*-(5-(2-phenylethynyl)pyrimidin-4-yl)-*L*-4-(*N,N*-  
25 dimethylcarbamyloxy)phenylalanine *tert*-butyl ester was dissolved in ethanol  
to which was added 10% palladium on carbon and sodium acetate (3.0 eq).  
The reaction mixture was hydrogenated at 40 psi hydrogen for approximately  
three hours, then filtered through a pad of Celite, and the filtrate  
concentrated. The residue was washed with 0.2 N citric acid, water,  
30 saturated NaHCO<sub>3</sub>, brine, dried (MgSO<sub>4</sub>), filtered and concentrated. Silica

-- 162 --

gel column chromatography using ethyl acetate/hexanes yielded the desired product.

#### Method Y

5

#### Preparation of

#### *N*-(6-Chloropyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-Butyl Ester

A solution 4,6-dichloropyrimidine (1.2 eq), L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester (1.0 eq), and  
10 triethylamine (1.05 eq) in ethanol was heated at reflux for 16 hours. The reaction mixture was cooled and concentrated, and the residue was taken-up in water and ethyl acetate. The organic phase was washed with 0.2 N citric acid, water, saturated NaHCO<sub>3</sub>, brine, dried (MgSO<sub>4</sub>), filtered and concentrated. The residue was purified by silica gel chromatography using  
15 ethyl acetate/hexanes to afford the title compound.

#### Method Z

#### Suzuki Coupling Procedure V

An ethyleneglycol dimethyl ether solution of  
20 tetrakis(triphenylphosphine)palladium (0.12 eq), *N*-(6-chloropyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester (1.0 eq.) and triphenylphosphine (0.05 eq) was stirred for approximately ten minutes. The boronic acid or ester (1.2 - 2.5 eq) and 2M Na<sub>2</sub>CO<sub>3</sub> (2.0 eq) were added, and the reaction was heated at 90°C for 16 to 72 hours. The reaction mixture  
25 was cooled and concentrated, and the residue was taken up in water and ethyl acetate. The organic phase was washed with 0.2 N citric acid, water, saturated NaHCO<sub>3</sub>, brine, dried (MgSO<sub>4</sub>), filtered and concentrated. The residue was purified by preparative thin layer chromatography on silica gel using ethyl acetate/hexanes to afford the desired product.

30



-- 163 --

## Method AA

## Preparation of

*N*-(6-(*N*-Alkylamino)pyrimidin-4-yl)-L-4-(*N,N*-  
dimethylcarbamyloxy)phenylalanine  
*tert*-Butyl Ester

5 A mixture of *N*-(6-chloropyrimidin-4-yl)-L-4-(*N,N*-  
dimethylcarbamyloxy)phenylalanine *tert*-butyl ester (1.0 eq) and an  
alkylamine (10.0 eq) was heated in a sealed tube at 120°C for 16 hours. The  
reaction mixture was cooled and diluted with ethyl acetate. The organic  
10 portion was washed with 0.2 N citric acid, water, saturated NaHCO<sub>3</sub>, brine,  
dried (MgSO<sub>4</sub>), filtered and concentrated. The residue was purified by silica  
gel chromatography using ethyl acetate/hexanes to afford the desired  
compound.

15

## Method BB

Preparation of 4-*N*-Alkylamino-5-bromo-2-chloropyrimidine

A methanol solution of 5-bromo-2,4-dichloropyrimidine (1.0 eq), the  
alkylamine (1.05 eq, typically L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine  
*tert*-butyl ester), and *N,N*-diisopropylethylamine (5.0 eq) was heated to 40°C  
20 for 16 hours. The reaction mixture was then concentrated, and the residue  
was taken up in ethyl acetate. The organic portion was washed with 0.2 N  
citric acid, water, saturated NaHCO<sub>3</sub>, brine, dried (MgSO<sub>4</sub>), filtered and  
concentrated. The crude material was purified by silica gel chromatography  
using ethyl acetate/hexanes to afford the desired compound.

25

## Method CC

Preparation of 4-*N*-Alkylamino-5-bromo-2-*N*-alkylaminopyrimidine

An isopropanol solution of the 4-*N*-alkylamino-5-bromo-2-  
chloropyrimidine (1.0 eq) and an alkylamine (5.0 eq) was heated in sealed  
30 tube at 130°C for 3-5 hours. The reaction mixture was then cooled and  
washed with 0.2 N citric acid, water, saturated NaHCO<sub>3</sub>, brine, dried  
(MgSO<sub>4</sub>), filtered and concentrated. The crude material was purified by

-- 164 --

silica gel chromatography using ethyl acetate/hexanes to afford the desired compound.

#### Method DD

##### 5                    4-N-Alkylamino-5-bromo-2-N-alkylaminopyrimidine

##### Suzuki Coupling Procedure

To an ethyleneglycol dimethyl ether solution of tetrakis(triphenylphosphine)palladium (0.04 eq) was added an 4-N-alkylamino-5-bromo-2-N-alkylaminopyrimidine (1.0 eq.). After stirring for approximately ten minutes, the boronic acid or ester (1.2 eq) and 2M Na<sub>2</sub>CO<sub>3</sub> (2.0 eq) was added, and the reaction flask was evacuated and then flushed with nitrogen gas. The reaction was heated at reflux for three to four hours. The reaction mixture was then cooled and diluted with water and ethyl acetate, and the organic phase was washed with 0.2 N citric acid, water, 15 saturated NaHCO<sub>3</sub>, brine, dried (MgSO<sub>4</sub>), filtered and concentrated. The residue was purified by either silica gel column or preparative thin layer chromatography using ethyl acetate/hexanes to afford the desired product.

#### Method EE

##### 20                    Preparation of

##### N-tert-Butoxycarbonyl-4-Iodo-L-phenylalanine Methyl Ester

The title compound was prepared from 4-iodo-L-phenylalanine by standard conditions described by Bodanszky and Bodanszky in *The Practice of Peptide Synthesis*; Springer-Verlag: Berlin, 1984.

25

30

-- 165 --

## Method FF

## Preparation of

*N*-tert-Butoxycarbonyl-4-(2,6-dimethoxyphenyl)-L-phenylalanineMethyl Ester

5        To a dimethylformamide solution of  
tetrakis(triphenylphosphine)palladium (0.02 - 0.05 eq) was added *N*-tert-  
butoxycarbonyl-4-(2,6-dimethoxyphenyl)-L-phenylalanine methyl ester (1.0  
eq.). After stirring for approximately ten minutes, 2,6-dimethoxyphenyl  
boronic acid (1.1 eq) and K<sub>3</sub>PO<sub>4</sub> (2.0 eq) were added, and the reaction was  
10       heated at 100°C for sixteen hours. The reaction mixture was then cooled,  
diluted with water and ethyl acetate, and the organic phase was washed with  
0.2 N citric acid, water, saturated NaHCO<sub>3</sub>, brine, dried (MgSO<sub>4</sub>), filtered  
and concentrated. Column chromatography on silica gel using ethyl  
acetate/hexanes afforded the desired product.

15

## Method GG

## Preparation of

4-(2,6-Dimethoxyphenyl)-L-phenylalanine Methyl EsterTrifluoroacetic Acid Salt

20       A methylene chloride solution of *N*-tert-butoxycarbonyl-4-(2,6-  
dimethoxyphenyl)-L-phenylalanine methyl ester was treated with  
trifluoroacetic acid for six hours at room temperature. Concentration of the  
volatiles yielded the title compound.

25

## Method HH

*tert*-Butyl Ester Cleavage Procedure III

A methylene chloride solution of the appropriate *tert*-butyl ester was  
treated with trifluoroacetic acid at room temperature. After 2-3 hours the  
volatiles were evaporated, and the residue was treated again with methylene

-- 166 --

chloride and trifluoroacetic acid. After 2-3 hours the volatiles were evaporated again to yield the desired compound.

#### Method II

##### Preparation of

*N*-(5-Allylpyrimidin-4-yl)-L-4-(*N,N*-dimethyl-carbamoyloxy)phenylalanine

##### *tert*-Butyl Ester

*N*-(5-Iodo-pyrimidin-4-yl)-L-4-(*N,N*-dimethyl-carbamoyloxy)phenylalanine *tert*-butyl ester (1.0 eq) was dissolved in dry DMF, with allyltributylstannane (1.1 eq), bis(triphenylphosphine)palladium dichloride (0.03 eq) and LiCl (3.0eq). The reaction mixture was flushed under nitrogen, and heated to 90°C for 2 hours. EtOAc was added, and the organic layer was washed with water and brine, and dried over MgSO<sub>4</sub>. After filtration and evaporation of the solvent under reduced pressure, the crude material was purified by column chromatography (silica gel) eluting with EtOAc/hexanes 1:3. The title material was isolated in good yields.

#### Method JJ

##### Preparation of

*N*-[5-propylpyrimidin-4-yl]-L-4-(*N,N*-dimethyl-carbamoyloxy)phenylalanine

##### *tert*-Butyl Ester

*N*-(5-Allylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester was dissolved in methanol and treated with a catalytic amount of 10% palladium on carbon. The mixture was shaken under 10 psi hydrogen gas for 3 hours. Upon filtration through a pad of Celite, and evaporation of the solvent under reduced pressure, the desired material was isolated as a foam.

-- 167 --

## Method KK

## Preparation of

*N*-(5-propylpyrimidin-4-yl)-L-4-(*N,N*-dimethyl-carbamoyloxy)phenylalanine

*N*-(5-Propylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)-

5 phenylalanine *tert*-butyl ester was treated with neat trifluoroacetic acid, and the mixture was stirred for 5 h at room temperature. Upon evaporation of the solvent under reduced pressure, the desired material was isolated as a foam.

## Method LL

10 Preparation of Dimethyl 2-Alkylmalonate

To a suspension of sodium hydride 60% dispersion in mineral oil (1.1 eq) in anhydrous THF was added slowly with stirring dimethyl malonate (1.1 eq), causing the evolution of gas. To the resulting solution was added a bromoalkane, iodoalkane, or trifluoromethanesulfonyloxyalkane (1.0 eq), and  
15 the mixture was heated to 50°C for 48 h, at which point TLC indicated consumption of the bromoalkane, iodoalkane, or trifluoromethanesulfonyloxyalkane. The mixture was diluted with diethyl ether and washed with 70% saturated sodium chloride. The organic extracts were treated with anhydrous magnesium sulfate, filtered, and evaporated to  
20 afford a dimethyl 2-alkylmalonate of sufficient purity for immediate conversion to a 5-alkyl-4,6-dihydropyrimidine.

## Method MM

Preparation of Diethyl 2-Alkylidenylmalonate

25 Procedure B (p. 2759) of Houve and Winberg (J. Org. Chem. 1980, 45(14), 2754-2763) was employed to react diethyl malonate with a ketone or an aldehyde to afford a diethyl 2-alkylidenylmalonate of sufficient purity for immediate conversion to a diethyl 2-alkylmalonate.

30

-- 168 --

## Method NN

Preparation of Diethyl 2-Alkylmalonate

A diethyl 2-alkylidenylmalonate and an equal mass 10% palladium on carbon were suspended in ethanol. The mixture was shaken under 55 psi hydrogen gas for 24 h, at which point TLC indicated consumption of the diethyl 2-alkylidenylmalonate. The mixture was filtered through Celite and evaporated to afford a diethyl 2-alkylmalonate of sufficient purity for immediate conversion to a 5-alkyl-4,6-dihydroxypyrimidine.

10

## Method OO

Preparation of 5-Alkyl-4,6-dihydroxypyrimidine

To a diethyl 2-alkylmalonate or a dimethyl 2-alkylmalonate (1.0 eq) was added formamidine acetate (1.0 eq) and 25% sodium methoxide in methanol (3.3 eq). The resulting slurry was stirred vigorously and heated to 60°C for 4 h, and then allowed to cool. The slurry was diluted with water, and acidified to pH = 2 by addition of HCl. The resulting precipitate was collected by filtration, washed with water, and dried under vacuum, to afford a 5-alkyl-4,6-dihydroxypyrimidine of sufficient purity for immediate conversion to a 5-alkyl-4,6-dichloropyrimidine.

20

## Method PP

Preparation of 5-Alkoxy-4-hydroxypyrimidine

The method (p. 308) of Anderson et al. (Org. Proc. Res. Devel. 1997, 1, 300-310) was employed to react a methyl alkoxyacetate, sodium methoxide, ethyl formate, and formamidine acetate to afford a 5-alkoxy-4-hydroxypyrimidine of sufficient purity for immediate conversion to a 5-alkoxy-4-chloropyrimidine.

30

-- 169 --

## Method QQ

## Preparation of

5-Alkyl-4,6-dichloropyrimidine or 5-Alkoxy-4-chloropyrimidine

To a 5-alkyl-4,6-dihydroxypyrimidine or a 5-alkoxy-4-hydroxypyrimidine (1.0 eq) were added phosphorus oxychloride (15.0 eq) and *N,N*-dimethylaniline (1.0 eq), and the mixture was heated to 100°C for 3 h, and then allowed to cool. The resulting solution was poured onto ice, and the mixture was extracted with dichloromethane. The organic extracts were treated with anhydrous magnesium sulfate, filtered, and evaporated to afford a 5-alkyl-4,6-dichloropyrimidine or a 5-alkoxy-4-chloropyrimidine of sufficient purity for immediate conversion to a 5-alkyl-4-*N*-alkylamino-6-chloropyrimidine or a 5-alkoxy-4-*N*-alkylaminopyrimidine.

## Method RR

## Preparation of

5-Alkyl-4-*N*-alkylamino-6-chloropyrimidine or  
5-Alkoxy-4-*N*-alkylaminopyrimidine

To a solution of a 5-alkyl-4,6-dichloropyrimidine or a 5-alkoxy-4-chloropyrimidine (1.0 eq) in ethanol were added an alkyl amine (1.2 eq, typically L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester) and diisopropylethylamine (2.0 eq). The mixture was sealed in a pressure tube and heated to 120°C for 48 h, at which point TLC indicated consumption of the 5-alkyl-4,6-dichloropyrimidine or the 5-alkoxy-4-chloropyrimidine. The mixture was evaporated, and the residue was partitioned between ethyl acetate and pH = 4.5 citrate buffer. The organic extracts were washed with saturated sodium chloride, treated with anhydrous magnesium sulfate, filtered, and evaporated. The residue was purified by chromatography on silica gel using ethyl acetate and hexanes to afford a pure 5-alkyl-4-*N*-alkylamino-6-chloropyrimidine or 5-alkoxy-4-*N*-alkylaminopyrimidine.

-- 170 --

## Method SS

Preparation of 5-Alkyl-4-N-alkylaminopyrimidine (Procedure I)

A suspension of 5-alkyl-4-N-alkylamino-6-chloropyrimidine (1.0 eq), and an equal mass 10% palladium on carbon, and sodium bicarbonate (5.0 eq) in methanol was shaken under 55 psi hydrogen gas for 16 h, at which point TLC indicated consumption of the 5-alkyl-4-N-alkylamino-6-chloropyrimidine. The mixture was filtered through Celite and evaporated to give a residue, which was partitioned between ethyl acetate and 70% saturated sodium chloride. The organic extracts were treated with anhydrous magnesium sulfate, filtered, and evaporated. The residue was purified by chromatography on silica gel using ethyl acetate and hexanes to afford a pure 5-alkyl-4-N-alkylaminopyrimidine.

## Method TT

Preparation of 5-Alkyl-4-N-alkylaminopyrimidine (Procedure II)

A suspension of 5-alkyl-4-N-alkylamino-6-chloropyrimidine (1.0 eq), sodium acetate (10.0 eq), and zinc powder (20.0 eq) in a 9:1 mixture of acetic acid and water was stirred vigorously at 40°C for 72 h, at which point TLC indicated partial consumption of the 5-alkyl-4-N-alkylamino-6-chloropyrimidine. The supernatant solution was decanted from remaining zinc and evaporated. The residue was partitioned between ethyl acetate and saturated sodium bicarbonate, and the organic extracts were treated with anhydrous magnesium sulfate, filtered, and evaporated. The residue was purified by chromatography on silica gel using ethyl acetate and hexanes to afford a pure 5-alkyl-4-N-alkylaminopyrimidine.



-- 171 --

## Method UU

## Preparation of

*N*-Benzyloxycarbonyl-L-Tyrosine *tert*-Butyl Ester

To a 0°C suspension of L-tyrosine *tert*-butyl ester (Bachem, 1.0 eq) and  
5 sodium bicarbonate (2.0 eq) in a 1:1 mixture of THF and water was added  
slowly with stirring benzyl chloroformate (1.1 eq). After the addition, the  
mixture was stirred at 0°C for 3 h and at room temperature for 24 h. The  
mixture was diluted with diethyl ether, and the aqueous layer was separated.  
The organic extracts were washed with saturated sodium chloride, treated  
10 with anhydrous magnesium sulfate, filtered, and evaporated to afford *N*-  
benzyloxycarbonyl-L-tyrosine *tert*-butyl ester of sufficient purity for  
immediate conversion of the tyrosine hydroxyl into a carbamate.

## Method VV

## 15 Preparation of

*N*-Benzyloxycarbonyl-L-4-(*N,N*-Dimethylcarbamyloxy)phenylalanine  
*tert*-Butyl Ester

A mixture of *N*-benzyloxycarbonyl-L-tyrosine *tert*-butyl ester (1.0 eq),  
4-dimethylaminopyridine (1.0 eq), triethylamine (1.5 eq),  
20 dimethylcarbamyloxychloride (1.2 eq), and dichloromethane was heated to 37°C  
for 16 h. The mixture was diluted with additional dichloromethane and  
washed sequentially with 1.0 M potassium bisulfate, water, saturated sodium  
bicarbonate, and saturated sodium chloride. The organic extracts were  
treated with anhydrous magnesium sulfate, filtered, and evaporated to afford  
25 *N*-benzyloxycarbonyl-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-  
butyl ester as a white solid of sufficient purity for immediate conversion to L-  
4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester.

-- 172 --

## Method WW

## Preparation of

L-4-(N,N-Dimethylcarbamyloxy)phenylalanine *tert*-Butyl Ester

A suspension of *N*-benzyloxycarbonyl-L-4-(*N,N*-  
5 dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and an equal mass of  
10% palladium on carbon in methanol was shaken under 55 psi hydrogen gas  
for 1 h, at which point TLC indicated consumption of the *N*-  
benzyloxycarbonyl-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl  
ester. The mixture was filtered through Celite and evaporated to afford L-4-  
10 (N,N-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester of sufficient purity  
for immediate use in reactions with chloropyrimidines.

## Method XX

## Preparation of

15 *N*-Benzyloxycarbonyl-L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine  
*tert*-Butyl Ester

To a stirred solution maintained at 0°C of *N*-benzyloxycarbonyl-L-  
tyrosine *tert*-butyl ester (1.0 eq) and triethylamine (2.5 eq) in  
dichloromethane was added 4-nitrophenyl chloroformate (1.0 eq). The  
20 mixture was stirred for 30 min at 0°C, and then 1-methylpiperazine (1.5 eq)  
was added, and then the mixture was stirred for 2 h while warming to room  
temperature. The mixture was diluted with ethyl acetate and washed five  
times with 10% potassium carbonate and once with saturated sodium  
chloride. The organic extracts were treated with anhydrous magnesium  
25 sulfate, filtered, and evaporated to afford *N*-benzyloxycarbonyl-L-4-(4-  
methylpiperazin-1-ylcarbonyloxy)phenylalanine *tert*-butyl ester of sufficient  
purity for immediate conversion to L-4-(4-methylpiperazin-1-  
ylcarbonyloxy)phenylalanine *tert*-butyl ester.

30

- 173 -

## Method YY

## Preparation of

L-4-(4-Methylpiperazin-1-ylcarbonyloxy)phenylalanine *tert*-Butyl Ester

A suspension of *N*-benzyloxycarbonyl-L-4-(4-methylpiperazin-1-ylcarbonyloxy)-phenylalanine *tert*-butyl ester and an equal mass of 10% palladium on carbon in methanol was shaken under 55 psi hydrogen gas for 1 h, at which point TLC indicated consumption of *N*-benzyloxycarbonyl-L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine *tert*-butyl ester. The mixture was filtered through Celite and evaporated to afford L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine *tert*-butyl ester of sufficient purity for immediate use in reactions with chloropyrimidines.

## Method ZZ

*tert*-Butyl Ester Cleavage Procedure IV

The *tert*-butyl ester was dissolved in 96% formic acid and heated to 40°C for 16 h, at which point TLC indicated consumption of the *tert*-butyl ester. The mixture was evaporated under a stream of air to give a residue, which was stored under high vacuum for 72 h to afford the pure carboxylic acid.

## Method AAA

Preparation of 2,4-Dichloro-5-nitropyrimidine

5-Nitrouracil was treated with phosphorus oxychloride and *N,N*-dimethylaniline, according to the procedure of Whittaker (J. Chem. Soc. 1951, 1565), to give 2,4-dichloro-5-nitropyrimidine as an orange oil, which was used without distillation immediately in the next step.

-- 174 --

## Method BBB

## Preparation of

*N*-(2-Chloro-5-nitropyrimidin-4-yl)-L-4-(*N,N*-  
dimethylcarbamoyloxy)phenylalanine  
*tert*-Butyl Ester

5 To a stirred solution of L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine  
*tert*-butyl ester (6.38 g, 20.69 mmol) and *N,N*-diisopropylethylamine (5.40  
mL, 4.01 g, 31.03 mmol) in 70 mL CH<sub>2</sub>Cl<sub>2</sub> at 0°C, was added a solution of  
2,4-dichloro-5-nitropyrimidine (3.25 g, 20.69 mmol) in 70 mL of CH<sub>2</sub>Cl<sub>2</sub>, at  
10 such a rate the temperature did not exceed 10°C. After the addition, the  
mixture was stirred at 0-10°C for 15 minutes, at which point TLC indicated  
conversion of 2,4-dichloro-5-nitropyrimidine. To the mixture were added  
100 mL 1 M KHSO<sub>4</sub> and 200 mL diethyl ether. The organic layer was  
15 separated, washed (H<sub>2</sub>O, sat. NaHCO<sub>3</sub>, and sat. NaCl), dried (MgSO<sub>4</sub>),  
filtered, and evaporated to give *N*-(2-chloro-5-nitropyrimidin-4-yl)-L-4-(*N,N*-  
dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester (9.52 g, 20.45 mmol,  
99%) as an orange oil, which was used immediately in the next step.

## Method CCC

## Preparation of

*N*-(5-Aminopyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine  
*tert*-Butyl Ester

20 A mixture of *N*-(2-chloro-5-nitropyrimidin-4-yl)-L-4-(*N,N*-  
dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester (9.52 g, 20.45 mmol),  
25 Degussa-type 20% palladium on carbon (9.52 g), NaHCO<sub>3</sub> (8.59 g, 102.2  
mmol), and 165 mL MeOH was shaken under 55 psi H<sub>2</sub> for 16 h, at which  
point TLC indicated conversion of *N*-(2-chloro-5-nitropyrimidin-4-yl)-L-4-  
(*N,N*-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester into a single  
product. The mixture was filtered through Celite, and the filtrate was  
30 evaporated to give a residue, which was dissolved by addition of 150 mL

-- 175 --

EtOAc and 75 mL H<sub>2</sub>O. The organic layer was separated, washed (sat. NaCl), dried (MgSO<sub>4</sub>), filtered, and evaporated to give *N*-(5-aminopyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester (7.14 g, 17.79 mmol, 87%) as an orange solid, which was used immediately in the next step.

## Method DDD

## Preparation of

*N*-(5-(*N*-4-Toluenesulfonylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine *tert*-Butyl Ester

To a stirred solution of *N*-(5-aminopyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester (1.00 g, 2.49 mmol) in 10 mL anhydrous pyridine at 0°C, was added in portions 4-toluenesulfonylchloride (0.474 g, 2.49 mmol). After the addition, the resulting red solution was stirred at 0°C for 3 h, at which point TLC indicated nearly complete conversion of *N*-(5-aminopyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester. To the mixture was added 3-dimethylaminopropylamine (0.325 mL, 0.264 g, 2.49 mmol), and the mixture was stirred for 30 min while warming to room temperature. The mixture was poured into 100 mL 1 M KHSO<sub>4</sub>, and extracted with 150 mL EtOAc. The organic layer was washed (2 x 1 M KHSO<sub>4</sub>, H<sub>2</sub>O, sat. NaHCO<sub>3</sub>, sat. NaCl), dried (MgSO<sub>4</sub>), filtered, and evaporated to give a brown residue, which was purified by flash chromatography using EtOAc/hexanes on silica gel, to give *N*-(5-(*N*-4-toluenesulfonylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester (1.01 g, 1.81 mmol, 73%) as a clear oil.

-- 176 --

## Method EEE

## Preparation of

*N*-(5-(*N*-Methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-Butyl Ester

5 To a stirred two-phase mixture of 45 mL 1 M NaOH and 25 mL diethyl ether at 0°C, was added in portions 1-methyl-3-nitro-1-nitrosoguanidine (1.33 g, 9.05 mmol). After stirring for 25 min, at which point evolution of N<sub>2</sub> had subsided, the bright yellow solution of diazomethane in diethyl ether  
10 was transferred by pipette to a stirred solution of *N*-(5-(*N*-4-toluenesulfonylamino)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester (1.01 g, 1.81 mmol) in 15 mL diethyl ether and 15 mL CH<sub>2</sub>Cl<sub>2</sub> at 0°C. After stirring for 15 min, at which point TLC indicated complete conversion of *N*-(5-(*N*-4-toluenesulfonylamino)pyrimidin-4-yl)-*L*-4-(*N,N*-  
15 dimethylcarbamyloxy)phenylalanine *tert*-butyl ester, excess AcOH was added to destroy unreacted diazomethane. The mixture was diluted with 100 mL diethyl ether, washed (2 x sat. NaHCO<sub>3</sub>, sat. NaCl), dried (MgSO<sub>4</sub>), filtered and evaporated to give a yellow residue, which was purified by flash  
20 chromatography using EtOAc/hexanes on silica gel, to give *N*-(5-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester (0.846 g, 1.48 mmol, 82%) as a clear oil.

25

## Method FFF

Preparation of Diethyl 2-(*N,N*-Dialkylamino)malonate

The appropriate amine (1.0 eq) was added to a 0°C solution of diethyl bromomalonate (1.0 eq) and *N,N*-diisopropylethyl amine (1.1 eq) in ethanol. The mixture was stirred and allowed to warm room temperature. After 16  
30 hours, the reaction mixture was concentrated and the residue was suspended

-- 177 --

in ethyl acetate and sat.  $\text{NaHCO}_3$ . The organic portion was washed with sat  $\text{NaHCO}_3$ , brine, dried ( $\text{MgSO}_4$ ) filtered and concentrated to yield the diethyl 2-(*N,N*-dialkylamino)malonate, of sufficient purity for immediate conversion to a 5-(*N,N*-dialkylamino)-4,6-dihydroxypyrimidine.

5

#### Method GGG

##### Preparation of 5-(*N,N*-Dialkylamino)-4,6-dihydroxypyrimidine

A suspension of a diethyl 2-(*N,N*-dialkylamino)malonate (1.0 eq), formamidine acetate (1.10 eq.) and 25% sodium methoxide in methanol (3.3 eq) was heated to 65°C for 3.5 hours. The reaction mixture was cooled and diluted with water. The mixture was acidified to pH = 4.5 by addition of dilute HCl. The resulting precipitate was collected by filtration, washed with water, and dried under vacuum to afford a 5-(*N,N*-dialkylamino)-4,6-dihydroxypyrimidine of sufficient purity for immediate conversion to a 5-(*N,N*-dialkylamino)-4,6-dichloropyrimidine. Alternatively, the acidified solution was evaporated to give a solid residue, which was extracted with boiling ethanol. The ethanol extracts were filtered and concentrated to give a residue, which was recrystallized from isopropyl alcohol to afford a 5-(*N,N*-dialkylamino)-4,6-dihydroxypyrimidine of sufficient purity for immediate conversion to a 5-(*N,N*-dialkylamino)-4,6-dichloropyrimidine.

15

20

#### Method HHH

##### Preparation of 5-(*N,N*-Dialkylamino)-4,6-dichloropyrimidine

A 5-(*N,N*-dialkylamino)-4,6-dihydroxypyrimidine (1.0 eq) was suspended in  $\text{POCl}_3$  (15.0 eq), and the mixture was heated to reflux for 16 hours. Then the mixture was cooled and carefully poured into a suspension of ethyl ether and aqueous  $\text{K}_2\text{CO}_3$ . The organic portion was washed with brine, dried ( $\text{MgSO}_4$ ), filtered and concentrated to yield a 5-(*N,N*-dialkylamino)-4,6-dichloro-pyrimidine of sufficient purity for immediate reaction with alkylamines.

25

30

-- 178 --

## Method III

## Preparation of

4-(N-Alkylamino)-5-(N,N-dialkylamino)-6-chloropyrimidine

A 5-(N,N-dialkylamino)-4,6-dichloropyrimidine (1.0 eq),  
5 L-4-(N,N-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester (1.5 eq) and  
N,N-diisopropyl ethylamine (1.5 eq) were dissolved in ethanol and heated to  
120°C in a sealed tube for 72 h. The cooled reaction mixture was  
concentrated, and the residue dissolved in ethyl acetate. The ethyl acetate  
solution was washed with 0.2 N citric acid, water, saturated NaHCO<sub>3</sub>, brine,  
10 dried (MgSO<sub>4</sub>), filtered and concentrated. The residue was purified by silica  
gel chromatography using ethyl acetate/hexanes to afford the 4-(N-  
alkylamino)-5-(N,N-dialkylamino)-6-chloropyrimidine.

## Method JJJ

15 Preparation of 4-(N-Alkylamino)-5-(N,N-dialkylamino)pyrimidine

A 4-(N-Alkylamino)-5-(N,N-dialkylamino)-6-chloropyrimidine (1.0 eq),  
an equal mass of 10% palladium on carbon. and NaHCO<sub>3</sub> (5.0 eq) were  
suspended in methanol. The reaction mixture was hydrogenated at 45 psi  
hydrogen for 16 hours and then filtered through a pad of Celite. The filtrate  
20 was concentrated, and the residue was dissolved in ethyl acetate. The ethyl  
acetate solution was washed with water, brine, dried (MgSO<sub>4</sub>), filtered and  
concentrated to yield an oil. The oil was purified by column  
chromatography on silica gel using ethyl acetate and hexanes to afford a pure  
4-(N-alkylamino)-5-(N,N-dialkylamino)pyrimidine.

25

## Method KKK

Suzuki Coupling Procedure V

To an ethyleneglycol dimethyl ether solution of  
tetrakis(triphenylphosphine) palladium (0.04 eq) was added N-(5-bromo-2-  
30 chloro-pyrimidin-4-yl)-L-4-(N,N-dimethylcarbamoyloxy)phenylalanine *tert*-



-- 179 --

butyl ester (1.5 eq.). After stirring for approximately ten minutes *o*-tolylboronic acid (1.5 eq) and 2M Na<sub>2</sub>CO<sub>3</sub> (2.0 eq) were added, and the reaction flask was evacuated and flushed with nitrogen gas. The reaction was heated to reflux for four hours. The reaction mixture was then cooled and  
5 diluted with water and methylene chloride. The organic phase was separated and washed with brine, dried (MgSO<sub>4</sub>), filtered and concentrated. The residue was purified by silica gel chromatography using ethyl acetate/hexanes to afford the desired product.

10

## Method LLL

## Preparation of

L-Phenylalanine Isopropyl Ester Hydrochloride orL-Tyrosine Isopropyl Ester Hydrochloride

Excess HCl gas was added with stirring to a suspension of L-  
15 phenylalanine or L-tyrosine in excess isopropanol. The mixture was heated to reflux for 16 h, and then the volatiles were evaporated under vacuum to give L-phenylalanine isopropyl ester hydrochloride or L-tyrosine isopropyl ester hydrochloride of sufficient purity for immediate use.

20

## Method MMM

Bromopyrimidine Debromination Procedure

The bromopyrimidine was dissolved in isopropyl alcohol to which was added 10% palladium on carbon. The reaction was hydrogenated at 45 psi hydrogen. Filtration and concentration of the filtrate yielded the desired  
25 dehalogenated pyrimidine.

## Method NNN

Preparation of 2-Isopropoxypyrimidine

A 2-chloropyrimidine was dissolved in isopropyl alcohol to which was  
30 added diisopropylamine. The reaction was heated in a sealed tube for ten

-- 180 --

days at 130°C. The cooled reaction mixture was concentrated, and the product purified via silica gel column chromatography to yield the 2-isopropoxypyrimidine.

5

## Method OOO

Heck Reaction Procedure III

To a dioxane/triethylamine (1:1 by volume) solution of the *N*-(5-iodopyridin-4-yl)-*L*-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine isopropyl ester (1.0 eq), triphenylphosphine (0.05 eq), copper (I) iodide (0.2 eq) was added phenylacetylene (4.0 eq). After flushing the solution for ten minutes with nitrogen gas, dichlorobis(triphenylphosphine)palladium (0.10 eq) was added, and the resulting reaction mixture heated to 50°C for 16 hours. The reaction mixture was then diluted with ethyl acetate and water, and the organic portion was washed with 0.2 N citric acid, water, saturated NaHCO<sub>3</sub>, brine, dried (MgSO<sub>4</sub>), filtered and concentrated. The residue was chromatographed on a silica gel column using ethyl acetate/hexanes to afford the desired product.

10

15

## Method PPP

20

## Preparation of

*N*-[5-(Phenyl)pyrimidin-4-yl]-*L*-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine  
*tert*-Butyl Ester

*N*-[5-iodopyrimidin-4-yl]-*L*-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester (123 mg, 0.2 mmol) was diluted in dry DMF (5 mL) under nitrogen with KOAc (3.0 eq, 73 mg), bis(pinacolato)diboron (1.1 eq, 63 mg), and a catalytic amount of [1,1'-bis(diphenylphosphino)-ferrocene]dichloropalladium (II) complex with dichloromethane (1:1). The reaction was heated for 2 hours at 100°C. To this was added, K<sub>3</sub>PO<sub>4</sub> (2.0 eq, 105 mg), iodobenzene (2.0 eq, 0.056 mL) and an additional catalytic amount of [1,1'-bis(diphenylphosphino)-ferrocene]dichloropalladium (II)

25

30

-- 181 --

complex with dichloromethane (1:1). The reaction mixture was stirred overnight at 100°C. EtOAc was added and the organic layer washed with brine, dried over MgSO<sub>4</sub>. Upon filtration, and evaporation of the solvent under reduced pressure, the crude material was eluted on column chromatography (silica gel) with EtOAc/hexanes 1:1. The desired material was isolated in good yields.

#### Method QQQ

##### Preparation of 2-Amino-3-Chloropyrazine

A mixture of 2,3-dichloropyrazine (Lancaster) and ammonium hydroxide was heated in a sealed tube at 100°C for 24 h resulting in a white precipitate. The precipitate was collected by filtration and dried under vacuum to afford 2-amino-3-chloropyrazine of sufficient purity for immediate conversion to 2-chloro-3-nitropyrazine.

#### Method RRR

##### Preparation of 2-Chloro-3-Nitropyrazine

The method (p. 1638) of Hartman et al. (J. Med. Chem. **1984**, 27(12), 1634-1639) was employed to convert 2-amino-3-chloropyrazine into 2-chloro-3-nitropyrazine of sufficient purity for immediate use.

#### Method SSS

##### Preparation of 4-Alkylamino-2-dialkylamino-5-nitropyrimidine

A solution of 1.0 eq 4-alkylamino-2-chloro-5-nitropyrimidine and 5.0 eq dialkylamine in THF was allowed to stand for 16 h. The mixture was diluted with ethyl acetate and then washed with pH = 4.5 citrate buffer and saturated sodium chloride. The organic extracts were treated with anhydrous magnesium sulfate, filtered, and evaporated to give a residue, which was purified by chromatography on silica gel using ethyl acetate and hexanes.

-- 182 --

## Method TTT

Preparation of L-4-(2,6-Dimethoxyphenyl)phenylalanine Methyl Ester

To a stirred solution (DMF, 66 mL) of N-Boc-L-(p-iodo)phenylalanine methyl ester (13.2 g, 32.7 mmol) prepared according to the procedure of Schwabacher et al., J. Org. Chem. 1994, 59, 4206-4210) was added  
5 Pd(PPh<sub>3</sub>)<sub>4</sub> (0.03 eq, 1.13 g, 1 mmol). The solution was stirred for 10 min and then 2,6-dimethoxyboronic acid (1.2 eq, 7.1 g, 39 mmol) and K<sub>3</sub>PO<sub>4</sub> (1.5 eq, 10.4 g, 49 mmol) were added. The reaction flask was evacuated and flushed with nitrogen. This process was repeated twice and the reaction  
10 mixture was then heated to 100°C under a stream of nitrogen for about 3.5 h at which time TLC showed the reaction to be complete (4.5:1 hexanes:EtOAc, R<sub>f</sub> = 0.2, UV active). The reaction mixture was cooled and partitioned between water and ethyl acetate (200 mL each). The organic portion was washed with 0.2N citric acid (3 X 100 mL), brine (1 X 100  
15 mL), dried (MgSO<sub>4</sub>), filtered and stripped to a thick reddish oil, about 13 g. The resulting product was chromatographed on silica gel eluting with 4.5:1 hexanes/EtOAc, R<sub>f</sub> = 0.2. The combined fractions were stripped and treated with methanol saturated with HCl to yield the title intermediate as the hydrochloride salt.

20

## Example 1

## Synthesis of

*N*-(2-Chloro-5-nitropyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

25

Step A - Preparation of 2,4-Dichloro-5-nitropyrimidine

5-Nitrouracil (Aldrich Chemical Company) was treated with phosphorous oxychloride and *N,N*-dimethylaniline according to the procedure described in Whittaker, J. Chem. Soc. 1951, 1565, to give 2,4-dichloro-5-  
30 nitropyrimidine as an orange oil which was used without distillation immediately in the next step.

-- 183 --

Step B - Preparation of *N*-(2-Chloro-5-nitropyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-Butyl Ester

To a stirred solution of L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester (6.38 g, 2069 mol) and *N,N*-diisopropylethylamine (5.40 mL, 4.01 g, 31.03 mol.) in 70 mL CH<sub>2</sub>Cl<sub>2</sub> at 0°C, was added a solution of 2,4-dichloro-5-nitropyrimidine (3.25 g, 20.69 mol.) in 70 mL CH<sub>2</sub>Cl<sub>2</sub> at such a rate that the temperature did not exceed 10°C. After the addition, the mixture was stirred at 0-10°C for 15 minutes, at which point TLC indicated conversion of the starting materials. To the mixture were added 100 mL 1 M KHSO<sub>4</sub> and 200 mL diethyl ether. The organic layer was separated, washed (H<sub>2</sub>O, sat. NaHCO<sub>3</sub>, and sat. NaCl), dried (MgSO<sub>4</sub>), filtered, and evaporated to give the title compound (9.52 g, 2045 mol., 99%) as an orange oil.

Step C - Preparation of *N*-(2-Chloro-5-nitropyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

The title compound was prepared by hydrolysis of the product from Step B using the procedure of Example 5.

Example 2

Synthesis of

*N*-[5-(*N*-4-Toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-Butyl Ester

Step A - Preparation of *N*-(5-Aminopyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-Butyl Ester

A mixture of *N*-(2-chloro-5-nitropyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester (9.52 g, 20.45 mol), Degussa-type 20% palladium on carbon (9.52 g), NaHCO<sub>3</sub> (8.59 g, 102.2 mol), and 165 mL MeOH was shaken under 55 psi for 16 h, at which point TLC indicated conversion of the starting material into a single product. The

mixture was filtered through Celite, and the filtrate was evaporated to give a residue, which was dissolved by addition of 150 mL EtOAc and 75 mL H<sub>2</sub>O. The organic layer was separated, washed (sat. NaCl), dried (MgSO<sub>4</sub>), filtered, and evaporated to give the title intermediate (7.14 g, 17.79 mol, 87%) as an orange solid, which was used immediately in the next step.

Step B - Preparation of *N*-[5-(*N*-4-Toluenesulfonyl-amino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-Butyl Ester

To a stirred solution of the product from Step A (100 g, 2.49 mol) in 10 mL anhydrous pyridine at 0 °C, was added in portions 4-toluenesulfonyl chloride (0.474 g, 2.49 mol). After the addition, the resulting red solution was stirred at 0 °C for 3 h, at which point TLC indicated nearly complete conversion of the starting material. To the mixture was added 3-dimethylaminopropylamine (0.325 mL, 0.264 g, 2.49 mol), and the mixture was stirred for 30 min while warming to room temperature. The mixture was poured into 100 mL 1 M KHSO<sub>4</sub>, and extracted with 150 mL EtOAc. The organic layer was washed (2 x 1 M KHSO<sub>4</sub>, H<sub>2</sub>O, sat. NaHCO<sub>3</sub>, sat. NaCl), dried (MgSO<sub>4</sub>), filtered, and evaporated to give a brown residue, which was purified by flash chromatography using EtOAc/hexanes on silica gel, to give the title compound (1.01 g, 1.81 mol., 73%) as a clear oil.

Example 3

Synthesis of  
*N*-[5-(*N*-4-Toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

The title compound was prepared by hydrolysis of *N*-[5-(*N*-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester using the procedure of Example 5.

-- 185 --

## Example 4

**Synthesis of**  
***N*-[5-(*N*-Methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl]-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-Butyl Ester**

5

To a stirred two-phase mixture of 45 mL 1 M NaOH and 25 mL diethyl ether at 0°C, was added in portions 1-methyl-3-nitro-1-nitrosoguanidine (1.33 g, 9.05 mol). After stirring for 25 min, at which point evolution of N<sub>2</sub> had subsided, the bright yellow solution of diazomethane in diethyl ether was transferred by pipette to a stirred solution of the product of Example 2 (1.01 g, 1.81 mol) in 15 mL diethyl ether and 15 mL CH<sub>2</sub>Cl<sub>2</sub> at 0°C. After stirring for 15 min, at which point TLC indicated complete conversion of the starting material, excess AcOH was added to destroy unreacted diazomethane. The mixture was diluted with 100 mL diethyl ether, washed (2 x sat. NaHCO<sub>3</sub>, sat. NaCl), dried (MgSO<sub>4</sub>), filtered and evaporated to give a yellow residue, which was purified by flash chromatography using EtOAc/hexanes on silica gel, to give the title compound (0.846 g, 1.48 mol, 82%) as a clear oil.

10

15

## Example 5

20

**Synthesis of**  
***N*-[5-(*N*-Methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl]-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine**

The product of Example 4 (0.400 g, 0.700 mol) was dissolved in 8 mL 96% formic acid, and the mixture was heated to 40°C for 16 h, at which point TLC indicated conversion of the starting material. Most of the formic acid was evaporated under a stream of N<sub>2</sub>, and then the residue was placed under high vacuum for 48 h to give the title compound (0.382 g, 0.700 mol, 100%) as a clear oil.

25

30 Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 8.33 (bs, 1H), 8.07 (bs, 1H), 7.64 (d, J = 8.1 Hz, 2H), 7.42 (d, J = 8.1 Hz, 2H), 7.36 (bs, 1H), 7.29 (bs, 2H), 6.99 (d,

30

-- 186 --

J = 7.5 Hz, 2H), 5.07-4.96 (m, 1H), 3.42-3.31 (m, 1H), 3.25-3.15 (m, 1H), 3.08 (s, 3H), 3.05 (bs, 3H), 2.96 (s, 3H), 2.44 (s, 3H).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ = 174.7, 174.6, 164.6, 157.8, 156.8, 152.9., 152.1, 146.5, 135.4, 135.1, 131.7, 131.3, 129.4, 123.2, 122.9, 55.8, 38.2, 37.1, 36.8, 36.7, 21.5.

Using the appropriate starting materials and reagents, the following additional compounds were prepared:

- 10 *N*-[5-(*N,N*-Di-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine (Example 6);
- N*-[5-[*N*-(1-*N'*-Methylpyrazol-4-ylsulfonyl)-*N*-methylamino]pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine (Example 7);
- 15 *N*-[5-(*N*-Methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine isopropyl ester (Example 8);
- 20 *N*-[5-(*N*-Methyl-*N*-3-pyridylsulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester (Example 9); and

#### Example 10

##### Synthesis of

25 *N*-(5-(*N*-Methyl-*N*-(1-butylpyrazol-4-yl)sulfonylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. 5-Nitouracil (Aldrich) was converted via Method AAA into 2,4-dichloro-5-nitopyrimidine. L-4-(*N,N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 2,4-dichloro-5-nitopyrimidine were coupled via Method BBB, and the product of this coupling was sequentially converted via Methods CCC, DDD (using 1-butyl-4-chlorosulfonylpyrazole), EEE and ZZ to give the title compound.



-- 187 --

Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  = 8.35 (s, 1H), 8.14 (s, 1H), 7.76 (s, 1H),  
7.61 (bs, 1H), 7.23 (bs, 2H), 6.98 (d, 2H), 5.01-4.94 (m, 1H), 4.19 (t, 2H),  
3.40 - 3.28 (m, 1H), 3.26-3.14 (m, 1H), 3.09 (s, 3H), 3.06 (bs, 3H), 2.96  
5 (s, 3H), 1.84 (pent., 2H), 1.29 (sext., 2H), 0.945 (t, 3H).

#### Example 11

10

##### Synthesis of

##### ***N*-(5-(2,4-Dimethoxypyrimidin-5-yl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine**

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via  
15 Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-  
phenylalanine *tert*-butyl ester. 4(3H)-pyrimidinone (Aldrich) was  
sequentially converted via Methods N and O into 4-chloro-5-iodopyrimidine.  
L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4-chloro-5-  
iodopyrimidine were coupled via Method P, and the coupled product was  
20 reacted with 2,4-dimethoxypyrimidin-5-yl boronic acid (Frontier Scientific,  
Inc.) via Method S. The product of this coupling was converted via Method  
KK to give the title compound.

#### Example 12

25

##### Synthesis of

##### ***N*-(5-(2,6-Difluorophenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine**

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via  
30 Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-  
phenylalanine *tert*-butyl ester. 4(3H)-Pyrimidinone (Aldrich) was  
sequentially converted via Methods N and O into 4-chloro-5-iodopyrimidine.  
L-4-(*N,N*-Dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4-chloro-  
5-iodopyrimidine were coupled via Method P, and the coupled product was

reacted with 2,6-difluorophenyl boronic acid (Lancaster Synthesis) via Method R. The product of this coupling was converted via Method HH to give the title compound.

5

### Example 13

#### Synthesis of

10

#### *N*-(5-(2-Hydroxymethylphenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. 4(3H)-Pyrimidinone (Aldrich) was sequentially converted via Methods N and O into 4-chloro-5-iodopyrimidine. L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4-chloro-5-iodopyrimidine were coupled via Method P, and the coupled product was reacted with 2-(hydroxymethyl)phenyl boronic acid (Lancaster Synthesis) via Method Q. The product of this coupling was converted via Method HH to give the title compound.

20

### Example 14

#### Synthesis of

25

#### *N*-(2-(*N*-Cyclohexylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with cyclohexylamine (Aldrich) via Method CC to give a product that was coupled

30

-- 189 --

with o-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

Physical data were as follows:

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 9.68 (s, 1H), 7.3 – 6.8 (m, 9H), 6.35 (m, 1H),  
5 4.73(m, 1H), 3.81 (bs, 1H), 3.6 – 3.0 (m, 2H), 3.09 (s, 3H), 3.0 (s, 3H),  
2.18 (s, 1.5H), 1.94 (s, 1.5H), 2.1 – 1.1 (m, 10H).

<sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 176.11, 175.94, 160.05, 159.79, 154.76,  
153.58, 150.05, 150.01, 139.26, 137.84, 137.63, 134.29, 134.15, 130.66,  
130.36, 130.11, 129.14, 126.70, 126.41, 121.25, 109.57, 109.39, 56.84,  
10 56.35, 50.15, 36.55, 36.32, 32.34, 31.99, 25.41, 24.86, 19.48, 19.27.

### Example 15

#### 15      **Synthesis of** ***N*-(2-(*N*-Methyl-*N*-(1-methylpiperidin-4-yl)amino)-5-(2-tolyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine**

*L*-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via  
Methods UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamyloxy)-  
20 phenylalanine *tert*-butyl ester. *L*-4-(*N,N*-Dimethylcarbamyloxy)-  
phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich)  
were coupled via Method BB. The product of this reaction was reacted with  
1-methyl-4-(*N*-methylamino)piperidine (Aldrich) via Method CC to give a  
product that was coupled with o-tolyl boronic acid (Aldrich) via Method DD.  
25 The product of this coupling was converted via Method ZZ to give the title  
compound.

Physical data were as follows:

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 8.82 (s, 2H), 8.43 (s, 1H), 7.62 (s, 1H),  
7.30 – 6.90 (m, 8H), 5.42 (br, 1H), 4.66 (br, 2H), 3.60 – 2.8 (m, 15H),  
30 2.66 (bs, 3H), 2.32 (br, 2H), 2.18 (s, 1.5H), 1.82 (brs, 3.5H).

***N*-(2-(*N*-Ethyl-*N*-isopropylamino)-5-(2-tolyl)pyrimidin-4-yl)-*L*-4-(*N*,*N*-dimethylcarbamoyloxy)phenylalanine**

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with *N*-ethyl-*N*-isopropylamine (Aldrich) via Method CC to give a product that was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

**Physical data were as follows:**

<sup>1</sup> H NMR (CDCl<sub>3</sub>): δ = 8.0 – 6.5 (br, 1H), 7.66 (s, 0.5H), 7.62 (s, 0.5H), 7.3 – 6.8 (m, 8H), 6.2 (m, 1H), 4.86 (br, 1H), 4.70 (m, 1H), 3.70 – 3.08 (m, 4H), 3.09 (s, 3H), 3.0 (s, 3H), 2.14 (bs, 1.5H), 1.92 (bs, 1.5H), 1.4 – 0.9 (br, 9H).

<sup>13</sup> C NMR (CDCl<sub>3</sub>): δ = 174.38, 174.19, 159.44, 159.16, 155.24, 154.68, 152.39, 150.02, 141.63, 137.77, 137.56, 134.30, 134.09, 130.79, 130.66, 130.54, 130.46, 130.41, 130.33, 130.08, 129.07, 126.54, 126.45, 126.38, 121.21, 121.16, 110.27, 110.01, 56.77, 56.36, 47.59, 36.80, 36.55, 36.32, 20.27, 20.18, 19.57, 19.38, 14.51.

### Example 17

***N*-(5-(2,4,6-Trimethylphenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine**

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-

-- 191 --

phenylalanine *tert*-butyl ester. 4(3H)-Pyrimidinone (Aldrich) was sequentially converted via Methods N and O into 4-chloro-5-iodopyrimidine. L-4-(*N,N*-Dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester and 4-chloro-5-iodopyrimidine were coupled via Method P, and the coupled product was  
5 reacted with 2,4,6-trimethylphenyl boronic acid (Frontier Scientific, Inc) via Method R. The product of this coupling was converted via Method HH to give the title compound.

Physical data were as follows:

<sup>1</sup> H NMR (CD<sub>3</sub>OD):  $\delta$  = 8.68 (d, 1H), 7.95 (d, 1H), 7.10 (d, 2H),  
10 7.09 – 6.95(m, 2H), 6.94 – 6.91 (m, 2H), 5.32 – 5.27 (m, 1H), 3.42 – 3.36 (m, 1H), 3.15 – 3.09 (m, 4H), 2.97 (s, 3H), 2.33 (s, 3H), 2.04 (s, 3H), 1.84 (s, 3H).

<sup>13</sup> C NMR (CD<sub>3</sub>OD):  $\delta$  = 172.9, 163.5, 161.5, 161.0, 156.7, 152.0,  
151.9, 142.6, 141.5, 138.9, 138.6, 135.3, 131.2, 130.4, 130.3, 126.5,  
15 123.0, 120.3, 56.4, 36.7, 36.6, 36.5, 21.2, 19.9, 19.7.

#### Example 18

##### Synthesis of 20 *N*-(5-Isopropylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamoyloxy)-  
25 phenylalanine *tert*-butyl ester. Diethyl 2-isopropylmalonate (Aldrich) was sequentially converted via Methods OO and QQ into 4,6-dichloro-5-isopropylpyrimidine. L-4-(*N,N*-Dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester and 4,6-dichloro-5-isopropylpyrimidine were coupled via Method RR, and the product of this coupling was sequentially converted via Methods  
30 SS and ZZ to give the title compound.

Physical data were as follows:

-- 192 --

<sup>1</sup> H NMR (CD<sub>3</sub>OD): δ = 8.44 (bs, 1H), 7.94 (bs, 1H), 7.22 (d, 2H), 6.94(d, 2H), 5.12 (dd, 1H), 3.46 (dd, 1H), 3.19 (dd, 1H), 3.07 (s, 3H), 2.95 (s, 3H), 3.00 – 2.88 (m, 1H), 1.25 (d, 3H), 1.13 (d, 3H).

<sup>13</sup> C NMR (CD<sub>3</sub>OD): δ = 175.60, 165.74, 163.78, 156.91, 152.38, 151.85, 141.88, 136.30, 131.43, 126.17, 122.87, 57.84, 37.48, 36.81, 36.64, 26.63, 21.09, 20.94.

### Example 19

#### Synthesis of *N*-(2-(*N*-Methyl-*N*-butylamino)-5-(2-tolyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

*L*-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. *L*-4-(*N,N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with *N*-methyl-*N*-butylamine (Aldrich) via Method CC to give a product that was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

Physical data were as follows:

<sup>1</sup> H NMR (CDCl<sub>3</sub>): δ = 12.5 – 11.4 (br, 1H), 7.6 (s, 0.5H), 7.58 (s, 0.5H), 7.3 – 6.8 (m, 8H), 6.3 (m, 1H), 4.7 (m, 1H), 3.7 – 2.9 (m, 4H), 3.08 (s, 3H), 3.01 (s, 6H), 2.13 (s, 1.5H), 1.91 (s, 1.5H), 1.57 (bs, 2H), 1.33 (m, 2H), 0.96 (t, 3H).

<sup>13</sup> C NMR (CDCl<sub>3</sub>): δ = 174.21, 174.06, 159.37, 159.22, 154.69, 153.52, 169.99, 141.87, 137.77, 137.54, 134.43, 130.78, 130.59, 130.10, 128.98, 126.51, 126.32, 121.17, 121.11, 110.20, 109.96, 56.82, 56.43, 50.03, 36.54, 36.32, 35.91, 29.27, 19.89, 19.52, 19.35, 13.84.

-- 193 --

## Example 20

Synthesis of  
*N*-(2-(*N*-Ethyl-*N*-propylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N*,*N*-  
dimethylcarbamoyloxy)phenylalanine

5

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N*,*N*-dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N*,*N*-Dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with *N*-ethyl-*N*-propylamine (Aldrich) via Method CC to give a product that was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

10

Physical data were as follows:

15

<sup>1</sup> H NMR (CDCl<sub>3</sub>): δ = 11.0 – 9.5 (br, 1H), 7.66 (s, 0.5H), 7.64 (s, 0.5H), 7.4 – 6.8 (m, 8H), 6.28 (m, 1H), 4.65 (m, 1H), 3.70 – 2.80 (m, 6H), 3.09 (s, 3H), 3.01 (s, 3H), 3.01 (s, 3H), 2.2 (s, 1.5H), 1.85 (s, 1.5H), 1.58 (bs, 2H), 1.05 (bs, 3H), 0.85 (bs, 3H).

20

<sup>13</sup> C NMR (CDCl<sub>3</sub>): δ = 174.26, 174.11, 159.36, 159.11, 154.70, 153.07, 149.96, 142.43, 137.80, 137.56, 134.54, 134.37, 130.84, 130.74, 130.57, 130.14, 128.86, 126.47, 126.29, 121.10, 121.06, 110.01, 109.71, 56.86, 56.49, 49.62, 63.20, 36.55, 36.32, 20.87, 19.61, 19.41, 12.63, 11.03.

25

## Example 21

Synthesis of  
*N*-(2-(*N*,*N*-Diethylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N*,*N*-  
dimethylcarbamoyloxy)phenylalanine

30

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N*,*N*-dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N*,*N*-Dimethylcarbamoyloxy)-

-- 194 --

phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with *N,N*-diethylamine (Aldrich) via Method CC to give a product that was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

Physical data were as follows:

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 12.2 (br, 1H), 7.63 (s, 0.5H), 7.60 (s, 0.5H), 7.40 – 6.80 (m, 8H), 6.28 (m, 1H), 4.70 (m, 1H), 3.80 – 2.90 (m, 6H), 3.06 (s, 3H), 2.98(s, 3H), 2.13 (s, 1.5H), 1.92 (s, 1.5H), 0.90 (s, 6H).

<sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 174.34, 174.15, 159.4, 159.1, 154.70, 152.66, 169.97, 142.06, 137.76, 137.55, 134.44, 134.27, 130.81, 130.57, 130.10, 128.95, 126.48, 126.32, 121.14, 121.08, 110.08, 109.80, 56.78, 56.37, 42.77, 36.53, 36.31, 19.57, 19.38, 12.77.

## Example 22

### Synthesis of *N*-(2-(*N*-Methyl-*N*-ethylamino)-5-(2-tolyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. *L*-4-(*N,N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with *N*-methyl-*N*-ethylamine (Aldrich) via Method CC to give a product that was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

Physical data were as follows:

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 12.5 (br, 2H), 8.23 (s, 1H), 7.50 (s, 0.5H), 7.44(s, 0.5H), 7.30 – 6.80 (m, 8H), 6.10 (m, 1H), 4.75 (m, 1H), 3.58 (bs,



-- 195 --

2H), 3.30 (m, 1H), 3.00 (m, 1H), 3.08 (s, 3H), 3.00 (s, 3H), 2.93(s, 3H),  
2.08 (s, 1.5H), 1.92 (s, 1.5H), 1.50 (s, 3H).

<sup>13</sup> C NMR (CDCl<sub>3</sub>): δ = 174.63, 174.34, 165.72, 159.96, 159.72,  
154.88, 152.62, 150.49, 150.45, 140.64, 137.90, 137.81, 133.83, 133.65,  
5 131.03, 130.95, 130.85, 130.63, 130.10, 130.04, 129.76, 129.62, 126.88,  
126.72, 121.70, 121.61, 110.69, 110.46, 56.65, 56.11, 45.16, 36.57, 36.35,  
35.17, 19.38, 19.17, 11.96.

#### No Example 23

10

#### Example 24

#### 15                                      Synthesis of                                     *N*-(5-Benzyloxypyrimidin-4-yl)-L-phenylalanine

Methyl 2-benzyloxyacetate (Aldrich) was sequentially converted via  
Methods PP and QQ into 4-chloro-5-benzyloxypyrimidine. L-4-  
phenylalanine *tert*-butyl ester (Bachem) and 4-chloro-5-benzyloxypyrimidine  
20 were coupled via Method RR, and the product of this coupling was converted  
via Method ZZ to give the title compound.

Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 8.54 (s, formate), 8.03 (s, 1H), 7.67 (s,  
1H), 7.37 - 7.31(m, 5H), 7.17 - 7.12 (m, 5H), 5.11 (s, 2H), 4.78 - 4.75  
25 (m, 1H), 3.35 - 3.11 (m, 2H).

<sup>13</sup>C NMR (CD<sub>3</sub>OD) : δ = 159.07, 143.16, 132.35, 130.64, 124.52,  
123.94, 123.83, 123.59, 123.11, 122.00, 99.47, 66.28, 50.32, 32.05.

30

-- 196 --

## Example 25

**Synthesis of**  
***N*-(5-Benzoyloxypyrimidin-4-yl)-L-4-(*N,N*-**  
**dimethylcarbamyloxy)phenylalanine**

5

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. Methyl 2-benzyloxyacetate (Aldrich) was sequentially converted via Methods PP and QQ into 4-chloro-5-  
10 benzyloxypyrimidine. L-4-(*N,N*-Dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4-chloro-5-benzyloxypyrimidine were coupled via Method RR, and the product of this coupling was converted via Method ZZ to give the title compound.

15

## Example 26

**Synthesis of**  
***N*-(5-(*N*-Methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl)-L-**  
**phenylalanine**

20

5-Nitrouracil (Aldrich) was converted via Method AAA into 2,4-dichloro-5-nitropyrimidine. L-4-Phenylalanine *tert*-butyl ester (Bachem) and  
25 2,4-dichloro-5-nitropyrimidine were coupled via Method BBB, and the product of this coupling was sequentially converted via Methods CCC, DDD, EEE and ZZ to give the title compound.

30

## Example 27

**Synthesis of**  
***N*-(5-(*N*-Methyl-*N*-3-pyridinesulfonylamino)pyrimidin-4-yl)-L-4-(*N,N*-**  
**dimethylcarbamyloxy)phenylalanine**

35

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-

-- 197 --

phenylalanine *tert*-butyl ester. 5-Nitrouracil (Aldrich) was converted via Method AAA into 2,4-dichloro-5-nitropyrimidine. L-4-(*N,N*-Dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester and 2,4-dichloro-5-nitropyrimidine were coupled via Method BBB, and the product of this coupling was sequentially converted via Methods CCC, DDD (using 3-chlorosulfonylpyridine), EEE and ZZ to give the title compound.

Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  = 8.90 (d, 1H), 8.85 (d, 1H), 8.36 (s, 1H), 8.15 (d, 1H), 7.64 (dd, 1H), 7.53 (bs, 1H), 7.27 (bs, 2H), 6.99 (d, 2H), 5.04 – 4.87 (m, 1H), 3.40 – 3.28 (m, 1H), 3.26 – 3.16 (m, 1H), 3.13 (bs, 3H), 3.09 (s, 3H), 2.97 (s, 3H).

#### Example 28

15

##### Synthesis of *N*-(5-Phenylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester. 4(3H)-Pyrimidinone (Aldrich) was sequentially converted via Methods N and O to 4-chloro-5-iodopyrimidine. L-4-(*N,N*-Dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester and 4-chloro-5-iodopyrimidine were coupled via Method P, and the coupled product was reacted with phenyl boronic acid (Aldrich) via Method S. The product of this coupling was converted via Method HH to give the title compound.

Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  = 8.62 (s, 1H), 8.04 (s, 1H), 7.53 – 7.51 (m, 3H), 7.30 – 7.27 (m, 2H), 7.17 – 7.15 (m, 2H), 7.00 – 6.97 (m, 2H), 5.27 – 5.22 (m, 1H), 3.45 – 3.39 (m, 1H), 3.16 – 3.08 (m, 4H), 2.96 (s, 3H).

<sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  = 173.8, 163.7, 157.5, 152.8, 152.3, 142.4, 135.9, 132.2, 132.1, 131.8, 130.7, 123.9, 122.4, 57.7, 37.7, 37.5.

## Example 29

## Synthesis of

5 ***N*-(3-(*N*-Methyl-*N*-4-toluenesulfonylamino)pyrazin-2-yl)-*L*-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine**

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamoyloxy)-  
10 phenylalanine *tert*-butyl ester. 2,3-Dichloropyrazine (Lancaster) was converted via Method QQQ and RRR into 2-chloro-3-nitropyrazine. *L*-4-(*N,N*-Dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester and 2-chloro-3-nitropyrazine were coupled via Method BBB, and the product of this coupling was sequentially converted via Methods CCC, DDD, EEE and ZZ to give  
15 the title compound.

Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 8.07 (s, formate), 7.94 (d, 1H), 7.59 (d, 2H), 7.51 (d, 1H), 7.36 (d, 2H), 7.29 (d, 2H), 7.01 (d, 2H), 4.90 (m, 1H), 3.30 – 3.18 (m, 2H), 3.08 (s, 3H), 2.96 (s, 3H), 2.94 (s, 3H), 2.43 (s, 3H).

20 <sup>13</sup>C NMR (CD<sub>3</sub>OD): δ = 177.07, 169.41, 158.64, 150.92, 147.23, 145.92, 139.97, 137.14, 133.12, 129.62, 128.90, 125.69, 124.67, 124.08, 116.86, 49.99, 31.67, 31.28, 30.77, 30.62, 15.46.

## No Example 30

25

## Example 31

## Synthesis of

30 ***N*-(5-(2,2,2-Trifluoroethyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine**

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamoyloxy)-  
35 phenylalanine *tert*-butyl ester. 1-Trifluoromethanesulfonyloxy-2,2,2-

-- 199 --

trifluoroethane was sequentially converted via Methods LL, OO and QQ into 4,6-dichloro-5-(2,2,2-trifluoroethyl)pyrimidine. L-4-(*N,N*-Dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester and 4,6-dichloro-5-(2,2,2-trifluoroethyl)pyrimidine were coupled via Method RR, and the  
5 product of this coupling was sequentially converted via Methods SS and ZZ to give the title compound.

Physical data were as follows:

<sup>1</sup> H NMR (CD<sub>3</sub>OD):  $\delta$  = 8.41 (s, 1H), 8.09 (s, formate), 8.06 (s, 1H), 7.24 (d, 2H), 6.96 (d, 2H), 5.06 (m, 1H), 3.60 - 3.40 (m, 2H), 3.37 -  
10 3.11 (m, 2H), 3.08 (s, 3H), 2.96 (s, 3H).

<sup>13</sup> C NMR (CD<sub>3</sub>OD):  $\delta$  = 169.35, 158.91, 156.43, 151.33, 150.97, 148.87, 145.76, 130.21, 125.27, 116.80, 50.80, 31.34, 30.75, 30.60, 26.65, 26.23.

15

### Example 32

#### Synthesis of

*N*-(5-(*N*-Methyl-*N*-3-pyridinesulfonylamino)pyrimidin-4-yl)-L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine Isopropyl Ester  
20

L-Tyrosine (Aldrich) was sequentially converted via Methods LLL, UU, XX and YY into L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine isopropyl ester. 5-Nitrouracil (Aldrich) was converted via Method AAA into 2,4-dichloro-5-nitropyrimidine. L-4-(4-Methylpiperazin-1-  
25 ylcabonyloxy)phenylalanine isopropyl ester and 2,4-dichloro-5-nitropyrimidine were coupled via Method BBB, and the product of this coupling was sequentially converted via Methods CCC, DDD (using 3-chlorosulfonylpyridine) and EEE to give the title compound.

30

-- 200 --

## Example 33

**Synthesis of  
N-(5-Benzylpyrimidin-4-yl)-L-4-(N,N-  
dimethylcarbamyloxy)phenylalanine**

5

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(N,N-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. Diethyl 2-benzylmalonate (Aldrich) was sequentially converted via Methods OO and QQ into 4,6-dichloro-5-benzylpyrimidine. L-4-(N,N-Dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4,6-dichloro-5-benzylpyrimidine were coupled via Method RR, and the product of this coupling was sequentially converted via Methods SS and ZZ to give the title compound.

15

Physical data were as follows:

<sup>1</sup> H NMR (CD<sub>3</sub>OD):  $\delta$  = 8.41 (s, 1H), 8.13 (s, formate), 7.80 (s, 1H) 7.34 – 7.19 (m, 3H), 7.17 (d, 2H), 7.00 (d, 2H), 6.85 (d, 2H), 5.01 (m, 1H), 3.82 (m, 2H), 3.09 (s, 3H), 3.09 – 2.97 (m, 2H), 2.97 (s, 3H).

20

<sup>13</sup> C NMR (CD<sub>3</sub>OD):  $\delta$  = 159.31, 156.23, 150.88, 148.07, 145.70, 141.38, 131.56, 129.81, 125.30, 124.21, 124.01, 122.37, 116.81, 51.35, 31.68, 30.78, 30.61, 28.28.

## Example 34

25

**Synthesis of  
N-(5-(N-Methyl-N-3-pyridinesulfonylamino)pyrimidin-4-yl)-L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine *tert*-Butyl Ester**

30

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, XX and YY into L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine *tert*-butyl ester. 5-Nitrouracil (Aldrich) was converted via Method AAA into 2,4-dichloro-5-nitropyrimidine. L-4-(4-Methylpiperazin-1-ylcarbonyloxy)phenylalanine *tert*-butyl ester and 2,4-

-- 201 --

dichloro-5-nitropyrimidine were coupled via Method BBB, and the product of this coupling was sequentially converted via Methods CCC, DDD (using 3-chlorosulfonylpyridine) and EEE to give the title compound.

5

## Example 35

## Synthesis of

10

*N*-(5-(2-Trifluoromethylphenyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

15

*L*-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester. 4(3H)-Pyrimidinone (Aldrich) was sequentially converted via Methods N and O to 4-chloro-5-iodopyrimidine. *L*-4-(*N,N*-Dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4-chloro-5-iodopyrimidine were coupled via Method P and the coupled product was reacted with 2-trifluoromethylphenyl boronic acid (Aldrich) via Method Q. The product of this coupling was converted via Method HH to give the title compound.

20

Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 8.51 (s, 1H), 7.84 – 7.49 (m, 2H), 7.71 – 7.63 (m, 2H), 7.37 (d, 1H), 7.11 – 6.97 (m, 4H), 6.88 (d, 1H), 4.99 (s, 1H), 3.37 – 3.19 (m, 1H), 3.14 – 3.02 (m, 4H), 2.97 (s, 3H).

25

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ = 175.7, 175.5, 165.6, 161.9, 161.7, 158.6, 157.6, 157.5, 153.3, 153.1, 152.6, 152.5, 136.4, 136.2, 135.0, 134.9, 134.5, 133.1, 132.2, 131.9, 131.7, 128.9, 128.7, 127.8, 124.3, 123.6.

30

No Example 36

-- 202 --

## Example 37

**Synthesis of**  
***N*-(5-(2-*N,N*-Dimethylcarbamyloxy)pyrimidin-4-yl)-*L*-4-(*N,N*-**  
**dimethylcarbamyloxy)phenylalanine**

5                    *L*-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via  
Methods UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamyloxy)-  
phenylalanine *tert*-butyl ester. 4(3H)-Pyrimidinone (Aldrich) was  
10 sequentially converted via Methods N and O to 4-chloro-5-iodopyrimidine.  
*L*-4-(*N,N*-Dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4-chloro-  
5-iodopyrimidine were coupled via Method P and the coupled product was  
reacted with dimethylacrylamide (Aldrich) via Method U. The product of this  
reaction which was sequentially converted via Methods V and HH to give the  
15 title compound.

Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 8.56 (s, 1H), 8.06 (s, 1H), 7.32 (d, 2H),  
7.01 (d, 2H), 5.35 – 5.30 (m, 1H), 3.56 – 3.49 (m, 1H), 3.23 – 3.18 (m,  
1H), 3.11 (s, 3H), 3.02 (s, 3H), 2.99 (s, 3H), 2.97 (s, 3H), 2.88 (t, 2H),  
20 2.65 (t, 2H).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ = 174.5, 174.2, 152.7, 151.6, 142.6,  
136.5, 132.0, 123.8, 121.0, 57.8, 38.4, 37.9, 37.5, 36.9, 32.2, 24.6.

## 25                    Example 38

**Synthesis of**  
***N*-(5-(*N*-Methyl-*N*-3-(1-methylpyrazole)sulfonylamino)pyrimidin-4-yl)-*L*-**  
**4-(*N,N*-dimethylcarbamyloxy)phenylalanine Isopropyl Ester**

30                    *L*-Tyrosine (Aldrich) was sequentially converted via Methods LLL,  
UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine  
isopropyl ester. 5-Nitrouracil (Aldrich) was converted via Method AAA into  
2,4-dichloro-5-nitropyrimidine. *L*-4-(*N,N*-Dimethylcarbamyloxy)-  
35 phenylalanine isopropyl ester and 2,4-dichloro-5-nitropyrimidine were



-- 203 --

coupled via Method BBB, and the product of this coupling was sequentially converted via Methods CCC, DDD (using 1-methyl-3-chlorosulfonylpyrazole) and EEE to give the title compound.

Physical data were as follows:

5       <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ = 8.47 (s, 1H), 7.76 (s, 1H), 7.68 (bs, 2H), 7.19 (m, 2H), 7.04 (d, 2H), 6.17 (d, 1H), 5.03 (m, 2H), 3.95 (s, 3H), 3.31 - 3.12 (m, 2H), 3.08 (s, 3H), 3.06 (s, 3H), 2.99 (s, 3H), 1.24 (d, 3H), 1.21 (d, 3H).

10

### Example 39

#### Synthesis of

15    *N*-(6-Phenylpyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

*L*-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. *L*-4-(*N,N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 4,6-dichloropyrimidine (Aldrich) were coupled via Method Y and the coupled product was reacted with phenyl boronic acid (Aldrich) via Method Z. The product of this coupling was converted via Method HH to give the title compound.

Physical data were as follows:

25       <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 8.65 (s, 1H), 7.82 - 7.79 (m, 2H), 7.77 - 7.62 (m, 3H), 7.31 (d, 2H), 7.06 - 7.01 (m, 4H), 5.32 - 5.28 (m, 1H), 3.50 - 3.44 (m, 1H), 3.20 - 3.06 (m, 4H), 2.99 (s, 3H).

30       <sup>13</sup>C NMR (CD<sub>3</sub>OD): δ = 173.9, 165.7, 157.6, 154.9, 154.3, 152.8, 135.8, 134.6, 132.3, 132.2, 131.7, 129.2, 123.8, 104.6, 57.8, 38.8, 37.7, 37.5.

-- 204 --

## Example 40

## Synthesis of

***N*-(6-(2-Trifluoromethylphenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine**

5

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester and 4,6-dichloropyrimidine (Aldrich) were  
10 coupled via Method Y and the coupled product was reacted with 2-trifluoromethylphenyl boronic acid (Aldrich) via Method Z. The product of this coupling was converted via Method HH to give the title compound.

Physical data were as follows:

15 <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 8.46 (s, 1H), 7.95 – 7.82 (m, 1H), 7.73 – 7.67 (m, 2H), 7.50 – 7.48 (m, 1H), 7.29 (d, 2H), 7.03 (d, 2H), 6.65 (s, 1H), 5.05 (s, 1H), 3.39 (m, 1H), 3.16 – 3.12 (m, 4H), 3.00 (s, 3H).  
<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ = 176.0, 164.3, 158.8, 157.7, 152.6, 136.6, 139.0, 132.9, 132.1, 131.4, 130.1, 129.7, 128.2, 128.2, 123.6, 38.8,  
20 37.7, 37.5.

## Example 41

## Synthesis of

***N*-(6-(2-Hydroxymethylphenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine**

25

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester and 4,6-dichloropyrimidine (Aldrich) were  
30 coupled via Method Y and the coupled product was reacted with 2-(hydroxymethyl)phenyl boronic acid (Lancaster Synthesis) via Method Z.

-- 205 --

The product of this coupling was converted via Method HH to give the title compound.

Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  = 8.48 (s, 1H), 8.09 (s, 1H), 7.61 – 7.44 (m, 4H), 7.29 (d, 2H), 7.02 (d, 2H), 6.71 (s, 1H), 5.27 (s, 2H), 5.10 – 5.02 (m, 1H), 3.42 – 3.41 (m, 1H), 3.16 – 3.12 (m, 4H), 2.99 (s, 3H).

<sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$  = 175.7, 165.6, 164.7, 158.0, 157.6, 152.6, 141.6, 138.5, 136.7, 135.8, 132.2, 131.9, 131.7, 131.4, 131.3, 123.7, 64.9, 64.3, 38.9, 37.7, 37.5.

#### Example 42

##### Synthesis of *N*-(5-Cyclohexylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. Cyclohexanone (Aldrich) was sequentially converted via Methods MM, NN, OO and QQ into 4,6-dichloro-5-cyclohexylpyrimidine. L-4-(*N,N*-Dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4,6-dichloro-5-cyclohexylpyrimidine were coupled via Method RR, and the product of this coupling was sequentially converted via Methods SS and ZZ to give the title compound.

Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  = 8.41 (bs, 1H), 7.89 (bs, 1H), 7.21 (d, 2H), 6.94 (d, 2H), 5.12 (dd, 1H), 3.47 (dd, 1H), 3.19 (dd, 1H), 3.06 (s, 3H), 2.95 (s, 3H), 3.0 (m, 1H), 2.88 – 2.57 (bs, 1H), 2.5 (bs, 1H), 1.95 – 1.67(m, 1H).

-- 206 --

<sup>13</sup> C NMR (CD<sub>3</sub>OD):  $\delta$  = 175.68, 165.82, 156.87, 152.10, 151.88, 141.96, 136.30, 131.44, 125.38, 122.89, 57.86, 37.44, 36.81, 36.64, 36.30, 32.65, 32.13, 27.29, 27.25, 26.95.

5

## Example 43

## Synthesis of

*N*-(2-(*N*-Methyl-*N*-2-furanmethylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

10

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with N-methylfurfurylamine (Salor) via Method CC to give a product that was coupled with o-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

Physical data were as follows:

<sup>1</sup> H NMR (CD<sub>3</sub>OD):  $\delta$  = 7.43 – 7.35 (m, 2H), 7.35 – 7.2 (m, 2H), 7.2 – 7.0 (m, 4H), 7.0 – 6.9 (m, 2H), 6.42 (d, 1H), 6.39 (d, 1H), 4.85 (m, 1H), 3.3 – 3.1 (m, 7H), 3.09 (s, 3H), 2.98 (s, 3H), 2.16 (s, 3H), 1.89 (s, 3H).

25

## Example 44

## Synthesis of

*N*-(2-(*N*-Methyl-*N*-4-chlorophenylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

30

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich)

-- 207 --

were coupled via Method BB. The product of this reaction was reacted with *N*-methyl-4-chloroaniline (Aldrich) via Method CC to give a product that was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

5           Physical data were as follows:

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  = 8.17 (s, 1H), 7.56 – 7.34 (m, 8H), 7.1 – 6.97 (m, 4H), 3.50 (m, 2H), 3.13 (s, 3H), 2.1 (s, 3H), 2.17 (s, 3H), 1.94 (s, 3H).

10

#### Example 45

#### Synthesis of *N*-(5-(3-Thienyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

15

*L*-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. 4(3H)-Pyrimidinone (Aldrich) was sequentially converted via Methods N and O into 4-chloro-5-iodopyrimidine. *L*-4-(*N,N*-Dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4-chloro-5-iodopyrimidine were coupled via Method P and the coupled product was reacted with 3-thiophenyl boronic acid (Frontier Scientific, Inc.) via Method S. The product of this coupling was converted via Method KK to give the title compound.

25

Physical data were as follows:

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  = 8.62 (s, 1H), 8.13 (s, 1H), 7.62 (m, 1H), 7.59 (m, 1H), 7.20 (d, 2H), 7.09 (d, 1H), 7.01 (d, 2H), 3.47 – 3.13 (m, 2H), 3.13 (s, 3H), 2.97 (s, 3H).

30

$^{13}\text{C}$  NMR :  $\delta$  = 173.22, 162.83, 156.84, 152.17, 151.43, 141.46, 135.22, 131.54, 131.35, 129.96, 127.99, 127.90, 123.24, 117.13, 56.87, 36.82, 36.64.

## Example 46

**Synthesis of  
N-(5-(2-Thienyl)pyrimidin-4-yl)-L-4-(N,N-  
dimethylcarbamyloxy)phenylalanine**

5

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(N,N-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. 4(3H)-Pyrimidinone (Aldrich) was sequentially converted via Methods N and O to 4-chloro-5-iodopyrimidine. L-4-(N,N-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4-chloro-5-iodopyrimidine were coupled via Method P, and the coupled product was reacted with 2-thiophenyl boronic acid (Frontier Scientific, Inc.) via Method S. The product of this coupling was converted via Method KK to give the title compound.

15

Physical data were as follows:

<sup>1</sup> H NMR (CD<sub>3</sub>OD):  $\delta$  = 8.10 (s, 1H), 7.67 (s, 1H), 7.19 (d, 1H), 6.73 (m, 4H), 6.49 (m, 2H), 4.80 (m, 1H), 2.89 (m, 1H), 2.70 (m, 1H), 2.60 (s, 3H), 2.45 (s, 3H).

20

<sup>13</sup> C NMR (CD<sub>3</sub>OD):  $\delta$  = 173.07, 162.72, 156.80, 152.13, 151.74, 142.30, 135.07, 131.58, 131.14, 130.69, 130.38, 129.92, 123.19, 115.18, 56.94, 36.87, 36.81, 36.62, 28.74.

## Example 47

25

**Synthesis of  
N-(2-(N-Methyl-N-2-hydroxyethylamino)-5-(2-fluorophenyl)pyrimidin-4-yl)-L-4-(N,N-dimethylcarbamyloxy)phenylalanine**

30

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(N,N-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. L-4-(N,N-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with

-- 209 --

2-(*N*-methylamino)ethanol (Aldrich) via Method CC to give a product that was coupled with 2-fluorophenyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method KK to give the title compound.

5

#### Example 48

##### Synthesis of

10 ***N*-(5-(Piperidin-1-yl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine**

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-  
15 phenylalanine *tert*-butyl ester. Piperidine (Aldrich) was sequentially converted via Methods FFF, GGG and HHH into 4,6-dichloro-5-piperidin-1-ylpyrimidine. L-4-(*N,N*-Dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4,6-dichloro-5-piperidin-1-ylpyrimidine were coupled via Method III, and the product of this coupling was sequentially converted via Methods JJJ  
20 and ZZ into the title compound.

#### Example 49

25

##### Synthesis of

***N*-(5-(1-Propylbutyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine**

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via  
30 Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. 4-Heptanone (Aldrich) was sequentially converted via Methods MM, NN, OO and QQ into 4,6-dichloro-5-(1-propylbutyl)pyrimidine. L-4-(*N,N*-Dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4,6-dichloro-5-(1-propylbutyl)pyrimidine were coupled via

-- 210 --

Method RR, and the product of this coupling was sequentially converted via Methods SS and ZZ to give the title compound.

#### Example 50

5

#### Synthesis of *N*-(2-(*N*-Methyl-*N*-cyclobutylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N*,*N*-dimethylcarbamyloxy)phenylalanine

10 Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N*,*N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N*,*N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with  
15 N-methylcyclobutylamine (prepared by the Method of Giardina *et al.* J. Med. Chem. **1994**, 37(21), 3482 – 3491) via Method CC to give a product that was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

20

#### Example 51

#### Synthesis of *N*-(2-(*N*,*N*-Bis-(2-hydroxyethyl)amino)pyrimidin-4-yl)-L-4-(*N*,*N*-dimethylcarbamyloxy)phenylalanine

25

A byproduct was isolated by chromatography of the crude product of Example 52, and the byproduct was converted via Method KK into the title compound.

30

Physical data were as follows:

<sup>1</sup> H NMR (CD<sub>3</sub>OD): δ = 7.59 (d, 1H), 7.25 (d, 2H), 7.02 (d, 2H), 6.18 (d, 1H), 3.76 (brs, 8H), 2.97 (s, 8H).

<sup>13</sup> C NMR (CD<sub>3</sub>OD): δ = 174.1, 163.7, 155, 152, 142.1, 135.2, 131.3, 123.7, 99, 60.5, 56.8, 53.2, 37.5, 36.8, 36.6.

35



### Example 52

## Synthesis of

**5** *N*-(2-(*N,N*-bis(2-Hydroxyethyl)amino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine

10 Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with diethanolamine (Aldrich) via Method CC to give a product that was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The product of this  
15 coupling was converted via Method KK to give the title compound.

Physical data were as follows:

<sup>1</sup> H NMR (CD<sub>3</sub>OD): δ = 7.48 – 7.31 (m, 5H), 7.15 – 6.98 (m, 4H), 4.9 (m, 1H), 4.63 (m, 1H), 3.83 (d, 8H), 3.1 (s, 8H), 1.9 (d, 3H).

20 <sup>13</sup>C NMR (CD<sub>3</sub>OD): δ = 173.8, 162.3, 154.6, 152.6, 140.9, 139.6, 139.4, 135.9, 135.8, 132.2, 132.0, 131.4, 131.2, 131.1, 128, 123.2, 123.1, 66.8, 60.6, 56.9, 56.4, 53.2, 52.8, 36.8, 36.6, 36.3, 19.5.

### No Example 53

### Example 54

## Synthesis of

30 ***N*-(2-(*N*-Methyl-*N*-phenylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N*,*N*-dimethylcarbamoyloxy)phenylalanine**

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamyloxy)-

-- 212 --

phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with *N*-methylaniline (Aldrich) via Method CC to give a product that was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The product of this  
5 coupling was converted via Method ZZ to give the title compound.

Physical data were as follows:

<sup>1</sup> H NMR (CD<sub>3</sub>OD):  $\delta$  = 7.57 – 6.99 (m, 14H), 4.99 (m, 1H), 3.49 (s, 3H), 3.11 (m, 5H), 2.98 (s, 3H), 2.16 (s, 3H).

<sup>13</sup> C NMR (CD<sub>3</sub>OD):  $\delta$  = 183.07, 173.72, 173.49, 162.55, 156.82,  
10 153.97, 152.07, 142.25, 141.06, 140.91, 139.53, 139.40, 135.50, 135.39, 132.21, 132.16, 132.05, 131.52, 131.31, 130.53, 128.44, 128.11, 128.00, 123.13, 123.04, 113.18, 56.95, 56.49, 40.02, 39.96, 37.14, 36.83, 36.65, 19.56, 19.47.

15

#### Example 55

##### Synthesis of

20 *N*-(2-(Isopropoxy)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamyloxy)-  
25 phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this coupling was sequentially converted via Methods NNN, DD (using *o*-tolyl boronic acid, Aldrich) and ZZ to give the title compound.

Physical data were as follows:

30 <sup>1</sup> H NMR (CDCl<sub>3</sub>):  $\delta$  = 7.77 (bs, 1H), 7.40 – 6.8 (m, 9H), 6.43 (d, 0.5H) 6.27(d, 0.5H), 6.78 (m, 1H), 6.16 (m, 1H), 3.09 (s, 3H), 3.00 (s, 3H), 3.40 – 2.80 (m, 4H), 2.20 (s, 1.5H), 1.94 (s, 1.5H), 1.23 (m, 6H).

-- 213 --

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 176.28, 176.15, 160.03, 159.78, 154.77, 153.65, 150.01, 169.97, 139.20, 137.81, 137.64, 134.39, 134.25, 130.71, 130.47, 130.12, 129.15, 126.69, 126.46, 121.24, 121.18, 109.56, 56.81, 56.34, 63.19, 36.90, 36.56, 36.32, 22.19, 21.99, 21.95, 19.51, 19.27.$

5

## Example 56

## Synthesis of

*N*-(2-(*N*-Methyl-*N*-3-methylbutylamino)-5-(2-tolyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine

10

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester. *L*-4-(*N,N*-Dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with *N*-methyl-*N*-isoamylamine (Pfaltz-Bauer) via Method CC to give a product that was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

15

20

Physical data were as follows:

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 7.6$  (s, 0.5H), 7.56 (s, 0.5H), 7.30 – 6.80 (m, 8H) 6.30(bm, 1H), 7.00 – 6.00 (br, 1H), 4.63 (m, 1H), 3.09 (s, 3H), 3.01 (s, 6H), 3.80 – 2.80 (m, 4H), 2.13 (s, 1.5H), 1.90 (s, 1.5H), 1.61 (m, 1H), 1.51 (bs, 2H), 0.96 (d, 6H).

25

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 174.03, 173.87, 159.28, 159.04, 154.71, 153.67, 150.00, 142.10, 137.81, 137.53, 134.39, 134.22, 130.78, 130.58, 130.13, 128.96, 126.52, 126.30, 121.19, 121.13, 110.11, 109.91, 56.80, 56.40, 48.75, 36.55, 36.33, 35.80, 25.92, 22.54, 22.48, 19.53, 19.34.$

30

-- 214 --

## Example 57

**Synthesis of**  
***N*-(2-(*N*-Methylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine**

5

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with *N*-methylamine (Aldrich) via Method CC to give a product that was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

10

15

Physical data were as follows:

<sup>1</sup> H NMR (CDCl<sub>3</sub>): δ = 10.0 – 8.0 (br, 1H), 9.42 (bs, 1H), 8.24 (s, 1H), 7.4 – 6.8 (m, 10H), 5.93 (m, 1H), 4.85 (m, 1H), 3.2 – 2.8 (m, 1H), 3.37 (m, 1H), 3.12 (s, 1.5H), 3.11 (s, 1.5H), 3.03 (s, 1.5H), 3.02 (s, 1.5H), 2.95 (s, 3H), 2.13 (s, 1.5H), 1.83 (s, 1.5H).

20

## Example 58

**Synthesis of**  
***N*-(2-(2-tolyl)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine**

25

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with *o*-tolyl boronic acid (Aldrich) via Method KKK. The product of this coupling was converted via Method ZZ to give the title compound.

30

Physical data were as follows:

-- 215 --

<sup>1</sup> H NMR (CDCl<sub>3</sub>): δ = 8.14 (d, 1H), 7.68 (d, 1H), 7.4 – 6.8 (m, 12H), 5.42(m, 1H), 4.94 (m, 1H), 3.11 (s, 3H), 3.02 (s, 3H), 3.4 – 2.8 (m, 2H), 2.49 (s, 3H), 2.11 (s, 1.5H), 1.91 (s, 1.5H).

5

## Example 59

## Synthesis of

10 *N*-(2-(*N*-Methyl-*N*-2-hydroxyethylamino)-5-(2-tolyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. *L*-4-(*N,N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with 2-(methylamino)-ethanol (Aldrich) via Method CC to give a product that was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

15

Physical data were as follows:

20 <sup>1</sup> H NMR (CD<sub>3</sub>OD): δ = 7.4 – 6.94 (m, 4H), 4.82 (m, 1H), 3.8 (brs, 4H), 3.23/3.26 (s, rotamers, 3H), 2.98/3.7 (s, rotamers, 6H), 1.93/2.14 (s, rotamers, 3H).

25

## Example 60

## Synthesis of

30 *N*-(2-(*N*-Methyl-*N*-2-methylpropylamino)-5-(2-tolyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. *L*-4-(*N,N*-Dimethylcarbamyloxy)-

-- 216 --

phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with N-methyl isobutylamine (Aldrich) via Method CC to give a product that was coupled with o-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

Physical data were as follows:

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 10.5 – 9.8 (br, 1H), 7.63 (d, 1H), 7.3 – 6.8 (m, 8H), 6.35(m, 1H), 4.65 (m, 1H), 3.6 – 2.8 (m, 4H), 3.08 (s, 3H), 3.01 (s, 6H), 2.13(s, 1.5H), 2.06 (bs, 1H), 1.25 (s, 1.5H), 0.9 (s, 6H).

<sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 174.13, 173.97, 159.17, 158.9, 154.7, 153.99, 149.96, 142.00, 137.76, 137.53, 134.50, 134.33, 130.80, 130.58, 130.15, 128.95, 126.51, 126.30, 121.15, 121.11, 110.25, 109.99, 57.46, 56.90, 56.51, 36.89, 36.55, 36.32, 27.08, 19.87, 19.53, 19.38.

#### Example 61

##### Synthesis of

##### *N*-(2-(*N*-Methyl-*N*-propylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with N-methyl-N-propylamine (Aldrich) via Method CC to give a product that was coupled with o-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

Physical data were as follows:

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 10.5 – 9.5 (br, 1H), 7.6 (d, 1H), 7.38 – 6.7 (m, 8H), 6.3 (m, 1H), 4.7 (m, 1H), 3.7 – 3.0 (m, 4H), 3.09 (s, 3H), 3.01 (s, 6H), 2.13 (s, 1.5H), 1.92 (s, 1.5H), 1.59 (bs, 2H), 0.89 (bs, 3H).

-- 217 --

<sup>13</sup> C NMR (CDCl<sub>3</sub>): δ = 174.22, 174.06, 159.26, 159.0, 154.7, 153.76, 149.97, 142.22, 137.78, 137.53, 134.53, 134.36, 130.80, 130.73, 130.51, 130.12, 128.93, 126.50, 126.30, 121.16, 121.10, 110.13, 109.87, 56.90, 56.52, 51.72, 36.55, 36.33, 35.96, 20.45, 19.56, 19.37, 11.06.

5

## Example 62

## Synthesis of

10 *N*-(2-(*N,N*-Dimethylamino)-5-(2-tolyl)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. *L*-4-(*N,N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with *N,N*-dimethylamine (Aldrich) via Method CC to give a product that was coupled with *o*-tolylboronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

15  
20

Physical data were as follows:

<sup>1</sup> H NMR (CDCl<sub>3</sub>): δ = 11.0 – 9.5 (br, 1H), 7.62 (d, 1H), 7.3 – 6.8 (m, 8H), 6.22(m, 1H), 4.72 (m, 1H), 3.5 – 3.0 (m, 2H), 3.8 (s, 6H), 3.01 (s, 3H), 2.12 (s, 1.5H), 1.94 (s, 1.5H).

25 <sup>13</sup> C NMR (CDCl<sub>3</sub>): δ = 174.49, 174.3, 159.4, 158.93, 154.72, 149.93, 140.30, 137.75, 137.60, 134.67, 134.50, 130.92, 130.80, 130.51, 130.11, 128.87, 126.48, 126.32, 121.15, 121.08, 109.87, 109.69, 56.86, 56.49, 37.51, 36.87, 36.55, 36.34, 19.50, 19.38.

30

-- 218 --

## Example 63

Synthesis of  
*N*-(2-(*N*-Methyl-*N*-cyclohexylamino)-5-(3-pyridyl)pyrimidin-4-yl)-L-4-  
(*N,N*-dimethylcarbamyloxy)phenylalanine

5 Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamyloxy)-  
10 phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with *N*-methyl-*N*-cyclohexylamine (Aldrich) via Method CC to give a product that was coupled with 3-pyridyl boronic acid 1,3-propanediol cyclic ester (Lancaster Synthesis) via Method DD. The product of this coupling was  
15 converted via Method HH to give the title compound.

Physical data were as follows:

<sup>1</sup> H NMR (CD<sub>3</sub>OD):  $\delta$  = 8.83 – 8.78 (m, 1H), 8.56 (brs, 1H), 8.09 – 7.95 (m, 2H), 7.76 – 7.73 (m, 1H), 7.22 (d, 2H), 7.06 (d, 2H), 4.85 (m, 1H), 3.45 – 3.38 (m, 1H), 3.18 – 3.11 (m, 4H), 3.06 (s, 3H), 2.99 (sm,  
20 overlapping 4H), 1.92 (m, 2H), 1.76 – 1.57 (m, 8H).

<sup>13</sup> C NMR (CD<sub>3</sub>OD):  $\delta$  = 173.7, 161.5, 161.4, 160.9, 157.0, 152.0, 146.0, 145.7, 145.6, 143.3, 136.0, 132.2, 131.3, 128.1, 123.4, 107.8, 57.8, 57.4, 36.8, 36.6, 36.1, 30.6, 30.0, 26.4, 26.2.

25

## Example 64

Synthesis of  
*N*-(5-(2-Phenyl-2,2-difluoroethyl)pyrimidin-4-yl)-L-4-(*N,N*-  
dimethylcarbamyloxy)phenylalanine

30

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. 1-Trifluoromethanesulfonyloxy-2,2-difluoro-2-phenylethane was sequentially converted via Methods LL, OO and QQ into



-- 219 --

4,6-dichloro-5-(2,2-difluoro-2-phenylethyl)pyrimidine. L-4-(*N,N*-Dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester and 4,6-dichloro-5-(2,2-difluoro-2-phenylethyl)pyrimidine were coupled via Method RR, and the product of this coupling was sequentially converted via Methods TT and ZZ to give the title compound.

Physical data were as follows:

<sup>1</sup> H NMR (CD<sub>3</sub>OD):  $\delta$  = 8.37 (s, 1H), 7.79 (s, 1H), 7.44 (s, 5H), 7.25 (d, 2H), 6.98 (d, 2H), 5.07 (dd, 1H), 3.62 – 3.32 (m, 3H), 3.14 (dd, 1H) 3.08 (s, 3H), 2.96 (s, 3H).

#### Example 65

##### Synthesis of

*N*-(5-(2-Phenyl-2,2-difluoroethyl)-6-chloropyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester. 1-Trifluoromethanesulfonyloxy-2,2-difluoro-2-phenylethane was sequentially converted via Methods LL, OO and QQ into 4,6-dichloro-5-(2,2-difluoro-2-phenylethyl)pyrimidine. L-4-(*N,N*-Dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester and 4,6-dichloro-5-(2,2-difluoro-2-phenylethyl)pyrimidine were coupled via Method RR, and the product of this coupling was converted via Method ZZ to give the title compound.

Physical data were as follows:

<sup>1</sup> H NMR (CD<sub>3</sub>OD):  $\delta$  = 8.18 (s, 1H), 7.42 – 7.41 (m, 5H), 7.26 (d, 2H), 7.0 (d, 2H), 5.03 (dd, 1H), 3.72 – 3.45 (m, 2H), 3.34 (dd, 1H), 3.19 (dd, 1H), 3.08 (s, 3H), 2.96 (s, 3H).

-- 220 --

## Example 66

**Synthesis of**  
***N*-(5-(2-Phenylethyl)pyrimidin-4-yl)-L-4-(*N,N*-**  
**dimethylcarbamyloxy)phenylalanine**

5

4(3H)-Pyrimidinone (Aldrich) was sequentially converted via Methods N and O into 4-chloro-5-iodopyrimidine. L-Phenylalanine *tert*-butyl ester hydrochloride (Bachem) and 4-chloro-5-iodopyrimidine were coupled via  
10 Method P. The product of this reaction was converted via Method W to a product that was sequentially converted via Methods X and HH to give the title compound.

Physical data were as follows:

<sup>1</sup> H NMR (CD<sub>3</sub>OD): δ = 8.55 (d, 1H), 7.64 (d, 1H), 7.35 – 7.19  
15 (m, 8H), 7.01 – 6.98 (m, 2H), 5.46 – 5.41 (m, 1H), 5.34 – 3.60 (m, 1H),  
3.29 – 3.23 (m, 1H), 2.94 – 2.75 (m, 4H).

<sup>13</sup> C NMR (CD<sub>3</sub>OD): δ = 174.3, 164.3, 151.5, 141.8, 141.7,  
139.2, 131.0, 130.6, 130.5, 130.4, 128.9, 128.4, 120.6, 57.8, 38.4, 34.0,  
30.7.

20

## Example 67

**Synthesis of**  
***N*-(2-(*N*-Methyl-*N*-cyclohexylamino)pyrimidin-4-yl)-L-4-(*N,N*-**  
25 **dimethylcarbamyloxy)phenylalanine**

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamyloxy)-  
30 phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with *N*-methyl-*N*-cyclohexylamine (Aldrich) via Method CC to give a product which was sequentially converted via Methods MMM and ZZ to give the title compound.

-- 221 --

Physical data were as follows:

<sup>1</sup> H NMR (CDCl<sub>3</sub>): δ = 11.20 (bs, 2H), 8.44 (s, 1H), 7.76 (bs, 1H), 7.50 (br, 1H), 7.18 (d, 2H), 6.96 (d, 2H), 5.91 (bs, 1H), 4.83 (bs, 1H), 4.53 (br, 1H), 3.20 (m, 2H), 3.08 (s, 3H), 2.98 (s, 6H), 2.00 – 1.00 (m, 10H).

<sup>13</sup> C NMR (CDCl<sub>3</sub>): δ = 176.18, 171.50, 167.75, 162.44, 156.31, 154.49, 151.52, 135.83, 131.61, 122.85, 58.04, 56.87, 38.02, 37.79, 31.16, 31.00, 26.68.

#### Example 68

#### Synthesis of *N*-(5-Propylpyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine

*L*-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester. 4(3H)-Pyrimidinone (Aldrich) was sequentially converted via Methods N and O into 4-chloro-5-iodopyrimidine. *L*-4-(*N,N*-Dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester and 4-chloro-5-iodopyrimidine were coupled via Method P, and the product of this coupling was sequentially converted via Methods II, JJ and KK to give the title compound.

Physical data were as follows:

<sup>1</sup> H NMR (CD<sub>3</sub>OD): δ = 8.51 (s, 1H), 7.97 (s, 1H), 7.26 (d, 2H), 6.97 (d, 2H), 5.36 (m, 1H), 3.51 (m, 1H), 3.23 (m, 1H), 3.16 (s, 3H), 2.95 (s, 3H), 2.47 (m, 2H), 1.57 (m, 2H), 0.99 (m, 3H).

<sup>13</sup> C NMR (CD<sub>3</sub>OD): δ = 173.48, 163.61, 151.97, 150.75, 140.68, 135.74, 133.14, 131.30, 123.02, 120.85, 56.96, 36.99, 36.76, 36.58, 29.87, 21.02, 13.67.

-- 222 --

## Example 69

**Synthesis of**  
***N*-(5-(2-Methoxyphenyl)pyrimidin-4-yl)-L-4-(*N,N*-**  
**dimethylcarbamyloxy)phenylalanine**

5 L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via  
Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-  
phenylalanine *tert*-butyl ester. 4(3H)-Pyrimidinone (Aldrich) was  
10 sequentially converted via Methods N and O to 4-chloro-5-iodopyrimidine.  
L-4-(*N,N*-Dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4-chloro-  
5-iodopyrimidine were coupled via Method P and the coupled product was  
reacted with 2-methoxyphenyl boronic acid (Lancaster Synthesis) via Method  
Q. The product of this coupling was converted via Method HH to give the  
15 title compound.

Physical data were as follows:

<sup>1</sup> H NMR (CD<sub>3</sub>OD):  $\delta$  = 8.64 (s, 1H), 8.05 (s, 1H), 7.61 – 7.55  
(m, 1H), 7.27 – 7.13 (m, 5H), 6.99 (d, 2H), 5.36 – 5.32 (m, 1H), 3.73 (s,  
3H), 3.46 – 3.40 (m, 1H), 4.20 – 3.13 (m, 4H), 3.02 (s, 3H).

20 <sup>13</sup> C NMR (CD<sub>3</sub>OD):  $\delta$  = 173.8, 163.5, 159.5, 157.5, 152.8,  
152.1, 143.0, 135.9, 134.2, 133.9, 132.2, 123.8, 123.4, 120.5, 120.0,  
113.7, 57.5, 57.1, 37.9, 37.7, 37.5.

25

## Example 70

**Synthesis of**  
***N*-(5-(2-Fluorophenyl)pyrimidin-4-yl)-L-4-(*N,N*-**  
**dimethylcarbamyloxy)phenylalanine**

30

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via  
Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-  
phenylalanine *tert*-butyl ester. 4(3H)-Pyrimidinone (Aldrich) was

5 The product of this coupling was converted via Method HH to give the title compound.

### Example 71

10 **Synthesis of**  
***N*-(2-(*N*-Methyl-*N*-isopropylamino)-5-(2-tolyl)pyrimidin-4-yl)-*L*-4-(*N*,*N*-dimethylcarbamyloxy)phenylalanine**

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via  
15 Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-  
phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamyloxy)-  
phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich)  
were coupled via Method BB. The product of this reaction was reacted with  
*N*-methyl-*N*-isopropylamine (Aldrich) via Method CC to give a product that  
20 was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The  
product of this coupling was converted via Method ZZ to give the title  
compound.

**Physical data were as follows:**

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 10.5 – 9.5 (br, 1H), 7.59 (d, 1H), 7.30 – 6.70 (m, 8H), 6.3 (m, 1 H), 4.92 (bs, 1H), 4.7 (m, 1H), 3.50 – 3.0 (m, 2H), 3.08 (s, 3H), 3.00 (s, 3H), 2.83 (s, 3H), 2.13 (s, 1.5H), 1.93 (s, 1.5H) 1.15 (d, 6H).

<sup>13</sup> C NMR (CDCl<sub>3</sub>): δ = 174.31, 174.15, 159.21, 158.95, 154.70, 153.41, 149.92, 141.98, 137.79, 137.56, 134.59, 134.41, 130.59, 130.17, 128.95, 126.51, 126.32, 121.15, 110.26, 110.02, 56.87, 56.50, 46.86, 36.82, 36.55, 36.31, 28.18, 19.50, 19.39.

-- 224 --

## Example 72

**Synthesis of**  
***N*-(2-(*N*-Isopropylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine**

5

Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with isopropylamine (Aldrich) via Method CC to give a product that was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

15

Physical data were as follows:

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 9.57 (s, 1H), 8.31 (s, 1H), 7.40 – 6.80 (m, 8H), 6.19 (m, 1H), 4.79 (m, 1H), 4.15 (m, 1H), 3.4 – 3.0 (m, 2H), 3.10 (s, 3H), 3.01 (s, 3H), 2.16 (s, 1.5H), 1.41 (s, 1.5H), 1.24 (s, 6H).

20

<sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 176.07, 175.8, 166.23, 160.23, 159.99, 154.79, 153.50, 158.06, 139.38, 137.86, 137.66, 134.10, 133.93, 130.77, 130.61, 130.26, 130.01, 129.25, 126.71, 126.50, 121.46, 121.36, 109.59, 109.37, 56.77, 56.22, 43.31, 36.57, 36.34, 22.12, 21.96, 19.47, 19.22.

25

No Examples 73-77

## Example 78

**Synthesis of**  
***N*-(5-(2-Phenylethyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine Isopropyl Ester**

30

L-Tyrosine was sequentially converted via Methods LLL, UU, VV and WW into L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine isopropyl ester.

-- 225 --

4(3H)-Pyrimidinone (Aldrich) was sequentially converted via Methods N and O to 4-chloro-5-iodopyrimidine. L-4-(*N,N*-Dimethylcarbamoyloxy)phenylalanine isopropyl ester and 4-chloro-5-iodopyrimidine were coupled via Method P, and the coupled product was converted via Method OOO to a product, which was converted via Method X to give the title compound.

Physical data were as follows:

<sup>1</sup> H NMR (CDCl<sub>3</sub>): δ = 8.50 (s, 1H), 7.91 (s, 1H), 7.31 – 7.20 (m, 3H), 7.42 – 7.00 (m, 6H), 5.19 – 5.17 (m, 1H), 5.08 – 5.02 (m, 2H), 3.23 – 3.17 (m, 2H), 3.06 (s, 3H), 2.99 (s, 3H), 2.83 – 2.78 (m, 2H), 2.65 – 2.60 (m, 2H), 1.75 – 1.23 (m, 6H).

<sup>13</sup> C NMR (CDCl<sub>3</sub>): δ = 171.8, 159.2, 156.7, 153.5, 150.7, 140.5, 130.3, 128.7, 128.5, 126.4, 121.8, 117.1, 69.4, 54.2, 36.9, 36.6, 36.5, 33.6, 29.8, 21.7, 21.6.

#### Example 79

##### Synthesis of *N*-(3-(*N*-Methyl-*N*-4-toluenesulfonylamino)pyrazin-2-yl)-*L*-phenylalanine Isopropyl Ester

*L*-Phenylalanine (Aldrich) was converted via Method LLL to *L*-phenylalanine isopropyl ester hydrochloride. 2,3-Dichloropyrazine (Lancaster) was converted via Method QQQ and RRR into 2-chloro-3-nitropyrazine. *L*-Phenylalanine isopropyl ester hydrochloride and 2-chloro-3-nitropyrazine were coupled via Method BBB, and the product of this coupling was sequentially converted via Methods CCC, DDD and EEE to give the title compound.

Physical data were as follows:

<sup>1</sup> H NMR (CDCl<sub>3</sub>): δ = 7.91 (d, 1H), 7.59 (d, 2H), 7.51 (d, 1H), 7.31 – 7.23 (m, 7H), 6.08 (d, 1H), 5.01 – 4.97 (m, 1H), 4.92 – 4.89 (m, 1H) 3.24 (d, 2H), 2.97 (s, 3H), 2.43 (s, 3H), 1.21 – 1.12 (m, 6H).

-- 226 --

<sup>13</sup> C NMR (CDCl<sub>3</sub>): δ = 167.32, 147.440, 139.85, 137.38, 133.25, 131.98, 128.68, 126.17, 125.17, 125.06, 124.41, 124.11, 122.58, 64.38, 50.65, 33.49, 32.41, 17.16, 17.08, 17.03.

5

## Example 80

**Synthesis of  
N-(5-(2-Phenylethyl)pyrimidin-4-yl)-L-phenylalanine Isopropyl Ester**

10

L-Phenylalanine isopropyl ester hydrochloride was prepared by Method LLL. 4(3H)-Pyrimidinone (Aldrich) was sequentially converted via Methods N and O to 4-chloro-5-iodopyrimidine. L-Phenylalanine isopropyl ester hydrochloride and 4-chloro-5-iodopyrimidine were coupled via Method P and the coupled product sequentially converted via Methods OOO and X to give the title compound.

15

Physical data were as follows:

<sup>1</sup> H NMR (CDCl<sub>3</sub>): δ = 8.51 (s, 1H), 7.92 (s, 1H), 7.30 – 7.15 (m, 5H), 7.14 – 7.06 (m, 4H), 5.16 (m, 1H), 5.09 – 5.01 (m, 2H), 3.31 – 3.16 (m, 2H), 2.79 – 2.74 (m, 2H), 2.62 – 2.57 (m, 2H), 1.15 – 1.20 (m, 6H).

20

<sup>13</sup> C NMR (CDCl<sub>3</sub>): δ = 171.7, 159.1, 156.7, 153.5, 140.5, 136.1, 129.4, 128.6, 128.5, 128.3, 127.1, 126.4, 117.0, 69.3, 54.2, 37.6, 33.7, 30.0, 21.7, 21.6.

## No Example 81

25

## Example 82

**Synthesis of  
N-(5-(N-Methyl-N-3-pyridinesulfonylamino)pyrimidin-4-yl)-L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine**

30

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, XX and YY into L-4-(4-methylpiperazin-1-ylcarbonyloxy)-



-- 227 --

phenylalanine *tert*-butyl ester. 5-Nitrouracil (Aldrich) was converted via Method AAA into 2,4-dichloro-5-nitropyrimidine. L-4-(4-Methylpiperazin-1-ylcarbonyloxy)phenylalanine *tert*-butyl ester and 2,4-dichloro-5-nitropyrimidine were coupled via Method BBB, and the product of this coupling was sequentially converted via Methods CCC, DDD (using 3-chlorosulfonylpyridine), EEE and ZZ to give the title compound.

No Examples 83-84

10 Example 85

**Synthesis of  
*N*-(2-(*N*-Methyl-*N*-cyclohexylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine**

15 L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N* dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. L-4-(*N,N*-Dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester and 5-bromo-2,4-dichloropyrimidine (Aldrich) were coupled via Method BB. The product of this reaction was reacted with *N*-methyl-*N*-cyclohexylamine (Aldrich) via Method CC to give a product that was coupled with *o*-tolyl boronic acid (Aldrich) via Method DD. The product of this coupling was converted via Method ZZ to give the title compound.

25 Physical data were as follows:

<sup>1</sup> H NMR (CDCl<sub>3</sub>): δ = 10.0 – 9.08 (br, 1H), 7.55 (s, 0.5H), 7.52 (s, 0.5H), 7.20 – 6.31 (m, 8H), 6.36 (br, 1H), 4.69 (m, 2H), 3.40 (m, 1H), 3.15 (m, 1H), 3.06 (brs, 3H), 2.98 (brs, 3H), 2.84 (brs, 3H), 2.11 (s, 1.5H), 2.00 – 1.00 (brm, 11.5 H).

30 <sup>13</sup> C NMR (CDCl<sub>3</sub>): δ = 164.10, 159.20, 159.00, 154.79, 153.50, 150.03, 137.68, 137.48, 134.48, 130.66, 130.22, 129.01, 126.62, 126.40,

-- 228 --

121.16, 110.20, 57.00, 56.58, 55.50, 36.62, 36.39, 29.91, 29.52, 25.41,  
19.60, 19.65.

No Examples 86-87

5

Example 88

10

Synthesis of  
*N*-(5-(2-Tolyl)pyrimidin-4-yl)-L-4-(*N,N*-  
dimethylcarbamoyloxy)phenylalanine Isopropyl Ester

L-Tyrosine (Aldrich) was sequentially converted via Methods LLL,  
UU, VV and WW into L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine  
isopropyl ester. 4(3H)-Pyrimidinone (Aldrich) was sequentially converted via  
Methods N and O into 4-chloro-5-iodopyrimidine. L-4-(*N,N*-  
dimethylcarbamoyloxy)-phenylalanine isopropyl ester and 4-chloro-5-  
iodopyrimidine were coupled via Method P. The product of this coupling  
was reacted with o-tolyl boronic acid via Method Q to afford the title  
compound.

20

Physical data were as follows:

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 8.58 (s, 1H), 7.99 (s, 1H), 7.76 – 7.33 (m,  
3H), 7.13 (m, 0.5H), 7.03 – 6.95 (m, 4H), 4.97 – 4.87 (m, 3H), 3.08 –  
2.99 (m, 8H), 2.09 (s, 2H), 1.92 (s, 1.5H), 1.24 – 1.12 (m, 6H).

25

<sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 171.4, 171.2, 158.8, 158.5, 157.5, 154.7,  
153.6, 153.5, 150.5, 137.1, 137.0, 132.9, 132.3, 132.5, 130.8, 130.7,  
130.0, 129.8, 129.7, 128.9, 126.6, 126.5, 121.6, 119.5, 119.4, 69.0, 54.5,  
54.0, 36.9, 36.8, 36.6, 36.4, 21.65, 21.60, 19.3, 19.2.

30

-- 229 --

## Example 89

Synthesis of  
*N*-(5-(3-Nitrophenyl)pyrimidin-4-yl)-*L*-4-(*N,N*-  
dimethylcarbamyloxy)phenylalanine

5

*L*-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into *L*-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. 4(3H)-pyrimidinone (Aldrich) was  
10 sequentially converted via Methods N and O to 4-chloro-5-iodopyrimidine. *L*-4-(*N,N*-Dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4-chloro-5-iodopyrimidine were coupled via Method P, and the coupled product was reacted with 3-nitrophenyl boronic acid (Aldrich) via Method T. The product of this coupling was converted via Method HH to give the title compound.

15

Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 8.67 (s, 1H), 8.41 - 8.38 (m, 1H), 8.28 - 8.27 (m, 1H), 8.17 (s, 1H), 7.82 - 7.77 (m, 1H), 7.67 - 7.65 (m, 1H), 7.20 (d, 2H), 7.02 (d, 2H), 5.33 - 5.28 (m, 1H), 3.47 - 3.411 (m, 1H), 3.12 - 3.04 (m, 4H), 2.97 (s, 3H).

20

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ = 173.7, 163.6, 157.6, 152.8, 152.7, 151.2, 143.8, 137.4, 136.3, 134.2, 133.1, 132.2, 126.7, 126.3, 124.0, 120.3, 58.0, 37.7, 37.6, 37.5.

25

No Example 90

30

35

-- 230 --

## Example 91

Synthesis of  
*N*-(5-(3-Pyridyl)pyrimidin-4-yl)-L-4-(*N,N*-  
dimethylcarbamyloxy)phenylalanine

5  
10  
15  
L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. 4(3H)-Pyrimidinone (Aldrich) was sequentially converted via Methods N and O to 4-chloro-5-iodopyrimidine. L-4-(*N,N*-Dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4-chloro-5-iodopyrimidine were coupled via Method P, and the coupled product was reacted with 3-pyridyl boronic acid 1,3-propanediol cyclic ester (Lancaster Synthesis) via Method Q. The product of this coupling was converted via Method HH to give the title compound.

## Example 92

Synthesis of  
*N*-(5-(2-Phenylethyl)pyrimidin-4-yl)-L-4-(*N,N*-  
dimethylcarbamyloxy)phenylalanine

20  
25  
30  
L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamyloxy)-phenylalanine *tert*-butyl ester. 4(3H)-Pyrimidinone (Aldrich) was sequentially converted via Methods N and O into 4-chloro-5-iodopyrimidine. L-4-(*N,N*-Dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4-chloro-5-iodopyrimidine were coupled via Method P. The product of this reaction was sequentially converted via Methods W, X, and HH to give the title compound.

Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 8.52 (s, 1H), 7.67 (s, 1H), 7.34 – 7.19 (m, 5H), 7.08 – 6.99 (m, 4H), 5.50 – 5.42 (m, 1H), 5.59 – 5.53 (m, 1H), 3.26 – 3.21 (m, 1H), 3.09 (s, 2H), 2.99 (s, 3H), 2.94 – 2.85 (m, 4H).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ = 174.2, 164.2, 157.5, 152.7, 151.4, 141.8, 141.7, 136.5, 132.0, 130.5, 130.4, 128.4, 123.8, 120.5, 57.8, 37.9, 37.6, 37.5, 34.1, 30.6.

### Example 93

10 **Synthesis of**  
***N*-(2-*N,N*-Dimethylamino-5-(*N*-methyl-*N*-4-**  
**toluenesulfonylamino)pyrimidin-4-yl)-*L*-phenylalanine**

5-Nitrouracil (Aldrich) was converted via Method AAA into 2,4-dichloro-5-nitropyrimidine. L-Phenylalanine *tert*-butyl ester (Bachem) and 2,4-dichloro-5-nitropyrimidine were coupled via Method BBB, and the product of this coupling was sequentially converted via Methods SSS (using dimethylamine), CCC, DDD, EEE and ZZ to give the title compound.

**Physical data were as follows:**

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 8.15 (s, formate), 7.65 (m, 2H), 7.41 (d, 2H), 7.40 - 7.19 (m, 5H), 7.02 - 6.92 (m, 1H), 4.90 (m, 1H), 3.40 - 3.10 (m, 2H), 3.09 - 2.92(m, 9H), 2.43 (s, 3H).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ = 177.07, 159.64, 154.70, 152.25, 144.10, 141.97, 141.33, 140.25, 132.57, 129.02, 125.21, 124.82, 123.57, 123.42, 121.88, 107.64, 51.08, 33.71, 32.72, 31.76, 15.49.

### Example 94

### Synthesis of N-(5-(2-Tolyl)pyrimidin-4-yl)-L-4-(N,N- dimethylcarbamoyloxy)phenylalanine

L-Tyrosine *tert*-butyl ester (Bachem) was sequentially converted via Methods UU, VV and WW into L-4-(*N,N*-dimethylcarbamoyloxy)-phenylalanine *tert*-butyl ester. 4(3H)-Pyrimidinone (Aldrich) was sequentially converted via Methods N and O into 4-chloro-5-iodopyrimidine.

L-4-(*N,N*-Dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and 4-chloro-5-iodopyrimidine were coupled via Method P and the coupled product was reacted with *o*-tolyl boronic acid (Aldrich) via Method Q. The product of this coupling was converted via Method HH to give the title compound.

5        Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ = 8.75 – 8.65 (d, 1H), 8.05 – 8.03 (d, 1H), 7.51 – 7.35 (m, 3H), 7.26 – 7.11 (m, 3H), 7.02 – 6.97 (m, 2H), 5.38 – 5.27 (m, 2H), 3.50 – 3.39 (m, 1H), 3.21 – 3.07 (m, 4H), 3.02 (s, 3H), 2.21 – 1.93 (s, 3H).

10        <sup>13</sup>C NMR (CD<sub>3</sub>OD) : δ = 173.8, 173.6, 164.0, 163.8, 157.5, 152.7, 152.6, 143.0, 142.8, 139.7, 139.5, 136.1, 135.9, 133.2, 133.0, 132.4, 132.2, 132.1, 131.9, 131.1, 129.0, 128.9, 123.8, 123.7, 122.2, 122.0, 57.6, 57.4, 37.8, 37.7, 37.5, 37.4, 20.3, 20.2.

15        Additionally, using the procedures described herein and the appropriate starting materials, the following additional compounds can be prepared:

20        *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2-methoxyphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 95),

*N*-(2-(*N*-methyl-*N*-isopropylamino)-5-(2-fluorophenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 96),

25        *N*-(2-(*N*-methyl-*N*-isopropylamino)-5-(2-fluorophenyl)pyrimidin-4-yl)-L-4-(2-methoxyphenyl)phenylalanine (Example 97),

30        *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2,6-difluorophenyl)pyrimidin-4-yl)-L-4-(2,6-difluorophenyl)phenylalanine (Example 98),

*N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2-hydroxymethylphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 99),

35        *N*-(2-(*N,N*-bis-(2-hydroxyethyl)amino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 100),

- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2-trifluoromethylphenyl)pyrimidin-4-yl)-L-4-(2-cyanophenyl)phenylalanine (Example 101),
- 5      *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(3-thienyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 102),
- 10      *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2-thienyl)pyrimidin-4-yl)-L-4-(4-trifluoromethylphenyl)phenylalanine (Example 103),
- 15      *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(3-pyridyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 104),
- 20      *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(3-nitrophenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 105),
- 25      *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2,6-dichlorophenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 106),
- 30      *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(4-pyridyl)pyrimidin-4-yl)-L-4-(3-hydroxymethylphenyl)phenylalanine (Example 107),
- 35      *N*-(2-(*N*-ethyl-*N*-isopropylamino)-5-(2,6-dimethoxyphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 108),
- 40      *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2,3-dichlorophenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 109),
- 45      *N*-(2-(*N*-methyl-*N*-ethylamino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(2-cyanophenyl)phenylalanine (Example 110),
- 50      *N*-(2-(*N*-methyl-*N*-isopropylamino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(3-pyridyl)phenylalanine (Example 111),
- 55      *N*-(2-(*N,N*-bis-(2-hydroxyethyl)amino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(2-cyanophenyl)phenylalanine (Example 112),
- 60      *N*-(2-(*N*-methyl-*N*-(1-methylpiperidin-4-yl)amino)-5-(2-cyanophenyl)pyrimidin-4-yl)-L-4-(2,6-difluorophenyl)phenylalanine (Example 113),
- 65      *N*-(2-(*N*-ethyl-*N*-isopropylamino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(*o*-tolyl)phenylalanine (Example 114),

- N*-(2-(*N*-methyl-*N*-4-chlorophenylamino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 115),
- 5      *N*-(5-(*N*-methyl-*N*-2-(phenyl)ethylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine (Example 116),
- 10      *N*-(5-(*N*-methyl-*N*-hexylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine (Example 117),
- 15      *N*-(5-(*N*-methyl-*N*-isopropylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine (Example 118),
- 20      *N*-(5-(*N*-methyl-*N*-isopropylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine (Example 119),
- 25      *N*-(5-(*N*-methyl-*N*-*tert*-butylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine (Example 120),
- 30      *N*-(5-(*N*-ethyl-*N*-isopropylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine (Example 121),
- 35      *N*-(5-(*N*-methyl-*N*-2-(4-pyridyl)ethyl-pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine (Example 122),
- 40      *N*-(5-(*N*-methyl-*N*-2-(phenyl)ethylamino)pyrimidin-4-yl)-L-4-(4-(2,6-dimethoxyphenyl)phenylalanine (Example 123),
- 45      *N*-(5-(*N*-methyl-*N*-hexylamino)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 124),
- 50      *N*-(5-(*N*-methyl-*N*-isopropylamino)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 125),
- 55      *N*-(5-(*N*-methyl-*N*-isopropylamino)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 126),
- 60      *N*-(5-(*N*-methyl-*N*-*tert*-butylamino)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 127),
- 65      *N*-(5-(*N*-ethyl-*N*-isopropylamino)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 128),
- 70      *N*-(5-(*N*-methyl-*N*-2-(4-pyridyl)ethyl-pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine (Example 129)



*N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-ethylpyrimidin-4-yl)-*L*-4-(*N*,*N*-dimethylcarbamoyloxy)phenylalanine (Example 130).

5

### Example 131

## Synthesis of

**N-(4-(N,N-Di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-L-tyrosine**

10      **Step A:      Preparation of 3,4-Diethyloxy-1-oxo-1,2,5-thiadiazole and 3,4-Diethyloxy-1,1-dioxo-1,2,5-thiadiazole**

The title intermediates were prepared according to the procedures described in R. Y. Wen *et al*, *J Org Chem.*, (1975) 40, 2743; and R. Y. Wen *et al*, *Org Prep Proceed.*, (1969) 1, 255.

15

**Step B: Preparation of 4-(N,N-Di-*n*-hexylamino)-3-ethoxy-1,1-dioxo-1,2,5-thiadiazole**

20 Dihexylamine (90 mg, 0.48 mmol) was added to a solution of 3,4-diethyloxy-1,1-dioxo-1,2,5-thiadiazole (100 mg, 0.48 mmol) in ethanol (5 mL) and the reaction stirred overnight at room temperature. The solvent was removed under reduced pressure and the residue absorbed onto silica gel, and purified by flash column chromatography (silica, hexane:EtOAc 3:1) to yield the title intermediate (120 mg, 72%).

25            Physical data were as follows:

MS ( EI, m/e) 345.

**Step C: Preparation of N-(4-(N,N-Di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-L-tyrosine tert-Butyl Ester**

30

A solution of 4-(N,N-di-*n*-hexylamino)-3-ethoxy-1,1-dioxo-1,2,5-thiadiazole (400 mg, 1.02 mmol) and L-tyrosine t-butyl ester (261 mg, 1.1 mmol) in EtOH (10 mL) was stirred at room temperature for 36 hrs. The solvent was removed under reduced pressure residue purified by flash column

-- 236 --

chromatography (silica, hexane:EtOAc 3:1 then 1:1) to give the title compound as a white waxy solid (400 mg, 73%).

Physical data were as follows:

Anal. Calc'd for  $C_{27}H_{44}N_4O_5$  S.0.55EtOAc: C, 59.93; H, 8.34; N, 9.57. Found: C, 59.84; H, 8.44; N, 9.62.

**Step D: Preparation of N-(4-(N,N-Di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-L-tyrosine**

The compound from Step C (100 mg, 0.19 mmol) was dissolved in formic acid and the mixture stirred at room temperature for 36 hrs. Excess formic acid was removed under reduced pressure to yield the title compound as a white solid (90 mg, 98 %).

Physical data were as follows:

Anal. Calc'd for  $C_{23}H_{36}N_4O_5$  S: C, 57.48; H, 7.55; N, 11.66. Found: C, 57.04; H, 7.23; N, 11.38.

**Example 132**

**Synthesis of  
N-(4-(N,N-Di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-L-4-(N,N-dimethylcarbamoyloxy)phenylalanine**

**Step A: Preparation of N-(4-(N,N-Di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-L-4-(N,N-dimethylcarbamoyloxy)-phenylalanine tert-Butyl Ester**

N-(4-(N,N-Di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-L-tyrosine tert-butyl ester (180 mg, 0.34 mmol) was dissolved in pyridine (5 ml). Dimethylcarbamoyl chloride (108 mg, 1 mmol) was added dropwise and the mixture stirred at room temperature overnight. Pyridine was removed under high vacuum (low water bath temperature), the residue absorbed onto silica gel and purified by flash column chromatography (silica, hexane:EtOAc 2:1) to yield the title compound (140 mg, 68 %).

5           The compound from Step A (140 mg, 0.23 mmol) was dissolved in formic acid and the mixture stirred at room temperature overnight. Excess formic acid was removed under reduced pressure to yield the title compound as a white solid (110 mg, 87 %).

Physical data were as follows:

10                    Anal. Calc'd for  $C_{26}H_{41}N_5O_6$  S: C, 56.6; H, 7.49; N, 12.69.  
                      Found: C, 56.67; H, 7.4; N, 12.46.

### Example 133

**15** **Synthesis of**  
**N-(4-(N,N-Di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-L-4-(4-**  
**methylpiperazin-1-ylcarbonyloxy)phenylalanine**

20      **Step A:      Preparation of N-(4-(N,N-Di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine tert-Butyl Ester**

A solution of N-(4-(N,N-di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-L-tyrosine tert-butyl ester (500 mg, 0.93 mmol), and p-nitrophenyl chloroformate (179 mg, 0.89 mmol) in dichloromethane (20 mL) was cooled to 0°C under an argon atmosphere. Triethylamine (235 mg, 2.32 mmol) was added dropwise and the mixture stirred at 0°C for 30 mins, then allowed to warm to room temperature for a further 40 mins. The mixture was recooled to 0°C and N-methylpiperazine (90 mg, 0.89 mmol) added. The mixture was allowed to warm to room temperature and stirred for three hours.

The mixture was diluted with diethyl ether (150 mL) and the organic solution washed with 10% potassium carbonate solution until no further yellow color was produced in the aqueous phase. The organic layer was separated, dried ( $\text{MgSO}_4$ ) and the solvent removed under reduced pressure. The residue was

-- 238 --

purified by flash column chromatography (silica, EtOAc:MeOH:Et<sub>3</sub>N 94:5:1) to give the title compound as a pale yellow foam (310 mg, 50 %).

Physical data were as follows:

Anal. Calc'd for C<sub>33</sub>H<sub>54</sub>N<sub>6</sub>O<sub>6</sub> S: C, 59.79; H, 8.21; N, 12.68.

5 Found: C, 59.47; H, 8.25; N, 12.49

Step B: **Preparation of N-(4-(N,N-Di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine**

10

The compound from Step A (200 mg, 0.3 mmol) was dissolved in formic acid (5 mL) and the mixture stirred at room temperature for 48 hrs. Excess formic acid was removed under reduced pressure and the residue recrystallized from EtOAc/MeOH to yield the title compound as an off-white solid (120 mg, 67 %).

15

Physical data were as follows:

Anal. Calc'd for C<sub>29</sub>H<sub>46</sub>N<sub>6</sub>O<sub>6</sub> S.0.75H<sub>2</sub>O: C, 56.15; H, 7.72; N, 13.55. Found: C, 56.1; H, 7.44; N, 13.46.

20

#### Example 134

**Synthesis of N-[4-(2-(3-Methylphenylaminocarbonylamino)eth-1-ylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl]-L-4-(N,N-dimethylcarbamyloxy)phenylalanine**

25

Step A: **Preparation of N-(4-Ethoxy-1,1-dioxo-1,2,5-thiadiazol-3-yl)-L-tyrosine tert-Butyl Ester**

A solution of 3,4-diethyloxy-1,1-dioxo-1,2,5-thiadiazole (400 mg, 1.94 mmol) and L-tyrosine t-butyl ester (1.25 g, 5.2 mmol) in ethanol (25 mL) was stirred at room temperature overnight. Solvent was removed under reduced pressure and the product used in further transformations without further purification (Yield 790 mg).

30

-- 239 --

**Step B: Preparation of 2-(3-Methylphenylaminocarbonylamino)eth-1-ylamine**

5 N-Boc-Ethylene diamine (800 mg, 5 mmol) and m-tolyl isocyanate (665 mg, 5 mmol) were dissolved in acetonitrile and the mixture stirred at room temperature for 4 hrs. Solvent was removed under reduced pressure and the residue absorbed onto silica gel; prior to purification by flash column chromatography (silica, hexane:EtOAc 1:1) to yield the desired compound as a white solid (300 mg, 21 %) (MS ( +ESI, m/e) 294 (M+H)<sup>+</sup>). The N-Boc  
10 protected compound (300 mg, 1.02 mmol) was dissolved in formic acid (10 ml) and the mixture stirred at room temperature overnight. Excess acid was removed to yield the formate salt of the title compound as a white foam (210 mg).

**15 Step C: Preparation of N-[4-(2-(3-Methylphenylamino-carbonylamino)eth-1-ylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl]-L-tyrosine tert-Butyl Ester**

To a solution of N-(4-ethoxy-1,1-dioxo-1,2,5-thiadiazol-3-yl)-L-tyrosine tert-butyl ester from Step A (150 mg, 0.38 mmol) and the formate  
20 salt of 2-(3-methylphenylaminocarbonylamino)eth-1-ylamine from Step B (210 mg, 0.89 mmol) in ethanol (10 mL) was added triethylamine (133 mg, 1.44 mmol). The reaction was stirred at room temperature overnight. Solvent was removed under reduced pressure and the residue purified by  
25 flash column chromatography (silica, 5 % MeOH in EtOAc) to give the title compound (130 mg, 91 %).

Physical data were as follows:

MS ( +ESI, m/e) 545 (M+H)<sup>+</sup>.

-- 240 --

Step D: **Preparation of N-[4-(2-(3-Methylphenylamino-carbonylamino)eth-1-ylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine tert-Butyl Ester**

5

The intermediate from Step C (130 mg, 0.24 mmol) was dissolved in pyridine (5 mL). Dimethylcarbamoyl chloride (77 mg, 0.72 mmol) was added dropwise and the mixture heated at 50°C under an argon atmosphere overnight. Pyridine was removed under reduced pressure, the residue  
10 absorbed onto silica gel and purified by flash column chromatography (silica, hexane:EtOAc 1:2, then 5% MeOH in EtOAc) to yield the title compound (140 mg, 93 %).

Physical data were as follows:

MS ( +ESI, m/e) 616 (M+H)<sup>+</sup>.

15

Step E: **Preparation of N-[4-(2-(3-Methylphenylamino-carbonylamino)eth-1-ylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine**

20

The compound from Step D (120 mg, 0.19 mmol) was dissolved in formic acid (10 mL) and the mixture stirred at room temperature for 36 hrs. Excess acid was removed to yield the title compound as a pale yellow foam (100 mg, 93 %).

Physical data were as follows:

25

MS ( +ESI, m/e) 560 (M+H)<sup>+</sup>.

30

-- 241 --

## Example 135

## Synthesis of

5        **N-(4-(N,N-Dimethylamino)-1-oxo-1,2,5-thiadiazol-3-yl)-L-4-(N,N-dimethylcarbamoyloxy)phenylalanine *tert*-Butyl Ester**

Step A:        **Preparation of N-(4-Ethoxy-1-oxo-1,2,5-thiadiazol-3-yl)-L-tyrosine *tert*-Butyl Ester**

10        A solution of 3,4-diethoxy-1-oxo-1,2,5-thiadiazole (1 g, 0.52 mmol) and L-tyrosine *t*-butyl ester (1.25 g, 0.52 mmol) in ethanol (25 mL) was stirred at room temperature for 60 hr. Solvent was removed under reduced pressure and the residue purified by flash column chromatography (silica, hexane:EtOAc 1:1 to give the title intermediate (1.75 g, 88 %).

15        Physical data were as follows:  
MS ( +ESI, m/e) 382 (M+H)<sup>+</sup>.

20        Step B:        **Preparation of N-(4-Ethoxy-1-oxo-1,2,5-thiadiazol-3-yl)-L-4-(N,N-dimethylcarbamoyloxy)phenylalanine *tert*-Butyl Ester**

The intermediate from Step A (400 mg, 1.05 mmol) was dissolved in pyridine (10 mL) and dimethylcarbamoyl chloride (338 mg, 3.15 mmol) was added. The reaction was stirred at room temperature under an inert atmosphere overnight. TLC indicated large amounts of unreacted starting material so the mixture was heated at 50°C for a further 48 hrs. Excess pyridine was removed under reduced pressure and the residue purified by flash column chromatography (silica, hexane:EtOAc 1:1 to give the title intermediate (280 mg, 59 %).

25        Physical data were as follows:  
30        MS ( +ESI, m/e) 453 (M+H).

**Step C: Preparation of N-(4-(N,N-Dimethylamino)-1-oxo-1,2,5-thiadiazol-3-yl)-L-4-(N,N-dimethylcarbamyloxy)phenylalanine *tert*-Butyl Ester**

5 A 2M solution of dimethylamine in THF (5 mL, 10 mmol) was added to a solution of the compound from Step B (180 mg, 0.35 mmol) in ethanol (10 mL). The reaction was stirred at room temperature overnight and solvent removed under reduced pressure. Residue was purified by flash column chromatography (silica, EtOAc:MeOH:Et<sub>3</sub>N 90:10:1) to give the title  
10 compound as a white foam (140 mg, 88 %).

Physical data were as follows:

Anal. Calc'd for  $C_{22}H_{29}N_5O_5$  S: C, 53.2; H, 6.47; N, 15.51.

Found: C, 52.94; H, 6.18; N, 15.34.

15 **Example 136**

### Synthesis of *N*-(5-(2,2,2-Trifluoroethyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine

20 Substituting L-4-(2,6-dimethoxyphenyl)phenylalanine methyl ester from Method TTT for L-4-(N,N-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester and following the procedure described for the preparation of Example 31 yielded the title compound.

25 Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 8.41 (s, 1H), 8.05 (s, 1H), 7.24 (t, 1H), 7.2 (d, 2H), 7.1 (d, 2H), 6.67 (d, 2H), 5.1 (dd, 1H), 3.65 (s, 6H), 3.61-3.42 (m, 2H), 3.36 (dd, 1H), 3.2 (dd, 1H).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ 175.8, 162.3, 159.2, 157.9, 155.8, 136.9, 134.4, 132.2, 130.0, 129.5, 127.4, 120.9, 109.6, 105.7, 56.8, 56.2, 37.9, 32.6.



***N*-(2-(*N*-Cyclohexyl-*N*-methyl)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine**

Substituting L-4-(2,6-dimethoxyphenyl)phenylalanine methyl ester from Method TTT for L-4-(N,N-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester and following the procedure described for the preparation of Example 14 yielded the title compound.

**Physical data were as follows:**

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 7.37-7.19 (m, 5.5H), 7.09-7.02 (m, 4H), 6.94 (d, 0.5H), 6.68 (d, 2H), 4.79-4.74 (m, 0.5H), 4.69-4.65 (m, 0.5H), 3.67 (s, 3H), 3.65 (s, 3H), 3.44-3.33 (m, 1H), 3.02-2.95 (m, 4H), 2.19 (s, 1.5H), 1.85-1.71 (m, 6.5H), 1.57 (m, 4H), 1.29-1.2 (br s, 1H).

### Example 138

## Synthesis of

***N*-(5-(2-Fluorophenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine**

Substituting L-4-(2,6-dimethoxyphenyl)phenylalanine methyl ester from Method TTT for L-4-(N,N-dimethylcarbamoyloxy)phenylalanine *tert*-butyl ester and following the procedure described for the preparation of Example 70 yielded the title compound.

**Physical data were as follows:**

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 8.50 (s, 1H), 8.01 (s, 1H), 7.3-7.0 (m, 9H), 6.69 (d, 2H), 5.0 (m, 1H), 3.65 (s, 6H), 3.20-3.05 (m, 2H).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ 153.2, 151.6, 147.1, 130.2, 128.6, 126.7, 126.6, 126.5, 126.4, 126.3, 123.9, 123.5, 123.2, 120.5, 120.4, 111.7, 111.4, 99.6, 59.3, 31.7.

-- 244 --

## Example 139

## Synthesis of

*N*-(2-(*N*-Methyl-*N*-propyl)-5-(2-tolyl)pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine

Substituting *L*-4-(2,6-dimethoxyphenyl)phenylalanine methyl ester from Method TTT for *L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester and following the procedure described for the preparation of Example 61 yielded the title compound.

Physical data were as follows:

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 10.30-8.80 (br, 1H), 7.68 (s, 0.5H), 7.63 (s, 0.5H), 7.40-6.60 (m, 1H), 6.15 (m, 1H), 4.70 (m, 1H), 3.68 (s, 3H), 3.66 (s, 3H), 3.80-3.00 (m, 4H), 3.07 (s, 3H), 2.12 (s, 1.5H), 2.08 (s, 1.5H), 1.61 (bs, 2H), 0.87 (bs, 3H).

## Example 140

## Synthesis of

*N*-(3-Chloropyrazin-2-yl)-*L*-4-[1-(*tert*-butoxycarbonyl)piperidin-4-ylcarbonylamino]phenylalanine Ethyl Ester

Step A: Preparation of *N*-(3-Chloropyrazin-2-yl)-*L*-4-nitrophenylalanine

4-Nitrophenylalanine (50 mm, 10.59 mg) were stirred in absolute ethanol containing 1.0 eq (1.26 g) of sodium metal. The reaction mixture was stripped to a brown solid and the sodium salt was taken up in 200 mL of butanol containing 1.0 eq (7.45 g) 2,3-dichloropyrazine. The reaction mixture was refluxed overnight and the solvent was then removed under reduced pressure. The residue was taken up in ethyl acetate and washed with water (1 x), brine (1 x), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and stripped to give 15.5 g of the title intermediate as a brown oil.

Physical data were as follows:

Analytical: MS: (+)FAB [M+H] @ M/Z 323 with 1 Cl.

-- 245 --

**Step B: Preparation of N-(3-Chloropyrazin-2-yl)-L-4-nitrophenylalanine Ethyl Ester**

The intermediate from Step A was suspended in 300 mL of absolute ethanol. The reaction flask was placed in an ice bath and cooled to 0°C and HCl (g) was bubbled into reaction for 15 minutes. The gas tube was replaced with a drying tube and the reaction mixture was warmed to room temperature and stirred overnight. Ethanol was stripped off under reduced pressure to afford a dark brown residue which was taken up in ethyl acetate and washed with sat. NaHCO<sub>3</sub> (2x), H<sub>2</sub>O (1x), brine (1x), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and stripped to afford 15 g of a dark brown oil. This oil (8.0 g) was chromatographed on a silica 60 column packed in methylene chloride to provide 1.5 g (20% yield) of the title intermediate.

Physical data were as follows:

Analytical: MS: EI M<sup>+</sup> @ M/Z 350 1 Cl present.

**Step C: Preparation of N-(3-Chloropyrazin-2-yl)-L-4-aminophenylalanine Ethyl Ester**

The intermediate from Step B (0.75 g, 0.021 mol) was placed in a Paar hydrogenation bottle with 50 mL ethanol and 0.40 g of Pd/C catalyst. The bottle was placed on Paar shaker under 50 psi of H<sub>2</sub> for 3 hrs. The reaction mixture was then filtered through a sintered glass funnel (F) and the filtered catalyst was washed with ethanol. The combined filtrates were stripped to a yellow oil and the oil was taken up in ethyl acetate. A yellow precipitate formed and was filtered off. The filtrate was washed with NaHCO<sub>3</sub> solution (1 x), H<sub>2</sub>O (1x), brine (1x), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and stripped to afford the title intermediate as a yellow oil (0.340 g, 55% yield).

**Step D: Preparation of N-(3-Chloropyrazin-2-yl)-L-4-[1-(tert-butoxycarbonyl)piperidin-4-ylcarbonylamino]phenylalanine Ethyl Ester**

N-Boc-piperidine 4-carboxylic acid (0.253 g, 1.0 eq., 0.0011 mol) was stirred in 30 mL methylene chloride and reaction mixture was cooled to 0°C in ice bath. HOBt (0.224 g, 1.5 eq) was added and the mixture was stirred for 10 minutes then the intermediate from Step C (1 eq., 0.32 g) was added. The reaction mixture was stirred for 5 minutes and then 1,3-dicyclohexylcarbodiimide (0.25 g, 1.1 eq) was added. The reaction mixture was warmed to room temperature and stirred overnight. The reaction was then filtered and the filtrate was stripped to give a yellow solid. The solid was taken up in ethyl acetate and filtered. The ethyl acetate solution was washed with 10% citric acid (1 x), H<sub>2</sub>O (1 x), brine (1 x), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and stripped to afford a yellow oil (0.630 g; MS: EI M<sup>+</sup> @ M/Z 531 (1 chloro)). The yellow oil was chromatographed on a silica 60 column eluting with 3:1 hexane/ethyl acetate to afford 0.097 g of the title compound. This compound may also be used as an intermediate for other compounds of this invention.

Physical data were as follows:

Analytical: CHN: Theory (0.5 H<sub>2</sub>O): C, 57.71; H, 6.72; N, 12.94  
Found: C, 57.79; H, 6.32; N, 12.78. MS: M<sup>+</sup> @ M/Z 531 (1 Chloro).

20

#### Example A

#### *In vitro* Assay For Determining Binding of Candidate Compounds to VLA-4

An *in vitro* assay was used to assess binding of candidate compounds to  $\alpha_4\beta_1$  integrin. Compounds which bind in this assay can be used to assess VCAM-1 levels in biological samples by conventional assays (e.g., competitive assays). This assay is sensitive to IC<sub>50</sub> values as low as about 1nM.

30

The activity of  $\alpha_4\beta_1$  integrin was measured by the interaction of soluble VCAM-1 with Jurkat cells (e.g., American Type Culture Collection

Nos. TIB 152, TIB 153, and CRL 8163), a human T-cell line which expresses high levels of  $\alpha_4\beta_1$  integrin. VCAM-1 interacts with the cell surface in an  $\alpha_4\beta_1$  integrin-dependent fashion (Yednock, et al. J. Biol. Chem., 1995, 270:28740).

5

Recombinant soluble VCAM-1 was expressed as a chimeric fusion protein containing the seven extracellular domains of VCAM-1 on the N-terminus and the human IgG<sub>1</sub> heavy chain constant region on the C-terminus. The VCAM-1 fusion protein was made and purified by the manner described  
10 by Yednock, *supra*.

15

Jurkat cells were grown in RPMI 1640 supplemented with 10% fetal bovine serum, penicillin, streptomycin and glutamine as described by Yednock, *supra*.

Jurkat cells were incubated with 1.5 mM MnCl<sub>2</sub> and 5  $\mu$ g/mL 15/7 antibody for 30 minutes on ice. Mn<sup>+2</sup> activates the receptor to enhance ligand binding, and 15/7 is a monoclonal antibody that recognizes an activated/ligand occupied conformation of  $\alpha_4\beta_1$  integrin and locks the  
20 molecule into this conformation thereby stabilizing the VCAM-1/ $\alpha_4\beta_1$  integrin interaction. Yednock, et al., *supra*. Antibodies similar to the 15/7 antibody have been prepared by other investigators (Luque, et al, 1996, J. Biol. Chem. 271:11067) and may be used in this assay.

25

Cells were then incubated for 30 minutes at room temperature with candidate compounds, in various concentrations ranging from 66  $\mu$ M to 0.01  $\mu$ M using a standard 5-point serial dilution. 15  $\mu$ L soluble recombinant VCAM-1 fusion protein was then added to Jurkat cells and incubated for 30 minutes on ice. (Yednock et al., *supra*).

30

-- 248 --

Cells were then washed two times and resuspended in PE-conjugated goat F(ab')<sub>2</sub> anti-mouse IgG Fc (Immunotech, Westbrook, ME) at 1:200 and incubated on ice, in the dark, for 30 minutes. Cells were washed twice and analyzed with a standard fluorescence activated cell sorter ("FACS") analysis as described in Yednock, et al., *supra*.

Compounds having an IC<sub>50</sub> of less than about 15 μM possess binding affinity to α<sub>4</sub>β<sub>1</sub>.

When tested in this assay, each of the compound prepared in the above examples has or is expected to have an IC<sub>50</sub> of 15 μM or less (or is expected to be active *in vivo*).

#### Example B

##### ***In vitro* Saturation Assay For Determining Binding of Candidate Compounds to α<sub>4</sub>β<sub>1</sub>**

The following describes an *in vitro* assay to determine the plasma levels needed for a compound to be active in the Experimental Autoimmune Encephalomyelitis ("EAE") model, described in the next example, or in other *in vivo* models.

Log-growth Jurkat cells are washed and resuspended in normal animal plasma containing 20 μg/ml of the 15/7 antibody (described in the above example).

The Jurkat cells are diluted two-fold into either normal plasma samples containing known candidate compound amounts in various concentrations ranging from 66 μM to 0.01 μM, using a standard 12 point serial dilution for a standard curve, or into plasma samples obtained from the peripheral blood of candidate compound-treated animals.

-- 249 --

Cells are then incubated for 30 minutes at room temperature, washed twice with phosphate-buffered saline ("PBS") containing 2% fetal bovine serum and 1mM each of calcium chloride and magnesium chloride (assay medium) to remove unbound 15/7 antibody.

5

The cells are then exposed to phycoerythrin-conjugated goat F(ab')<sub>2</sub> anti-mouse IgG Fc (Immunotech, Westbrook, ME), which has been adsorbed for any non-specific cross-reactivity by co-incubation with 5% serum from the animal species being studied, at 1:200 and incubated in the dark at 4°C for 30 minutes.

10

Cells are washed twice with assay medium and resuspended in the same. They are then analyzed with a standard fluorescence activated cell sorter ("FACS") analysis as described in Yednock et al. J. Biol. Chem., 1995, 270:28740.

15

The data is then graphed as fluorescence versus dose, e.g., in a normal dose-response fashion. The dose levels that result in the upper plateau of the curve represent the levels needed to obtain efficacy in an *in vivo* model.

20

This assay may also be used to determine the plasma levels needed to saturate the binding sites of other integrins, such as the  $\alpha_9\beta_1$  integrin, which is the integrin most closely related  $\alpha_4\beta_1$  (Palmer et al, 1993, J. Cell Bio., 123:1289). Such binding is predictive of *in vivo* utility for inflammatory conditions mediated by  $\alpha_9\beta_1$  integrin, including by way of example, airway hyper-responsiveness and occlusion that occurs with chronic asthma, smooth muscle cell proliferation in atherosclerosis, vascular occlusion following angioplasty, fibrosis and glomerular scarring as a result of renal disease, aortic stenosis, hypertrophy of synovial membranes in rheumatoid arthritis,

25

30

and inflammation and scarring that occur with the progression of ulcerative colitis and Crohn's disease.

Accordingly, the above-described assay may be performed with a  
5 human colon carcinoma cell line, SW 480 (ATTC #CCL228) transfected with  
cDNA encoding  $\alpha_9$  integrin (Yokosaki et al., 1994, J. Biol. Chem.,  
269:26691), in place of the Jurkat cells, to measure the binding of the  $\alpha_9\beta_1$   
integrin. As a control, SW 480 cells which express other  $\alpha$  and  $\beta_1$  subunits  
may be used.

10

Accordingly, another aspect of this invention is directed to a method  
for treating a disease in a mammalian patient, which disease is mediated by  
 $\alpha_9\beta_1$ , and which method comprises administering to said patient a  
therapeutically effective amount of a compound of this invention. Such  
15 compounds are preferably administered in a pharmaceutical composition  
described herein above. Effective daily dosing will depend upon the age,  
weight, condition of the patient which factors can be readily ascertained by  
the attending clinician. However, in a preferred embodiment, the compounds  
are administered from about 20 to 500  $\mu\text{g/kg}$  per day.

20

### Example C

#### *In vivo* Evaluation

The standard multiple sclerosis model, Experimental Autoimmune (or  
25 Allergic) Encephalomyelitis ("EAE"), was used to determine the effect of  
candidate compounds to reduce motor impairment in rats or guinea pigs.  
Reduction in motor impairment is based on blocking adhesion between  
leukocytes and the endothelium and correlates with anti-inflammatory activity  
in the candidate compound. This model has been previously described by  
30 Keszthelyi et al., Neurology, 1996, 47:1053-1059, and measures the delay of  
onset of disease.



-- 251 --

Brains and spinal cords of adult Hartley guinea pigs were homogenized in an equal volume of phosphate-buffered saline. An equal volume of Freund's complete adjuvant (100 mg *mycobacterium tuberculosis* plus 10 ml Freund's incomplete adjuvant) was added to the homogenate. The mixture was emulsified by circulating it repeatedly through a 20 ml syringe with a peristaltic pump for about 20 minutes.

Female Lewis rats (2-3 months old, 170-220 g) or Hartley guinea pigs (20 day old, 180-200 g) were anesthetized with isoflurane and three injections of the emulsion, 0.1 ml each, were made in each flank. Motor impairment onset is seen in approximately 9 days.

Candidate compound treatment began on Day 8, just before onset of symptoms. Compounds were administered subcutaneously ("SC"), orally ("PO") or intraperitoneally ("IP"). Doses were given in a range of 10mg/kg to 200 mg/kg, bid, for five days, with typical dosing of 10 to 100 mg/kg SC, 10 to 50 mg/kg PO, and 10 to 100 mg/kg IP.

Antibody GG5/3 against  $\alpha_4\beta_1$  integrin (Keszthelyi et al., Neurology, 1996, 47:1053-1059), which delays the onset of symptoms, was used as a positive control and was injected subcutaneously at 3 mg/kg on Day 8 and 11.

Body weight and motor impairment were measured daily. Motor impairment was rated with the following clinical score:

0	no change
1	tail weakness or paralysis
2	hindlimb weakness
3	hindlimb paralysis
4	moribund or dead

-- 252 --

A candidate compound was considered active if it delayed the onset of symptoms, e.g., produced clinical scores no greater than 2 or slowed body weight loss as compared to the control.

5

#### Example D Asthma Model

Inflammatory conditions mediated by  $\alpha_4\beta_1$  integrin include, for example, airway hyper-responsiveness and occlusion that occurs with chronic  
10 asthma. The following describes an asthma model which can be used to study the *in vivo* effects of the compounds of this invention for use in treating asthma.

Following the procedures described by Abraham et al, J. Clin. Invest,  
15 93:776-787 (1994) and Abraham et al, Am J. Respir Crit Care Med, 156:696-703 (1997), both of which are incorporated by reference in their entirety. Compounds of this invention are formulated into an aerosol and administered to sheep which are hypersensitive to *Ascaris suum* antigen. Compounds which decrease the early antigen-induced bronchial response  
20 and/or block the late-phase airway response, e.g., have a protective effect against antigen-induced late responses and airway hyper-responsiveness ("AHR"), are considered to be active in this model.

Allergic sheep which are shown to develop both early and late  
25 bronchial responses to inhaled *Ascaris suum* antigen are used to study the airway effects of the candidate compounds. Following topical anesthesia of the nasal passages with 2% lidocaine, a balloon catheter is advanced through one nostril into the lower esophagus. The animals are then intubated with a cuffed endotracheal tube through the other nostril with a flexible fiberoptic  
30 bronchoscope as a guide.

-- 253 --

Pleural pressure is estimated according to Abraham (1994). Aerosols (see formulation below) are generated using a disposable medical nebulizer that provides an aerosol with a mass median aerodynamic diameter of  $3.2 \mu\text{m}$  as determined with an Andersen cascade impactor. The nebulizer is  
5 connected to a dosimeter system consisting of a solenoid valve and a source of compressed air (20 psi). The output of the nebulizer is directed into a plastic T-piece, one end of which is connected to the inspiratory port of a piston respirator. The solenoid valve is activated for 1 second at the beginning of the inspiratory cycle of the respirator. Aerosols are delivered at  
10  $V_T$  of 500 ml and a rate of 20 breaths/minute. A 0.5% sodium bicarbonate solution only is used as a control.

To assess bronchial responsiveness, cumulative concentration-response curves to carbachol can be generated according to Abraham (1994).  
15 Bronchial biopsies can be taken prior to and following the initiation of treatment and 24 hours after antigen challenge. Bronchial biopsies can be preformed according to Abraham (1994).

An *in vitro* adhesion study of alveolar macrophages can also be  
20 performed according to Abraham (1994), and a percentage of adherent cells is calculated.

#### Aerosol Formulation

A solution of the candidate compound in 0.5% sodium  
25 bicarbonate/saline (w/v) at a concentration of 30.0 mg/mL is prepared using the following procedure:

#### A. Preparation of 0.5% Sodium Bicarbonate / Saline Stock Solution: 100.0 mL

30

-- 254 --

Ingredient	Gram / 100.0 mL	Final Concentration
Sodium Bicarbonate	0.5 g	0.5%
Saline	q.s. ad 100.0 mL	q.s. ad 100%

## 5      Procedure:

1. Add 0.5g sodium bicarbonate into a 100 mL volumetric flask.
2. Add approximately 90.0 mL saline and sonicate until dissolved.
3. Q.S. to 100.0 mL with saline and mix thoroughly.

10

B. Preparation of 30.0 mg/mL Candidate Compound: 10.0 mL

Ingredient	Gram / 10.0 mL	Final Concentration
Candidate Compound	0.300 g	30.0 mg/mL
0.5% Sodium Bicarbonate / Saline Stock Solution	q.s. ad 10.0 mL	q.s ad 100%

15

## Procedure:

- 20      1. Add 0.300 g of the candidate compound into a 10.0 mL volumetric flask.
2. Add approximately 9.7 mL of 0.5% sodium bicarbonate / saline stock solution.
3. Sonicate until the candidate compound is completely dissolved.
- 25      4. Q.S. to 10.0 mL with 0.5% sodium bicarbonate / saline stock solution and mix thoroughly.

Using a conventional oral formulation, compounds of this invention would be active in this model.

30

-- 255 --

Example E  
**Allograft Model**

5        Allograft rejection, associated with infiltration of inflammatory cells,  
is the leading obstacle to long-term allograft survival. Cell surface adhesion  
molecules facilitate alloantigen recognition *in vitro* and may be critical for  
lymphocyte traffic *in vivo*. The following describes a model which can be  
used to study the *in vivo* effects of the compounds of this invention in the  
control of allograft rejection.

10

The following procedures are described in Coito et al.,  
*Transplantation* (1998) 65(6):699-706 and in Korom et al., *Transplantation*  
(1998) 65(6):854-859, both of which are incorporated by reference in their  
entirety.

15

Following the procedures described in Coito and Korom, male adult  
rats weighing approximately 200 - 250 g are used in this model. Lewis rats  
are used as the recipients of cardiac allografts from Lewis X Brown Norway  
rats. Hearts are transplanted into the abdominal great vessels using standard  
20        microvascular techniques.


A candidate compound is administered to the transplant recipient in a  
suitable pharmaceutical carrier for a 7-day course of treatment starting the  
day of the engraftment. Doses range from 0.3 to 30 mg/kg/day. Control  
25        recipients receive the pharmaceutical carrier only. The rats are euthanized  
and their cardiac allografts are analyzed as described in Coito and Korom.

Using conventional formulations, compounds of this invention would  
be active in this model.

30

## WHAT IS CLAIMED IS:

1. A method for treating a disease mediated by VLA-4 in a patient, which method comprises administering a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of formula Ia and/or Ib:



- 15 wherein, in formula Ia, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form an aryl, cycloalkenyl, heteroaryl or heterocyclic group having at least five atoms in the aryl, cycloalkenyl, heteroaryl or heterocyclic group and optionally containing or additionally containing in the case of heteroaryl and
- 20 heterocyclic groups 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heteroaryl or heterocyclic group is mono-cyclic;

- in formula Ib, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W' to which they are bound respectively, are joined to form a cycloalkyl, cycloalkenyl or heterocyclic group having at least five atoms in the cycloalkyl, cycloalkenyl or heterocyclic group and optionally containing or additionally containing in the case of the heterocyclic group 1 to 3
- 25 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heterocyclic group is mono-cyclic;

-- 257 --

and further wherein said aryl, cycloalkyl, cycloalkenyl, heteroaryl or heterocyclic group of formula Ia or Ib is optionally substituted, on any ring atom capable of substitution, with 1-3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen, hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where each R is independently hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, -N[S(O)<sub>2</sub>-R']<sub>2</sub> and -N[S(O)<sub>2</sub>-NR']<sub>2</sub> where each R' is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

$R^3$  and  $R^{3'}$  are independently selected from the group consisting of hydrogen, isopropyl,  $-\text{CH}_2\text{Z}$  where Z is selected from the group consisting of hydrogen, hydroxyl, acylamino, alkyl, alkoxy, aryloxy, aryl, aryloxyaryl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, carboxyl-substituted heterocyclic, cycloalkyl, substituted alkyl, substituted alkoxy, substituted aryl, substituted aryloxy, substituted aryloxyaryl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic, and

where  $R^3$  and  $R^{3'}$  are joined to form a substituent selected from the group consisting of  $=\text{CHZ}$  where Z is defined above provided that Z is not hydroxyl or thiol, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic and substituted heterocyclic;

Q is selected from the group consisting of  $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{S}(\text{O})-$ ,  $-\text{S}(\text{O})_2$ , and  $-\text{NR}^4-$ ;

$R^4$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic or, optionally,  $R^4$  and  $R^1$  or  $R^4$  and  $R^2$ , together with the atoms to which they are bound, are joined to form a heteroaryl, a substituted heteroaryl, a heterocyclic or a substituted heterocyclic group;

W is selected from the group consisting of nitrogen and carbon; and

W' is selected from the group consisting of nitrogen, carbon, oxygen, sulfur,  $\text{S}(\text{O})$ , and  $\text{S}(\text{O})_2$ ;

X is selected from the group consisting of hydroxyl, alkoxy, substituted alkoxy, alkenoxy, substituted alkenoxy, cycloalkoxy, substituted cycloalkoxy, cycloalkenoxo, substituted cycloalkenoxo, aryloxy, substituted aryloxy, heteroaryloxy, substituted heteroaryloxy, heterocycloxy,



-- 259 --


substituted heterocycloxy and -NR<sup>''</sup>R<sup>''</sup> where each R<sup>''</sup> is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;

and further wherein the compound of formula Ia and/or Ib has a binding affinity to VLA-4 as expressed by an IC<sub>50</sub> of about 15μM or less.

2. The method of Claim 1, wherein R<sup>3</sup> is -(CH<sub>2</sub>)<sub>x</sub>-Ar-R<sup>9</sup>, where Ar is aryl, substituted aryl, heteroaryl and substituted heteroaryl; R<sup>9</sup> is selected from the group consisting acyl, acylamino, acyloxy, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, oxycarbonylamino, oxythiocarbonylamino, thioamidino, thiocarbonylamino, aminosulfonylamino, aminosulfonyloxy, aminosulfonyl, oxysulfonylamino and oxysulfonyl; and x is an integer from 0 to 4; and R<sup>3'</sup> is hydrogen.

3. The method of Claim 2, wherein R<sup>3</sup> is a group of the formula:



wherein R<sup>9</sup> and x are as defined in Claim 2.

4. The method of Claim 3, wherein R<sup>9</sup> is in the *para* position of the phenyl ring and x is an integer from 1 to 4.

-- 260 --



5. The method of Claim 4, wherein  $R^9$  is selected from the group consisting of  $-O-Z-NR^{11}R^{11'}$  and  $-O-Z-R^{12}$  wherein  $R^{11}$  and  $R^{11'}$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, and where  $R^{11}$  and  $R^{11'}$  are joined to form a heterocycle or a substituted heterocycle,  $R^{12}$  is selected from the group consisting of heterocycle and substituted heterocycle, and Z is selected from the group consisting of  $-C(O)-$  and  $-SO_2-$ .

6. The method of Claim 5, wherein Z is  $-C(O)-$ .

7. The method of Claim 6, wherein  $R^9$  is  $-OC(O)NR^{11}R^{11'}$ .

8. The method of Claim 1, wherein Q is  $-NR^4-$ .

9. The method of Claim 1, wherein the compound has formula IIa or IIb:



30

wherein

ring A and ring B independently form a heteroaryl or substituted heteroaryl group having two nitrogen atoms in the heteroaryl ring;

5  $R^5$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

10  $R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $-SO_2R^{10}$  where  $R^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl;

15 or optionally, one of,  $R^4$  and ring A,  $R^4$  and  $R^5$ ,  $R^4$  and  $R^6$ , or  $R^5$  and  $R^6$ , together with the atoms to which they are bound, can be joined to form a heterocyclic or substituted heterocyclic ring;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof; and provided that ring B does not form a 6-amino or substituted amino pyrimidin-4-yl group.

20 10. The method of Claim 9, wherein ring A forms a pyridazine, pyrimidine or pyrazine ring, wherein the pyridazine, pyrimidine or pyrazine ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen, and


30 ring B forms a pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or a 1,1-dioxo-1,2,5-thiadiazole ring, wherein the pyridazine,

-- 262 --

pyrimidine or pyrazine ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.

11. The method of Claim 1, wherein the compound has formula IIIa, IIIc, IIIe or IIIf:


10



IIIa


15

20



IIIc


25



IIIe


30

5



IIIe

10



IIIf

15

wherein

20  $R^{4'}$  is selected from the group consisting of hydrogen and alkyl or, optionally, one of,  $R^{4'}$  and  $R^5$ ,  $R^{4'}$  and  $R^6$ ,  $R^5$  and  $R^6$ ,  $R^5$  and  $R^8$ , or  $R^6$  and  $R^8$ , together with the atoms to which they are bound, are joined to form a heterocyclic, a substituted heterocyclic, a heteroaryl or substituted heteroaryl group optionally containing from 1 to 3 additional hetero ring atoms selected from the group consisting of oxygen, nitrogen and sulfur;

25  $R^{4''}$  is selected from the group consisting of hydrogen and alkyl;

$R^5$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

$R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $-SO_2R^{10}$  where  $R^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl;

$R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$R^{16}$  and  $R^{17}$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen; and

$R^{18}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

$R^{20}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;


$R^{21}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;

$b$  is 1 or 2;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

12. The method of Claim 11, wherein the compound is selected from formula IIIId, IIIe or IIIf.

13. A method for treating a disease mediated by VLA-4 in a patient, which method comprises administering a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of formula IVa and/or IVb:



wherein, in formula IVa, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form an aryl, cycloalkenyl, heteroaryl or heterocyclic group having at least five atoms in the aryl, cycloalkenyl, heteroaryl or heterocyclic group and optionally containing or additionally containing in the case of heteroaryl and heterocyclic groups 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heteroaryl or heterocyclic group is mono-cyclic;

in formula IVb, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W' to which they are bound respectively, are joined to form a cycloalkyl, cycloalkenyl or heterocyclic group having at least five atoms in the cycloalkyl, cycloalkenyl or heterocyclic group and optionally containing or additionally containing in the case the heterocyclic group 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heterocyclic group is mono-cyclic;

and further wherein said aryl, cycloalkyl, cycloalkenyl, heteroaryl or heterocyclic group of formula IVa or IVb is optionally substituted, on any ring atom capable of substitution, with 1-3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocabonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen, hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{OSO}_2\text{-NRR}$  where each R is independently hydrogen or alkyl,  $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heterocyclic}$  where R is hydrogen or alkyl,  $-\text{N}[\text{S}(\text{O})_2\text{-R'}]_2$  and  $-\text{N}[\text{S}(\text{O})_2\text{-NR'}]_2$  where each R' is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;



$R^{13}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl, Cy, and Cy- $C_{1-10}$  alkyl, wherein alkyl is optionally substituted with one to four substituents independently selected from  $R^a$ ; and Cy is optionally substituted with one to four substituents independently selected from  $R^b$ ;

5  $R^{14}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, Cy, Cy- $C_{1-10}$  alkyl, Cy- $C_{2-10}$  alkenyl and Cy- $C_{2-10}$  alkynyl, wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents selected from phenyl and  $R^x$ , and Cy is optionally substituted with one to four substituents independently selected from  $R^y$ ;

10 or  $R^{13}$ ,  $R^{14}$  and the atoms to which they are attached together form a mono- or bicyclic ring containing 0-2 additional heteratoms selected from N, O and S;

$R^{15}$  is selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl, aryl- $C_{1-10}$  alkyl, heteroaryl, heteroaryl- $C_{1-10}$  alkyl, wherein  
15 alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents selected from  $R^x$ , and aryl and heteroaryl are optionally substituted with one to four substituents independently selected from  $R^y$ ;

or  $R^{14}$ ,  $R^{15}$  and the carbon to which they are attached form a 3-7 membered mono- or bicyclic ring containing 0-2 heteroatoms selected from  
20 N, O and S;

$R^a$  is selected from the group consisting of Cy and a group selected from  $R^x$ , wherein Cy is optionally substituted with one to four substituents independently selected from  $R^c$ ;

$R^b$  is selected from the group consisting of  $R^a$ ,  $C_{1-10}$  alkyl,  $C_{2-10}$   
25 alkenyl,  $C_{2-10}$  alkynyl, aryl  $C_{1-10}$  alkyl, heteroaryl  $C_{1-10}$  alkyl, wherein alkyl, alkenyl, alkynyl, aryl, heteroaryl are optionally substituted with a group independently selected from  $R^c$ ;

$R^c$  is selected from the group consisting of halogen,  $NO_2$ ,  $C(O)OR^f$ ,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy, aryl, aryl  $C_{1-4}$  alkyl, aryloxy, heteroaryl,  $NR^fR^g$ ,  
30  $R^fC(O)R^g$ ,  $NR^fC(O)NR^fR^g$ , and CN;

$R^d$  and  $R^e$  are independently selected from hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, Cy and Cy  $C_{1-10}$ alkyl, wherein alkyl, alkenyl, alkynyl and Cy are optionally substituted with one to four substituents independently selected from  $R^c$ ;

- 5           or  $R^d$  and  $R^e$  together with the atoms to which they are attached form a heterocyclic ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen;

$R^f$  and  $R^g$  are independently selected from hydrogen,  $C_{1-10}$  alkyl, Cy and Cy- $C_{1-10}$  alkyl wherein Cy is optionally substituted with  $C_{1-10}$  alkyl; or

- 10        $R^f$  and  $R^g$  together with the carbon to which they are attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

$R^h$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, cyano, aryl, aryl  $C_{1-10}$  alkyl, heteroaryl, heteroaryl  $C_{1-10}$  alkyl, and  $-SO_2R^i$ ; wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from  $R^a$ ; and aryl and heteroaryl are each optionally substituted with one to four substituents independently selected from  $R^b$ ;

- 15            $R^i$  is selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, and aryl; wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from  $R^c$ ;

- 20            $R^x$  is selected from the group consisting of  $-OR^d$ ,  $-NO_2$ , halogen,  $-S(O)_mR^d$ ,  $-SR^d$ ,  $-S(O)_2OR^d$ ,  $-S(O)_mNR^dR^e$ ,  $-NR^dR^e$ ,  $-O(CR^fR^g)_nNR^dR^e$ ,  $-C(O)R^d$ ,  $-CO_2R^d$ ,  $-CO_2(CR^fR^g)_nCONR^dR^e$ ,  $-OC(O)R^d$ ,  $-CN$ ,  $-C(O)NR^dR^e$ ,  $-NR^dC(O)R^e$ ,  $-OC(O)NR^dR^e$ ,  $-NR^dC(O)OR^e$ ,  $-NR^dC(O)NR^dR^e$ ,  $-CR^d(N-OR^e)$ ,  $CF_3$ , oxo,  $NR^dC(O)NR^dSO_2R^i$ ,  $NR^dS(O)_mR^e$ ,  $-OS(O)_2OR^d$ , and  $-OP(O)(OR^d)_2$ ;

- 25            $R^y$  is selected from the group consisting of  $R^x$ ,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl  $C_{1-10}$ alkyl, heteroaryl  $C_{1-10}$  alkyl, cycloalkyl,
- 30

heterocyclyl; wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from R<sup>a</sup>;

Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;

*m* is an integer from 1 to 2;

5 *n* is an integer from 1 to 10;

W is selected from the group consisting of carbon and nitrogen;

W' is selected from the group consisting of carbon, nitrogen, oxygen, sulfur, S(O) and S(O)<sub>2</sub>;

X' is selected from the group consisting of -C(O)OR<sup>d</sup>,  
10 -P(O)(OR<sup>d</sup>)(OR<sup>e</sup>), -P(O)(R<sup>d</sup>)(OR<sup>e</sup>), -S(O)<sub>m</sub>OR<sup>d</sup>, -C(O)NR<sup>d</sup>R<sup>h</sup>, and -5-tetrazolyl;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;

and further wherein the compound of formula IVa and/or IVb has a  
15 binding affinity to VLA-4 as expressed by an IC<sub>50</sub> of about 15 μM or less.

14. The method of Claim 13, wherein R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form a heteroaryl or substituted heteroaryl group having two nitrogen atoms in the  
20 heteroaryl ring.

15. The method of Claim 14, wherein R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form a pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-  
25 1,2,5-thiadiazole ring; wherein the pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl,  
30 substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.


-- 270 --

16. The method of Claim 13, wherein X' is -C(O)OR<sup>d</sup>.

17. The method of Claim 13, wherein the compound has formula Va, Vc, Vd, Ve or Vf:


5

10




Va

15



Vc

20





Vd

25

30

-- 271 --



15

wherein

20  $R^5$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

25  $R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $-SO_2R^{10}$  where  $R^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl; and

$R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl,

-- 272 --

substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$R^{16}$  and  $R^{17}$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen; and

$R^{18}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

$R^{20}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$R^{21}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;

$b$  is 1 or 2;


and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

18. The method of Claim 17, wherein the compound is selected from formula Vd, Ve or Vf.

19. A method for treating a disease mediated by VLA-4 in a patient, which method comprises administering a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of formula VIa and/or VIb:


-- 273 --

5



VIa

and



VIb

wherein, in formula VIa, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom  
 10 and W to which they are bound respectively, are joined to form an aryl,  
 cycloalkenyl, heteroaryl or heterocyclic group having at least five atoms in  
 the aryl, cycloalkenyl, heteroaryl or heterocyclic group and optionally  
 containing or additionally containing in the case of heteroaryl and  
 heterocyclic groups 1 to 3 heteroatoms selected from the group consisting of  
 15 oxygen, nitrogen and sulfur, and wherein the heteroaryl or heterocyclic group  
 is mono-cyclic;

in formula VIb, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W' to  
 which they are bound respectively, are joined to form a cycloalkyl,  
 cycloalkenyl or heterocyclic group having at least five atoms in the  
 20 cycloalkyl, cycloalkenyl or heterocyclic group and optionally containing or  
 additionally containing in the case of the heterocyclic group 1 to 3  
 heteroatoms selected from the group consisting of oxygen, nitrogen and  
 sulfur, and wherein the heterocyclic group is mono-cyclic;

and further wherein said aryl, cycloalkyl, cycloalkenyl, heteroaryl or  
 25 heterocyclic group of formula VIa or VIb is optionally substituted, on any  
 ring atom capable of substitution, with 1-3 substituents selected from the  
 group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl,  
 acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino,  
 alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino,  
 30 aminothiocabonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy,

substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen, hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where each R is independently hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, -N[S(O)<sub>2</sub>-R']<sub>2</sub> and -N[S(O)<sub>2</sub>-NR']<sub>2</sub> where each R' is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

R<sup>23</sup> is selected from the group consisting of hydrogen, C<sub>1-10</sub> alkyl optionally substituted with one to four substituents independently selected from R<sup>a</sup> and Cy optionally substituted with one to four substituents independently selected from R<sup>b</sup>;

R<sup>24</sup> is selected from the group consisting of Ar<sup>1</sup>-Ar<sup>2</sup>-C<sub>1-10</sub> alkyl, Ar<sup>1</sup>-Ar<sup>2</sup>-C<sub>2-10</sub> alkenyl, Ar<sup>1</sup>-Ar<sup>2</sup>-C<sub>2-10</sub> alkynyl, wherein Ar<sup>1</sup> and Ar<sup>2</sup> are independently aryl or heteroaryl each of which is optionally substituted with one to four substituents independently selected from R<sup>b</sup>; alkyl, alkenyl and



alkynyl are optionally substituted with one to four substituents independently selected from  $R^a$ ;

$R^{25}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl, aryl  $C_{1-10}$ alkyl, heteroaryl, and heteroaryl  $C_{1-10}$  alkyl, wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents selected from  $R^a$ , and aryl and heteroaryl are optionally substituted with one to four substituents independently selected from  $R^b$ ;

$R^a$  is selected from the group consisting of Cy,  $-OR^d$ ,  $-NO_2$ , halogen  $-S(O)_mR^d$ ,  $-SR^d$ ,  $-S(O)_2OR^d$ ,  $-S(O)_mNR^dR^e$ ,  $-NR^dR^e$ ,  $-O(CR^fR^g)_nNR^dR^e$ ,  $-C(O)R^d$ ,  $-CO_2R^d$ ,  $-CO_2(CR^fR^g)_nCONR^dR^e$ ,  $-OC(O)R^d$ ,  $-CN$ ,  $-C(O)NR^dR^e$ ,  $-NR^dC(O)R^e$ ,  $-OC(O)NR^dR^e$ ,  $-NR^dC(O)OR^e$ ,  $-NR^dC(O)NR^dR^e$ ,  $-CR^d(N-OR^e)$ ,  $CF_3$ , and  $-OCF_3$ ;

wherein Cy is optionally substituted with one to four substituents independently selected from  $R^c$ ;

$R^b$  is selected from the group consisting of  $R^a$ ,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl  $C_{1-10}$  alkyl, heteroaryl  $C_{1-10}$ alkyl,

wherein alkyl, alkenyl, aryl, heteroaryl are optionally substituted with a group independently selected from  $R^c$ ;

$R^c$  is selected from the group consisting of halogen, amino, carboxy,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy, aryl, aryl  $C_{1-4}$ alkyl, hydroxy,  $CF_3$ , and aryloxy;

$R^d$  and  $R^e$  are independently selected from hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, Cy and Cy  $C_{1-10}$ alkyl, wherein alkyl, alkenyl, alkynyl and Cy are optionally substituted with one to four substituents independently selected from  $R^c$ ; or  $R^d$  and  $R^e$  together with the atoms to which they are attached form a heterocyclic ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen;

$R^f$  and  $R^g$  are independently selected from hydrogen,  $C_{1-10}$  alkyl, Cy and Cy- $C_{1-10}$  alkyl; or  $R^f$  and  $R^g$  together with the carbon to which they are

attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

$R^{h'}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, cyano, aryl, aryl  $C_{1-10}$  alkyl, heteroaryl, 5 heteroaryl  $C_{1-10}$  alkyl, or  $-SO_2R^{i'}$ ;  
wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from  $R^{a'}$ ; and aryl and heteroaryl are each optionally substituted with one to four substituents independently selected from  $R^{b'}$ ;

10  $R^{i'}$  is selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, and aryl;

wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from  $R^{c'}$ ;

Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;

15  $X''$  is selected from the group consisting of  $-C(O)OR^d$ ,  $-P(O)(OR^d)(OR^e)$ ,  $-P(O)(R^d)(OR^e)$ ,  $-S(O)_mOR^d$ ,  $-C(O)NR^dR^{h'}$ , and -5-tetrazolyl;

$m$  is an integer from 1 to 2;

$n$  is an integer from 1 to 10;

20 and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;

and further wherein the compound of formula IVa and/or IVb has a binding affinity to VLA-4 as expressed by an  $IC_{50}$  of about  $15\mu M$  or less.

25 20. The method of Claim 19, wherein  $R^1$  and  $R^2$ , together with the carbon atom and W to which they are bound respectively, are joined to form a heteroaryl or substituted heteroaryl group having two nitrogen atoms in the heteroaryl ring.


-- 277 --

21. The method of Claim 20, wherein  $R^1$  and  $R^2$ , together with the carbon atom and W to which they are bound respectively, are joined to form a pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring; wherein the pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.


22. The method of Claim 19, wherein  $X''$  is  $-C(O)OR^d$ .

23. The method of Claim 19, wherein  $R^{24}$  is  $-\text{CH}_2-\text{Ar}^2-\text{Ar}^1$  and  $R^{25}$  is hydrogen.

24. The method of Claim 19, wherein the compound has formula VIIa, VIIc, VIId, VIIe or VIIf:




VIIa



VIIc


-- 278 --

5



VIIId


10



VIIe

15

20



VIIIf

wherein

25  $R^5$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

30  $R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl,

heteroaryl, substituted heteroaryl, and  $-SO_2R^{10}$  where  $R^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl; and

5  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

10  $R^{16}$  and  $R^{17}$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen; and

15  $R^{18}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

20  $R^{20}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$R^{21}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;

25  $b$  is 1 or 2;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

25. The method of Claim 24, wherein the compound is selected  
30 from formula VIId, VIIe or VIIf.



15

25

30

-- 281 --

consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl,  
 acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino,  
 alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino,  
 aminothiocabonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy,  
 5 substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen,  
 hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino,  
 guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted  
 thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted  
 thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl,  
 10 substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy,  
 substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy,  
 heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino,  
 oxythiocarbonylamino,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-}$   
 aryl,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted}$   
 15 heteroaryl,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{OSO}_2\text{-}$   
 NRR where each R is independently hydrogen or alkyl,  $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  
 $-\text{NRS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-}$   
 20 alkyl,  $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-}$   
 substituted aryl,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted}$   
 heteroaryl,  $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heterocyclic}$   
 where R is hydrogen or alkyl,  $-\text{N}[\text{S}(\text{O})_2\text{-R}']_2$  and  $-\text{N}[\text{S}(\text{O})_2\text{-NR}']_2$  where  
 each R' is independently selected from the group consisting of alkyl,  
 25 substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl,  
 heterocyclic and substituted heterocyclic;

$\text{R}^3$  is  $-(\text{CH}_2)_x\text{-Ar-R}^9$ , where Ar is aryl, substituted aryl, heteroaryl and  
 substituted heteroaryl;  $\text{R}^9$  is selected from the group consisting of acyl,  
 acylamino, acyloxy, aminoacyl, aminocarbonylamino,  
 30 aminothiocabonylamino, aminocarbonyloxy, oxycarbonylamino,

oxythiocarbonylamino, thioamidino, thiocarbonylamino, aminosulfonylamino, aminosulfonyloxy, aminosulfonyl, oxysulfonylamino and oxysulfonyl; and  $x$  is an integer from 0 to 4;

$R^{3'}$  is selected from the group consisting of hydrogen, isopropyl, 5  $-CH_2Z$  where  $Z$  is selected from the group consisting of hydrogen, hydroxyl, acylamino, alkyl, alkoxy, aryloxy, aryl, aryloxyaryl, carboxyl, carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl, carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic, 10 carboxyl-substituted heterocyclic, cycloalkyl, substituted alkyl, substituted alkoxy, substituted aryl, substituted aryloxy, substituted aryloxyaryl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

$Q$  is selected from the group consisting of  $-O-$ ,  $-S-$ ,  $-S(O)-$ ,  $-S(O)_2$ , 15 and  $-NR^4-$ ;

$R^4$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic 20 or, optionally,  $R^4$  and  $R^1$  or  $R^4$  and  $R^2$ , together with the atoms to which they are bound, are joined to form a heteroaryl, a substituted heteroaryl, a heterocyclic or a substituted heterocyclic group;

$W$  is selected from the group consisting of nitrogen and carbon; and

$W'$  is selected from the group consisting of nitrogen, carbon, oxygen, 25 sulfur,  $S(O)$ , and  $S(O)_2$ ;

$X$  is selected from the group consisting of hydroxyl, alkoxy, substituted alkoxy, alkenoxy, substituted alkenoxy, cycloalkoxy, substituted cycloalkoxy, cycloalkenoxo, substituted cycloalkenoxo, aryloxy, substituted aryloxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, 30 substituted heterocyclyloxy and  $-NR''R''$  where each  $R''$  is independently




-- 283 --

selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

5 and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;

and further wherein the compound of formula Ia and/or Ib has a binding affinity to VLA-4 as expressed by an  $IC_{50}$  of about  $15\mu M$  or less.

10 28. The compound of Claim 27, wherein  $R^3$  is a group of the formula:



15

wherein  $R^9$  and  $x$  are as defined in Claim 27; and  $R^{3'}$  is hydrogen.

20 29. The compound of Claim 28, wherein  $R^9$  is in the *para* position of the phenyl ring, and  $x$  is an integer from 1 to 4.

30. The compound of Claim 29, wherein  $R^9$  is selected from the group consisting of  $-O-Z-NR^{11}R^{11'}$  and  $-O-Z-R^{12}$  wherein  $R^{11}$  and  $R^{11'}$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, and where  $R^{11}$  and  $R^{11'}$  are joined to form a heterocycle or a substituted heterocycle,  $R^{12}$  is selected from the group consisting of heterocycle and substituted heterocycle, and  $Z$  is selected from the group consisting of  $-C(O)-$  and  $-SO_2-$ .

30

-- 284 --


31. The compound of Claim 30, wherein Z is -C(O)-.

32. The compound of Claim 31, wherein R<sup>9</sup> is -OC(O)NR<sup>11</sup>R<sup>11'</sup>.


5 33. The compound of Claim 27, wherein Q is -NR<sup>4</sup>-.

34. The compound of Claim 27, wherein the compound has formula IIa or IIb:

10



15



20

wherein

ring A and ring B independently form a heteroaryl or substituted  
25 heteroaryl group having two nitrogen atoms in the heteroaryl ring;

R<sup>5</sup> is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

R<sup>6</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and -SO<sub>2</sub>R<sup>10</sup> where R<sup>10</sup> is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl;

or optionally, one of, R<sup>4</sup> and ring A, R<sup>4</sup> and R<sup>5</sup>, R<sup>4</sup> and R<sup>6</sup>, or R<sup>5</sup> and R<sup>6</sup>, together with the atoms to which they are bound, can be joined to form a heterocyclic or substituted heterocyclic ring;


and enantiomers, diastereomers and pharmaceutically acceptable salts thereof; and provided that ring B does not form a 6-amino or substituted amino pyrimidin-4-yl group.

35. The compound of Claim 34, wherein ring A forms a pyridazine, pyrimidine or pyrazine ring, wherein the pyridazine, pyrimidine or pyrazine ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen, and

ring B forms a pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or a 1,1-dioxo-1,2,5-thiadiazole ring, wherein the pyridazine, pyrimidine or pyrazine ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.


36. The compound of Claim 27, wherein the compound has formula IIIa, IIIc, IIId, IIIe or IIIf:

5




IIIa

10



IIIc


15



III d

20


25



IIIe

30

5



IIIf

10

wherein

$R^{4'}$  is selected from the group consisting of hydrogen and alkyl or, optionally, one of,  $R^{4'}$  and  $R^5$ ,  $R^{4'}$  and  $R^6$ ,  $R^5$  and  $R^6$ ,  $R^5$  and  $R^8$ , or  $R^6$  and  $R^8$ , together with the atoms to which they are bound, are joined to form a heterocyclic, a substituted heterocyclic, a heteroaryl or substituted heteroaryl group optionally containing from 1 to 3 additional hetero ring atoms selected from the group consisting of oxygen, nitrogen and sulfur;

15

$R^{4''}$  is selected from the group consisting of hydrogen and alkyl;

20

$R^5$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

25

$R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $-SO_2R^{10}$  where  $R^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl;

30

$R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl,

-- 288 --

substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$R^{16}$  and  $R^{17}$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen; and

$R^{18}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

$R^{20}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;


$R^{21}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;

$b$  is 1 or 2;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

37. The compound of Claim 36, wherein the compound is selected from formula IIIId, IIIe or IIIf.

38. A compound of formula IVa:



wherein R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form a heteroaryl group having two nitrogen atoms in the heteroaryl ring;

5 and further wherein said heteroaryl group is optionally substituted, on any ring atom capable of substitution, with 1-3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, 10 aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen, hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted 15 thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>- 20 aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where each R is independently hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>- 25 heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, -N[S(O)<sub>2</sub>-R']<sub>2</sub> and -N[S(O)<sub>2</sub>-NR']<sub>2</sub> where each R' is 30 independently selected from the group consisting of alkyl, substituted alkyl,

aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

$R^{13}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl, Cy, and  $Cy-C_{1-10}$  alkyl, wherein alkyl is optionally substituted with one to four substituents independently selected from  $R^a$ ; and Cy is optionally substituted with one to four substituents independently selected from  $R^b$ ;

$R^{14}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, Cy,  $Cy-C_{1-10}$  alkyl,  $Cy-C_{2-10}$  alkenyl and  $Cy-C_{2-10}$  alkynyl, wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents selected from phenyl and  $R^x$ , and Cy is optionally substituted with one to four substituents independently selected from  $R^y$ ;

or  $R^{13}$ ,  $R^{14}$  and the atoms to which they are attached together form a mono- or bicyclic ring containing 0-2 additional heteratoms selected from N, O and S;

$R^{15}$  is selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl, aryl- $C_{1-10}$  alkyl, heteroaryl, heteroaryl- $C_{1-10}$  alkyl, wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents selected from  $R^x$ , and aryl and heteroaryl are optionally substituted with one to four substituents independently selected from  $R^y$ ;

or  $R^{14}$ ,  $R^{15}$  and the carbon to which they are attached form a 3-7 membered mono- or bicyclic ring containing 0-2 heteroatoms selected from N, O and S;

$R^a$  is selected from the group consisting of Cy and a group selected from  $R^x$ , wherein Cy is optionally substituted with one to four substituents independently selected from  $R^c$ ;

$R^b$  is selected from the group consisting of  $R^a$ ,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl  $C_{1-10}$  alkyl, heteroaryl  $C_{1-10}$  alkyl, wherein alkyl, alkenyl, alkynyl, aryl, heteroaryl are optionally substituted with a group independently selected from  $R^c$ ;

$R^c$  is selected from the group consisting of halogen,  $NO_2$ ,  $C(O)OR^f$ ,



-- 291 --

$C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy, aryl, aryl  $C_{1-4}$  alkyl, aryloxy, heteroaryl,  $NR^fR^g$ ,  $R^fC(O)R^g$ ,  $NR^fC(O)NR^fR^g$ , and CN;

$R^d$  and  $R^e$  are independently selected from hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, Cy and Cy  $C_{1-10}$  alkyl, wherein alkyl, alkenyl, alkynyl and Cy are optionally substituted with one to four substituents independently selected from  $R^c$ ;

or  $R^d$  and  $R^e$  together with the atoms to which they are attached form a heterocyclic ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen;

$R^f$  and  $R^g$  are independently selected from hydrogen,  $C_{1-10}$  alkyl, Cy and Cy- $C_{1-10}$  alkyl wherein Cy is optionally substituted with  $C_{1-10}$  alkyl; or  $R^f$  and  $R^g$  together with the carbon to which they are attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

$R^h$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, cyano, aryl, aryl  $C_{1-10}$  alkyl, heteroaryl, heteroaryl  $C_{1-10}$  alkyl, and  $-SO_2R^i$ ; wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from  $R^a$ ; and aryl and heteroaryl are each optionally substituted with one to four substituents independently selected from  $R^b$ ;

$R^i$  is selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, and aryl; wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from  $R^c$ ;

$R^x$  is selected from the group consisting of  $-OR^d$ ,  $-NO_2$ , halogen,  $-S(O)_mR^d$ ,  $-SR^d$ ,  $-S(O)_2OR^d$ ,  $-S(O)_mNR^dR^e$ ,  $-NR^dR^e$ ,  $-O(CR^fR^g)_nNR^dR^e$ ,  $-C(O)R^d$ ,  $-CO_2R^d$ ,  $-CO_2(CR^fR^g)_nCONR^dR^e$ ,  $-OC(O)R^d$ ,  $-CN$ ,  $-C(O)NR^dR^e$ ,  $-NR^dC(O)R^e$ ,  $-OC(O)NR^dR^e$ ,  $-NR^dC(O)OR^e$ ,  $-NR^dC(O)NR^dR^e$ ,  $-CR^d(N-OR^e)$ ,  $CF_3$ , oxo,  $NR^dC(O)NR^dSO_2R^i$ ,  $NR^dS(O)_mR^e$ ,  $-OS(O)_2OR^d$ , and  $-OP(O)(OR^d)_2$ ;

$R^y$  is selected from the group consisting of  $R^x$ ,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl  $C_{1-10}$ alkyl, heteroaryl  $C_{1-10}$  alkyl, cycloalkyl, heterocyclyl; wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from  $R^x$ ;

5        Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;

$m$  is an integer from 1 to 2;

$n$  is an integer from 1 to 10;

W is selected from the group consisting of carbon and nitrogen;

10        W' is selected from the group consisting of carbon, nitrogen, oxygen, sulfur, S(O) and S(O)<sub>2</sub>;

$X'$  is selected from the group consisting of -C(O)OR<sup>d</sup>, -P(O)(OR<sup>d</sup>)(OR<sup>e</sup>), -P(O)(R<sup>d</sup>)(OR<sup>e</sup>), -S(O)<sub>m</sub>OR<sup>d</sup>, -C(O)NR<sup>d</sup>R<sup>h</sup>, and -5-tetrazolyl;

15        and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;

and further wherein the compound of formula IV has a binding affinity to VLA-4 as expressed by an IC<sub>50</sub> of about 15 $\mu$ M or less;

and provided that when R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form a 2-arylpyrimidin-4-yl group and R<sup>14</sup> is hydrogen, then R<sup>15</sup> is not alkyl of from 1 to 6 carbon atoms optionally substituted with hydroxyl; and when R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form a 5-arylpyrazin-2-yl group and R<sup>14</sup> is hydrogen, then R<sup>15</sup> is not 4-hydroxybenzyl.

25

39.    The compound of Claim 38, wherein R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form a pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring; wherein the pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or a 1,1-dioxo-1,2,5-thiadiazole ring is optionally

30

-- 293 --

substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.


5

40. The compound of Claim 38, wherein X' is -C(O)OR<sup>d</sup>.


41. The compound of Claim 38, wherein the compound has formula Va, Vc, Vd, Ve or Vf:

10

15




20




25

30




5



Ve

10

15



Vf

wherein

20

$R^5$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

25

$R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $-SO_2R^{10}$  where  $R^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl; and

30

$R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

5         $R^{16}$  and  $R^{17}$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen; and

10         $R^{18}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

15         $R^{20}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;


20         $R^{21}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;

$b$  is 1 or 2;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

25        42. The compound of Claim 41, wherein the compound is selected from formula Vd, Ve or Vf.

43. A compound of formula VIa and/or VIb:



10            wherein, in formula VIa, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form an aryl, cycloalkenyl, heteroaryl or heterocyclic group having at least five atoms in the aryl, cycloalkenyl, heteroaryl or heterocyclic group and optionally containing or additionally containing in the case of heteroaryl and

15            heterocyclic groups 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heteroaryl or heterocyclic group is mono-cyclic;

              in formula VIb, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W' to which they are bound respectively, are joined to form a cycloalkyl, cycloalkenyl or heterocyclic group having at least five atoms in the cycloalkyl, cycloalkenyl or heterocyclic group and optionally containing or additionally containing in the case of the heterocyclic group 1 to 3

20            heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heterocyclic group is mono-cyclic;

25            and further wherein said aryl, cycloalkyl, cycloalkenyl, heteroaryl or heterocyclic group of formula VIa or VIb is optionally substituted, on any ring atom capable of substitution, with 1-3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino,

30

aminothiocabonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy,  
 substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen,  
 hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino,  
 guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted  
 5 thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted  
 thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl,  
 substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy,  
 substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy,  
 heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino,  
 10 oxythiocarbonylamino,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-}$   
 $\text{aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted}$   
 $\text{heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{OSO}_2\text{-}$   
 $\text{NRR}$  where each R is independently hydrogen or alkyl,  
 $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-}$   
 15  $\text{substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  
 $-\text{NRS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-}$   
 $\text{alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-}$   
 $\text{substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted}$   
 $\text{heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heterocyclic}$   
 20 where R is hydrogen or alkyl,  $-\text{N}[\text{S}(\text{O})_2\text{-R}']_2$  and  $-\text{N}[\text{S}(\text{O})_2\text{-NR}']_2$  where  
 each R' is independently selected from the group consisting of alkyl,  
 substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl,  
 heterocyclic and substituted heterocyclic;

$\text{R}^{23}$  is selected from the group consisting of hydrogen,  $\text{C}_{1-10}$  alkyl  
 25 optionally substituted with one to four substituents independently selected  
 from  $\text{R}^{\text{a}}$  and Cy optionally substituted with one to four substituents  
 independently selected from  $\text{R}^{\text{b}}$ ;

$\text{R}^{24}$  is selected from the group consisting of  $\text{Ar}^1\text{-Ar}^2\text{-C}_{1-10}$  alkyl,  
 $\text{Ar}^1\text{-Ar}^2\text{-C}_{2-10}$  alkenyl,  $\text{Ar}^1\text{-Ar}^2\text{-C}_{2-10}$  alkynyl, wherein  $\text{Ar}^1$  and  $\text{Ar}^2$  are  
 30 independently aryl or heteroaryl each of which is optionally substituted with

one to four substituents independently selected from  $R^b$ ; alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents independently selected from  $R^a$ ;

5  $R^{25}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl, aryl  $C_{1-10}$ alkyl, heteroaryl, and heteroaryl  $C_{1-10}$  alkyl, wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents selected from  $R^a$ , and aryl and heteroaryl are optionally substituted with one to four substituents independently selected from  $R^b$ ;

10  $R^a$  is selected from the group consisting of Cy,  $-OR^d$ ,  $-NO_2$ , halogen  $-S(O)_mR^d$ ,  $-SR^d$ ,  $-S(O)_2OR^d$ ,  $-S(O)_mNR^dR^c$ ,  $-NR^dR^c$ ,  $-O(CR^fR^g)_nNR^dR^c$ ,  $-C(O)R^d$ ,  $-CO_2R^d$ ,  $-CO_2(CR^fR^g)_nCONR^dR^c$ ,  $-OC(O)R^d$ ,  $-CN$ ,  $-C(O)NR^dR^c$ ,  $-NR^dC(O)R^c$ ,  $-OC(O)NR^dR^c$ ,  $-NR^dC(O)OR^c$ ,  $-NR^dC(O)NR^dR^c$ ,  $-CR^d(N-OR^c)$ ,  $CF_3$ , and  $-OCF_3$ ;

15 wherein Cy is optionally substituted with one to four substituents independently selected from  $R^c$ ;

$R^b$  is selected from the group consisting of  $R^a$ ,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl  $C_{1-10}$  alkyl, heteroaryl  $C_{1-10}$ alkyl,

20 wherein alkyl, alkenyl, aryl, heteroaryl are optionally substituted with a group independently selected from  $R^c$ ;

$R^c$  is selected from the group consisting of halogen, amino, carboxy,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy, aryl, aryl  $C_{1-4}$ alkyl, hydroxy,  $CF_3$ , and aryloxy;

25  $R^d$  and  $R^e$  are independently selected from hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, Cy and Cy  $C_{1-10}$ alkyl, wherein alkyl, alkenyl, alkynyl and Cy are optionally substituted with one to four substituents independently selected from  $R^c$ ; or  $R^d$  and  $R^e$  together with the atoms to which they are attached form a heterocyclic ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen;

30  $R^f$  and  $R^g$  are independently selected from hydrogen,  $C_{1-10}$  alkyl, Cy and Cy- $C_{1-10}$  alkyl; or  $R^f$  and  $R^g$  together with the carbon to which they are



attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

$R^{h'}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, cyano, aryl, aryl  $C_{1-10}$  alkyl, heteroaryl, heteroaryl  $C_{1-10}$  alkyl, or  $-SO_2R^{i'}$ ;

wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from  $R^{a'}$ ; and aryl and heteroaryl are each optionally substituted with one to four substituents independently selected from  $R^{b'}$ ;

$R^{i'}$  is selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, and aryl;

wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from  $R^{c'}$ ;

Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;

$X''$  is selected from the group consisting of  $-C(O)OR^d$ ,  $-P(O)(OR^d)(OR^e)$ ,  $-P(O)(R^d)(OR^e)$ ,  $-S(O)_mOR^d$ ,  $-C(O)NR^dR^h$ , and -5-tetrazolyl;

$m$  is an integer from 1 to 2;

$n$  is an integer from 1 to 10;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;

and further wherein the compounds of formula VIa and/or VIb have a binding affinity to VLA-4 as expressed by an  $IC_{50}$  of about  $15\mu M$  or less.

44. The compound of Claim 43, wherein  $R^1$  and  $R^2$ , together with the carbon atom and W to which they are bound respectively, are joined to form a heteroaryl or substituted heteroaryl group having two nitrogen atoms in the heteroaryl ring.

-- 300 --

45. The compound of Claim 44, wherein  $R^1$  and  $R^2$ , together with the carbon atom and W to which they are bound respectively, are joined to form a pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring; wherein the pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.

10


46. The compound of Claim 43, wherein  $X''$  is  $-C(O)OR^d$ .

47. The compound of Claim 43, wherein  $R^{24}$  is  $-\text{CH}_2-\text{Ar}^2-\text{Ar}^1$  and  $R^{25}$  is hydrogen.

15


48. The compound of Claim 43, wherein the compound has formula VIIa, VIIc, VIId, VIIe or VIIf:

20



VIIa


25



VIIc


30

5



VIIId


10



VIIe

15

20



VIIIf

wherein

25  $R^5$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

30  $R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl,

-- 302 --

heteroaryl, substituted heteroaryl, and  $\text{-SO}_2\text{R}^{10}$  where  $\text{R}^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl; and

5            $\text{R}^7$  and  $\text{R}^8$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$\text{R}^{16}$  and  $\text{R}^{17}$  are independently selected from the group consisting of  
10       hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen; and

$\text{R}^{18}$  is selected from the group consisting of alkyl, substituted alkyl,  
15       alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

$\text{R}^{20}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl,  
20       heterocyclic, substituted heterocyclic and halogen;

$\text{R}^{21}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;

25           ***b*** is 1 or 2;

          and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

49.       The compound of Claim 48, wherein the compound is selected  
30       from formula VIId, VIIe or VIIf.

50. A compound selected from the group consisting of:
- N*-(2-chloro-5-nitropyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 5 *N*-[5-(*N*-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester,
- N*-[5-(*N*-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 10 *N*-[5-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester,
- N*-[5-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 15 *N*-[5-(*N,N*-di-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-[5-[*N*-(1-*N'*-methylpyrazol-4-ylsulfonyl)-*N*-methylamino]pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 20 *N*-[5-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine isopropyl ester,
- 25 *N*-[5-(*N*-methyl-*N*-3-pyridylsulfonylamino)pyrimidin-4-yl]-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester,
- N*-(5-(*N*-methyl-*N*-(1-butylpyrazol-4-yl)sulfonylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 30 *N*-(5-(2,4-dimethoxypyrimidin-5-yl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(2,6-difluorophenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 35 *N*-(5-(2-hydroxymethylphenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 40 *N*-(2-(*N*-cyclohexylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,

- N*-(2-(*N*-methyl-*N*-(1-methylpiperidin-4-yl)amino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 5 *N*-(2-(*N*-ethyl-*N*-isopropylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 10 *N*-(5-isopropylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-butylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 15 *N*-(2-(*N*-ethyl-*N*-propylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 20 *N*-(2-(*N,N*-diethylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-ethylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 25 *N*-(5-benzyloxy)pyrimidin-4-yl)-L-phenylalanine,
- N*-(5-benzyloxy)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 30 *N*-(5-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl)-L-phenylalanine,
- N*-(5-(*N*-methyl-*N*-3-pyridinesulfonylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 35 *N*-(5-phenylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 40 *N*-(3-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrazin-2-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(2,2,2-trifluoroethyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,

- N*-(5-(*N*-methyl-*N*-3-pyridinesulfonylamino)pyrimidin-4-yl)-L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine isopropyl ester,
- 5 *N*-(5-benzylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(*N*-methyl-*N*-3-pyridinesulfonylamino)pyrimidin-4-yl)-L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine tert-butyl ester,
- 10 *N*-(5-(2-trifluoromethylphenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(2-*N,N*-dimethylcarbamylethyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 15 *N*-(5-(*N*-methyl-*N*-3-(1-methylpyrazole)sulfonylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine isopropyl ester,
- N*-(6-phenylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 20 *N*-(6-(2-trifluoromethylphenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(6-(2-hydroxymethylphenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 25 *N*-(5-cyclohexylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 30 *N*-(2-(*N*-methyl-*N*-2-furanmethylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-4-chlorophenylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 35 *N*-(5-(3-thienyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 40 *N*-(5-(2-thienyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-2-hydroxyethylamino)-5-(2-fluorophenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 45

- N*-(5-(piperidin-1-yl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 5 *N*-(5-(1-propylbutyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 10 *N*-(2-(*N*-methyl-*N*-cyclobutylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 15 *N*-(2-(*N,N*-bis-(2-hydroxyethyl)amino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 20 *N*-(2-(*N*-methyl-*N*-phenylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 25 *N*-(2-(*N*-methyl-*N*-3-methylbutylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 30 *N*-(2-(*N*-methylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 35 *N*-(2-(2-tolyl)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 40 *N*-(2-(*N*-methyl-*N*-2-hydroxyethylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 45 *N*-(2-(*N*-methyl-*N*-2-methylpropylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 40 *N*-(2-(*N*-methyl-*N*-propylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 45 *N*-(2-(*N,N*-dimethylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,



-- 307 --

- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(3-pyridyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- 5 *N*-(5-(2-phenyl-2,2-difluoroethyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- N*-(5-(2-phenyl-2,2-difluoroethyl)-6-chloropyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- 10 *N*-(5-(2-phenylethyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-cyclohexylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- 15 *N*-(5-propylpyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- N*-(5-(2-methoxyphenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- 20 *N*-(5-(2-fluorophenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- N*-(2-(*N*-Methyl-*N*-isopropylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- 25 *N*-(2-(*N*-isopropylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,
- 30 *N*-(5-(2-phenylethyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine isopropyl ester,
- N*-(3-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrazin-2-yl)-L-phenylalanine isopropyl ester,
- 35 *N*-(5-(2-phenylethyl)pyrimidin-4-yl)-L-phenylalanine isopropyl ester,
- N*-(5-(*N*-methyl-*N*-3-pyridinesulfonylamino)pyrimidin-4-yl)-L-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine,
- 40 *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamoyloxy)phenylalanine,

-- 308 --

- N*-(5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine isopropyl ester,
- 5 *N*-(5-(3-nitrophenyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(3-pyridyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 10 *N*-(5-(2-phenylethyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-*N,N*-dimethylamino-5-(*N*-methyl-*N*-4-toluenesulfonylamino)pyrimidin-4-yl)-L-phenylalanine,
- 15 *N*-(5-(2-tolyl)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2-methoxyphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- 20 *N*-(2-(*N*-methyl-*N*-isopropylamino)-5-(2-fluorophenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- N*-(2-(*N*-methyl-*N*-isopropylamino)-5-(2-fluorophenyl)pyrimidin-4-yl)-L-4-(2-methoxyphenyl)phenylalanine,
- 25 *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2,6-difluorophenyl)pyrimidin-4-yl)-L-4-(2,6-difluorophenyl)phenylalanine,
- 30 *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2-hydroxymethylphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- N*-(2-(*N,N*-bis-(2-hydroxyethyl)amino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- 35 *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2-trifluoromethylphenyl)pyrimidin-4-yl)-L-4-(2-cyanophenyl)phenylalanine,
- 40 *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(3-thienyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- 45

- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2-thienyl)pyrimidin-4-yl)-L-4-(4-trifluoromethylphenyl)phenylalanine,
- 5 *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(3-pyridyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(3-nitrophenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- 10 *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2,6-dichlorophenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(4-pyridyl)pyrimidin-4-yl)-L-4-(3-hydroxymethylphenyl)phenylalanine,
- 15 *N*-(2-(*N*-ethyl-*N*-isopropylamino)-5-(2,6-dimethoxyphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-(2,3-dichlorophenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- 20 *N*-(2-(*N*-methyl-*N*-ethylamino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(2-cyanophenyl)phenylalanine,
- N*-(2-(*N*-methyl-*N*-isopropylamino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(3-pyridyl)phenylalanine,
- 25 *N*-(2-(*N,N*-bis-(2-hydroxyethyl)amino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(2-cyanophenyl)phenylalanine,
- 30 *N*-(2-(*N*-methyl-*N*-(1-methylpiperidin-4-yl)amino)-5-(2-cyanophenyl)pyrimidin-4-yl)-L-4-(2,6-difluorophenyl)phenylalanine,
- N*-(2-(*N*-ethyl-*N*-isopropylamino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(*o*-tolyl)phenylalanine,
- 35 *N*-(2-(*N*-methyl-*N*-4-chlorophenylamino)-5-(2,4,6-trimethylphenyl)pyrimidin-4-yl)-L-4-(2,6-dimethoxyphenyl)phenylalanine,
- 40 *N*-(5-(*N*-methyl-*N*-2-(phenyl)ethylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- N*-(5-(*N*-methyl-*N*-hexylamino)pyrimidin-4-yl)-L-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 45

- N*-(5-(*N*-methyl-*N*-isopropylamino)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 5      *N*-(5-(*N*-methyl-*N-tert*-butylamino)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 10      *N*-(5-(*N*-ethyl-*N*-isopropylamino)pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 15      *N*-(5-(*N*-methyl-*N*-2-(4-pyridyl)ethyl-pyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 20      *N*-(5-(*N*-methyl-*N*-2-(phenyl)ethylamino)pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,
- 25      *N*-(5-(*N*-methyl-*N*-hexylamino)pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,
- 30      *N*-(5-(*N*-methyl-*N*-isopropylamino)pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,
- 35      *N*-(5-(*N*-methyl-*N-tert*-butylamino)pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,
- 40      *N*-(5-(*N*-ethyl-*N*-isopropylamino)pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,
- 45      *N*-(5-(*N*-methyl-*N*-2-(4-pyridyl)ethyl-pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,
- 50      *N*-(2-(*N*-methyl-*N*-cyclohexylamino)-5-ethylpyrimidin-4-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 55      *N*-(4-(*N,N*-di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-*L*-tyrosine,
- 60      *N*-(4-(*N,N*-di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine,
- 65      *N*-(4-(*N,N*-dimethylamino)-1-oxo-1,2,5-thiadiazol-3-yl)-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine *tert*-butyl ester,
- 70      *N*-[4-(2-(3-methylphenylaminocarbonylamino)ethyl-1-ylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl]-*L*-4-(*N,N*-dimethylcarbamyloxy)phenylalanine

-- 311 --

*N*-(4-(*N,N*-di-*n*-hexylamino)-1,1-dioxo-1,2,5-thiadiazol-3-yl)-*L*-4-(4-methylpiperazin-1-ylcarbonyloxy)phenylalanine,

5 *N*-(5-(2,2,2-trifluoroethyl)pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,

*N*-(2-(*N*-cyclohexyl-*N*-methyl)-5-(2-tolyl)pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,

10 *N*-(5-(2-fluorophenyl)pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,


15 *N*-(2-(*N*-methyl-*N*-propyl)-5-(2-tolyl)pyrimidin-4-yl)-*L*-4-(2,6-dimethoxyphenyl)phenylalanine,

*N*-(3-chloropyrazin-2-yl)-*L*-4-[1-(*tert*-butoxycarbonyl)piperidin-4-ylcarbonylamino]phenylalanine ethyl ester,

20 and pharmaceutically acceptable salts thereof.

51. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of formula Ia and/or Ib:

25



30

Ia

Ib

wherein, in formula Ia,  $R^1$  and  $R^2$ , together with the carbon atom and W to which they are bound respectively, are joined to form an aryl, cycloalkenyl, heteroaryl or heterocyclic group having at least five atoms in the aryl, cycloalkenyl, heteroaryl or heterocyclic group and optionally containing or additionally containing in the case of heteroaryl and

35

heterocyclic groups 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heteroaryl or heterocyclic group is mono-cyclic;

in formula Ib,  $R^1$  and  $R^2$ , together with the carbon atom and W' to  
5 which they are bound respectively, are joined to form a cycloalkyl, cycloalkenyl or heterocyclic group having at least five atoms in the cycloalkyl, cycloalkenyl or heterocyclic group and optionally containing or additionally containing in the case of the heterocyclic group 1 to 3  
heteroatoms selected from the group consisting of oxygen, nitrogen and  
10 sulfur, and wherein the heterocyclic group is mono-cyclic;

and further wherein said aryl, cycloalkyl, cycloalkenyl, heteroaryl or heterocyclic group of formula Ia or Ib is optionally substituted, on any ring atom capable of substitution, with 1-3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl,  
15 acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen, hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino,  
20 guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy,  
25 heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{OSO}_2\text{-NRR}$  where each R is independently hydrogen or alkyl,  $-\text{NRS}(\text{O})_2\text{-alkyl}$ , -

NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -  
 NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl,  
 -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-  
 alkyl, -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-  
 5 substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted  
 heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic  
 where R is hydrogen or alkyl, -N[S(O)<sub>2</sub>-R']<sub>2</sub> and -N[S(O)<sub>2</sub>-NR']<sub>2</sub> where  
 each R' is independently selected from the group consisting of alkyl,  
 substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl,  
 10 heterocyclic and substituted heterocyclic;

R<sup>3</sup> is -(CH<sub>2</sub>)<sub>x</sub>-Ar-R<sup>9</sup>, where Ar is aryl, substituted aryl, heteroaryl and  
 substituted heteroaryl; R<sup>9</sup> is selected from the group consisting of acyl,  
 acylamino, acyloxy, aminoacyl, aminocarbonylamino,  
 aminothiocarbonylamino, aminocarbonyloxy, oxycarbonylamino,  
 15 oxythiocarbonylamino, thioamidino, thiocarbonylamino,  
 aminosulfonylamino, aminosulfonyloxy, aminosulfonyl, oxysulfonylamino  
 and oxysulfonyl, and x is an integer from 0 to 4;

R<sup>3'</sup> is selected from the group consisting of hydrogen, isopropyl,  
 -CH<sub>2</sub>Z where Z is selected from the group consisting of hydrogen, hydroxyl,  
 20 acylamino, alkyl, alkoxy, aryloxy, aryl, aryloxyaryl, carboxyl,  
 carboxylalkyl, carboxyl-substituted alkyl, carboxyl-cycloalkyl, carboxyl-  
 substituted cycloalkyl, carboxylaryl, carboxyl-substituted aryl,  
 carboxylheteroaryl, carboxyl-substituted heteroaryl, carboxylheterocyclic,  
 carboxyl-substituted heterocyclic, cycloalkyl, substituted alkyl, substituted  
 25 alkoxy, substituted aryl, substituted aryloxy, substituted aryloxyaryl,  
 substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic and  
 substituted heterocyclic;

Q is selected from the group consisting of -O-, -S-, -S(O)-, -S(O)<sub>2</sub>,  
 and -NR<sup>4</sup>-;

-- 314 --

R<sup>4</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic or, optionally, R<sup>4</sup> and R<sup>1</sup> or R<sup>4</sup> and R<sup>2</sup>, together with the atoms to which they are bound, are joined to form a heteroaryl, a substituted heteroaryl, a heterocyclic or a substituted heterocyclic group;

W is selected from the group consisting of nitrogen and carbon; and


W' is selected from the group consisting of nitrogen, carbon, oxygen, sulfur, S(O), and S(O)<sub>2</sub>;

X is selected from the group consisting of hydroxyl, alkoxy, substituted alkoxy, alkenoxy, substituted alkenoxy, cycloalkoxy, substituted cycloalkoxy, cycloalkenoxo, substituted cycloalkenoxo, aryloxy, substituted aryloxy, heteroaryloxy, substituted heteroaryloxy, heterocycloxy, substituted heterocycloxy and -NR<sup>n</sup>R<sup>m</sup> where each R<sup>n</sup> is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;

and further wherein the compound of formula Ia and/or Ib has a binding affinity to VLA-4 as expressed by an IC<sub>50</sub> of about 15 μM or less.

52. The pharmaceutical composition of Claim 51, wherein R<sup>3</sup> is a group of the formula:



30



-- 315 --

wherein R<sup>9</sup> and x are as defined in Claim 47; and R<sup>3'</sup> is hydrogen.

53. The pharmaceutical composition of Claim 52, wherein R<sup>9</sup> is in the *para* position of the phenyl ring; and x is an integer from 1 to 4.

5

54. The pharmaceutical composition of Claim 53, wherein R<sup>9</sup> is selected from the group consisting of -O-Z-NR<sup>11</sup>R<sup>11'</sup> and -O-Z-R<sup>12</sup> wherein R<sup>11</sup> and R<sup>11'</sup> are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, and where R<sup>11</sup> and R<sup>11'</sup> are joined to form a heterocycle or a substituted heterocycle, R<sup>12</sup> is selected from the group consisting of heterocycle and substituted heterocycle, and Z is selected from the group consisting of -C(O)- and -SO<sub>2</sub>-.

15

55. The pharmaceutical composition of Claim 54, wherein Z is -C(O)-.

56. The pharmaceutical composition of Claim 55, wherein R<sup>9</sup> is -OC(O)NR<sup>11</sup>R<sup>11'</sup>.

20


57. The pharmaceutical composition of Claim 51, wherein Q is -NR<sup>4</sup>-.

58. The pharmaceutical composition of Claim 51, wherein the compound has formula IIa or IIb:


25

-- 316 --

5



10



15

wherein

ring A and ring B form a heteroaryl or substituted heteroaryl group having two nitrogen atoms in the heteroaryl ring;

20

$R^5$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

25

$R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $-SO_2R^{10}$  where  $R^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl;

-- 317 --


or optionally, one of, R<sup>4</sup> and ring A, R<sup>4</sup> and R<sup>5</sup>, R<sup>4</sup> and R<sup>6</sup>, or R<sup>5</sup> and R<sup>6</sup>, together with the atoms to which they are bound, can be joined to form a heterocyclic or substituted heterocyclic ring;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof; and provided that ring B does not form a 6-amino or substituted amino pyrimidin-4-yl group.

59. The pharmaceutical composition of Claim 58, wherein ring A forms a pyridazine, pyrimidine or pyrazine ring, wherein the pyridazine, pyrimidine or pyrazine ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen, and

ring B forms a pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or a 1,1-dioxo-1,2,5-thiadiazole ring, wherein the pyridazine, pyrimidine or pyrazine ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.


60. The pharmaceutical composition of Claim 51, wherein the compound has formula IIIa, IIIc, IIIe or IIIf:



IIIa


-- 318 --

5




IIIc

10



IIIId


15



IIIe

20

25



IIIf

30

wherein

$R^{4'}$  is selected from the group consisting of hydrogen and alkyl or, optionally, one of,  $R^{4'}$  and  $R^5$ ,  $R^{4'}$  and  $R^6$ ,  $R^5$  and  $R^6$ ,  $R^5$  and  $R^8$ , or  $R^6$  and  $R^8$ , together with the atoms to which they are bound, are joined to form a heterocyclic, a substituted heterocyclic, a heteroaryl or substituted heteroaryl group optionally containing from 1 to 3 additional hetero ring atoms selected from the group consisting of oxygen, nitrogen and sulfur;

$R^{4'}$  is selected from the group consisting of hydrogen and alkyl;

$R^5$  is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

$R^6$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $-SO_2R^{10}$  where  $R^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl;

$R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$R^{16}$  and  $R^{17}$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen; and

-- 320 --

$R^{18}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

5  $R^{20}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

10  $R^{21}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;


$b$  is 1 or 2;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

15

61. The pharmaceutical composition of Claim 60, wherein the compound is selected from formula IIIId, IIIe or IIIf.

20 62. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of formula IVa:



wherein R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form a heteroaryl group having two nitrogen atoms;

and further wherein said heteroaryl group is optionally substituted, on  
5 any ring atom capable of substitution, with 1-3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy,  
10 substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen, hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl,  
15 substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino, -OS(O)<sub>2</sub>-alkyl, -OS(O)<sub>2</sub>-substituted alkyl, -OS(O)<sub>2</sub>-aryl, -OS(O)<sub>2</sub>-substituted aryl, -OS(O)<sub>2</sub>-heteroaryl, -OS(O)<sub>2</sub>-substituted  
20 heteroaryl, -OS(O)<sub>2</sub>-heterocyclic, -OS(O)<sub>2</sub>-substituted heterocyclic, -OSO<sub>2</sub>-NRR where each R is independently hydrogen or alkyl, -NRS(O)<sub>2</sub>-alkyl, -NRS(O)<sub>2</sub>-substituted alkyl, -NRS(O)<sub>2</sub>-aryl, -NRS(O)<sub>2</sub>-substituted aryl, -NRS(O)<sub>2</sub>-heteroaryl, -NRS(O)<sub>2</sub>-substituted heteroaryl, -NRS(O)<sub>2</sub>-heterocyclic, -NRS(O)<sub>2</sub>-substituted heterocyclic, -NRS(O)<sub>2</sub>-NR-alkyl,  
25 -NRS(O)<sub>2</sub>-NR-substituted alkyl, -NRS(O)<sub>2</sub>-NR-aryl, -NRS(O)<sub>2</sub>-NR-substituted aryl, -NRS(O)<sub>2</sub>-NR-heteroaryl, -NRS(O)<sub>2</sub>-NR-substituted heteroaryl, -NRS(O)<sub>2</sub>-NR-heterocyclic, -NRS(O)<sub>2</sub>-NR-substituted heterocyclic where R is hydrogen or alkyl, -N[S(O)<sub>2</sub>-R']<sub>2</sub> and -N[S(O)<sub>2</sub>-NR']<sub>2</sub> where each R' is independently selected from the group consisting of alkyl, substituted alkyl,

aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

5  $R^{13}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl, Cy, and Cy- $C_{1-10}$  alkyl, wherein alkyl is optionally substituted with one to four substituents independently selected from  $R^a$ ; and Cy is optionally substituted with one to four substituents independently selected from  $R^b$ ;

10  $R^{14}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, Cy, Cy- $C_{1-10}$  alkyl, Cy- $C_{2-10}$  alkenyl and Cy- $C_{2-10}$  alkynyl, wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents selected from phenyl and  $R^x$ , and Cy is optionally substituted with one to four substituents independently selected from  $R^y$ ;

or  $R^{13}$ ,  $R^{14}$  and the atoms to which they are attached together form a mono- or bicyclic ring containing 0-2 additional heteratoms selected from N, O and S;

15  $R^{15}$  is selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl, aryl- $C_{1-10}$  alkyl, heteroaryl, heteroaryl- $C_{1-10}$  alkyl, wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents selected from  $R^x$ , and aryl and heteroaryl are optionally substituted with one to four substituents independently selected from  $R^y$ ;

20 or  $R^{14}$ ,  $R^{15}$  and the carbon to which they are attached form a 3-7 membered mono- or bicyclic ring containing 0-2 heteroatoms selected from N, O and S;

25  $R^a$  is selected from the group consisting of Cy and a group selected from  $R^x$ , wherein Cy is optionally substituted with one to four substituents independently selected from  $R^c$ ;

$R^b$  is selected from the group consisting of  $R^a$ ,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl  $C_{1-10}$  alkyl, heteroaryl  $C_{1-10}$  alkyl, wherein alkyl, alkenyl, alkynyl, aryl, heteroaryl are optionally substituted with a group independently selected from  $R^c$ ;

30  $R^c$  is selected from the group consisting of halogen,  $NO_2$ ,  $C(O)OR^f$ ,



-- 323 --

$C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy, aryl, aryl  $C_{1-4}$  alkyl, aryloxy, heteroaryl,  $NR^fR^g$ ,  $R^fC(O)R^g$ ,  $NR^fC(O)NR^fR^g$ , and CN;

$R^d$  and  $R^e$  are independently selected from hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, Cy and Cy  $C_{1-10}$  alkyl, wherein alkyl, alkenyl, alkynyl and Cy are optionally substituted with one to four substituents independently selected from  $R^c$ ;

or  $R^d$  and  $R^e$  together with the atoms to which they are attached form a heterocyclic ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen;

$R^f$  and  $R^g$  are independently selected from hydrogen,  $C_{1-10}$  alkyl, Cy and Cy- $C_{1-10}$  alkyl wherein Cy is optionally substituted with  $C_{1-10}$  alkyl; or  $R^f$  and  $R^g$  together with the carbon to which they are attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

$R^h$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, cyano, aryl, aryl  $C_{1-10}$  alkyl, heteroaryl, heteroaryl  $C_{1-10}$  alkyl, and  $-SO_2R^i$ ; wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from  $R^a$ ; and aryl and heteroaryl are each optionally substituted with one to four substituents independently selected from  $R^b$ ;

$R^i$  is selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, and aryl; wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from  $R^c$ ;

$R^x$  is selected from the group consisting of  $-OR^d$ ,  $-NO_2$ , halogen,  $-S(O)_mR^d$ ,  $-SR^d$ ,  $-S(O)_2OR^d$ ,  $-S(O)_mNR^dR^e$ ,  $-NR^dR^e$ ,  $-O(CR^fR^g)_nNR^dR^e$ ,  $-C(O)R^d$ ,  $-CO_2R^d$ ,  $-CO_2(CR^fR^g)_nCONR^dR^e$ ,  $-OC(O)R^d$ ,  $-CN$ ,  $-C(O)NR^dR^e$ ,  $-NR^dC(O)R^e$ ,  $-OC(O)NR^dR^e$ ,  $-NR^dC(O)OR^e$ ,  $-NR^dC(O)NR^dR^e$ ,  $-CR^d(N-OR^e)$ ,  $CF_3$ , oxo,  $NR^dC(O)NR^dSO_2R^i$ ,  $NR^dS(O)_mR^e$ ,  $-OS(O)_2OR^d$ , and  $-OP(O)(OR^d)_2$ ;

-- 324 --

$R^y$  is selected from the group consisting of  $R^x$ ,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl  $C_{1-10}$ alkyl, heteroaryl  $C_{1-10}$  alkyl, cycloalkyl, heterocyclyl; wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from  $R^x$ ;

5        Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;

$m$  is an integer from 1 to 2;

$n$  is an integer from 1 to 10;

W is selected from the group consisting of carbon and nitrogen;

W' is selected from the group consisting of carbon, nitrogen, oxygen,  
10        sulfur, S(O) and S(O)<sub>2</sub>;

X' is selected from the group consisting of -C(O)OR<sup>d</sup>,  
-P(O)(OR<sup>d</sup>)(OR<sup>e</sup>), -P(O)(R<sup>d</sup>)(OR<sup>e</sup>), -S(O)<sub>m</sub>OR<sup>d</sup>, -C(O)NR<sup>d</sup>R<sup>h</sup>, and -5-  
tetrazolyl;

and enantiomers, diastereomers and pharmaceutically acceptable salts  
15        thereof;

and further wherein the compound of formula IV has a binding  
affinity to VLA-4 as expressed by an IC<sub>50</sub> of about 15 μM or less;

and provided that when R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and  
W to which they are bound respectively, are joined to form a 2-  
20        arylpyrimidin-4-yl group and R<sup>14</sup> is hydrogen, then R<sup>15</sup> is not alkyl of from 1  
to 6 carbon atoms optionally substituted with hydroxyl; and when R<sup>1</sup> and R<sup>2</sup>,  
together with the carbon atom and W to which they are bound respectively,  
are joined to form a 5-arylpyrazin-2-yl group and R<sup>14</sup> is hydrogen, then R<sup>15</sup> is  
not 4-hydroxybenzyl.

25



63.        The pharmaceutical composition of Claim 62, wherein R<sup>1</sup> and  
R<sup>2</sup>, together with the carbon atom and W to which they are bound  
respectively, are joined to form a pyridazine, pyrimidine, pyrazine, 1-oxo-  
1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring; wherein the pyridazine,  
30        pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or a 1,1-dioxo-1,2,5-

-- 325 --

thiadiazole ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.


64. The pharmaceutical composition of Claim 62, wherein X' is -C(O)OR<sup>d</sup>.

65. The pharmaceutical composition of Claim 62, wherein the compound has formula Va, Vc, Vd, Ve or Vf:




-- 326 --

5



Vd


10



Ve

15

20



Vf

wherein

25

R<sup>5</sup> is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, heteroaryl and substituted heteroaryl;

30

R<sup>6</sup> is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted

cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and  $-\text{SO}_2\text{R}^{10}$  where  $\text{R}^{10}$  is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl; and

$\text{R}^7$  and  $\text{R}^8$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$\text{R}^{16}$  and  $\text{R}^{17}$  are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen; and

$\text{R}^{18}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

$\text{R}^{20}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;


$\text{R}^{21}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;

$b$  is 1 or 2;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

66. The pharmaceutical composition of Claim 65, wherein the compound is selected from formula Vd, Ve or Vf.

67. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of formula VIa and/or VIb:



15 wherein, in formula VIa, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W to which they are bound respectively, are joined to form an aryl, cycloalkenyl, heteroaryl or heterocyclic group having at least five atoms in the aryl, cycloalkenyl, heteroaryl or heterocyclic group and optionally containing or additionally containing in the case of heteroaryl and

20 heterocyclic groups 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heteroaryl or heterocyclic group is mono-cyclic;

25 in formula VIb, R<sup>1</sup> and R<sup>2</sup>, together with the carbon atom and W' to which they are bound respectively, are joined to form a cycloalkyl, cycloalkenyl or heterocyclic group having at least five atoms in the cycloalkyl, cycloalkenyl or heterocyclic group and optionally containing or additionally containing in the case of the heterocyclic group 1 to 3 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, and wherein the heterocyclic group is mono-cyclic;

and further wherein said aryl, cycloalkyl, cycloalkenyl, heteroaryl or heterocyclic group of formula VIa or VIb is optionally substituted, on any ring atom capable of substitution, with 1-3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, acyl, acylamino, thiocarbonylamino, acyloxy, amino, substituted amino, amidino, alkyl amidino, thioamidino, aminoacyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aryl, substituted aryl, aryloxy, substituted aryloxy, aryloxyaryl, substituted aryloxyaryl, cyano, halogen, hydroxyl, nitro, oxo, carboxyl, cycloalkyl, substituted cycloalkyl, guanidino, guanidinosulfone, thiol, thioalkyl, substituted thioalkyl, thioaryl, substituted thioaryl, thiocycloalkyl, substituted thiocycloalkyl, thioheteroaryl, substituted thioheteroaryl, thioheterocyclic, substituted thioheterocyclic, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, cycloalkoxy, substituted cycloalkoxy, heteroaryloxy, substituted heteroaryloxy, heterocyclyloxy, substituted heterocyclyloxy, oxycarbonylamino, oxythiocarbonylamino,  $-\text{OS}(\text{O})_2\text{-alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{OS}(\text{O})_2\text{-aryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{OS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{OS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{OS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{OSO}_2\text{-NRR}$  where each R is independently hydrogen or alkyl,  $-\text{NRS}(\text{O})_2\text{-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-substituted heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted alkyl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted aryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heteroaryl}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-heterocyclic}$ ,  $-\text{NRS}(\text{O})_2\text{-NR-substituted heterocyclic}$  where R is hydrogen or alkyl,  $-\text{N}[\text{S}(\text{O})_2\text{-R'}]_2$  and  $-\text{N}[\text{S}(\text{O})_2\text{-NR'}]_2$  where each R' is independently selected from the group consisting of alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

$R^{23}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl optionally substituted with one to four substituents independently selected from  $R^a$  and Cy optionally substituted with one to four substituents independently selected from  $R^b$ ;

- 5  $R^{24}$  is selected from the group consisting of  $Ar^1-Ar^2-C_{1-10}$  alkyl,  $Ar^1-Ar^2-C_{2-10}$  alkenyl,  $Ar^1-Ar^2-C_{2-10}$  alkynyl, wherein  $Ar^1$  and  $Ar^2$  are independently aryl or heteroaryl each of which is optionally substituted with one to four substituents independently selected from  $R^b$ ; alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents independently  
10 selected from  $R^a$ ;

- $R^{25}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl, aryl  $C_{1-10}$ alkyl, heteroaryl, and heteroaryl  $C_{1-10}$  alkyl, wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents selected from  $R^a$ , and aryl and heteroaryl are  
15 optionally substituted with one to four substituents independently selected from  $R^b$ ;

- $R^a$  is selected from the group consisting of Cy,  $-OR^d$ ,  $-NO_2$ , halogen  $-S(O)_mR^d$ ,  $-SR^d$ ,  $-S(O)_2OR^d$ ,  $-S(O)_mNR^dR^e$ ,  $-NR^dR^e$ ,  $-O(CR^fR^g)_nNR^dR^e$ ,  $-C(O)R^d$ ,  $-CO_2R^d$ ,  $-CO_2(CR^fR^g)_nCONR^dR^e$ ,  $-OC(O)R^d$ ,  $-CN$ ,  
20  $-C(O)NR^dR^e$ ,  $-NR^dC(O)R^e$ ,  $-OC(O)NR^dR^e$ ,  $-NR^dC(O)OR^e$ ,  $-Nr^dC(O)NR^dR^e$ ,  $-CR^d(N-OR^e)$ ,  $CF_3$ , and  $-OCF_3$ ;

wherein Cy is optionally substituted with one to four substituents independently selected from  $R^e$ ;

- $R^b$  is selected from the group consisting of  $R^a$ ,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, aryl  $C_{1-10}$  alkyl, heteroaryl  $C_{1-10}$ alkyl,  
25 wherein alkyl, alkenyl, aryl, heteroaryl are optionally substituted with a group independently selected from  $R^e$ ;

$R^e$  is selected from the group consisting of halogen, amino, carboxy,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy, aryl, aryl  $C_{1-4}$ alkyl, hydroxy,  $CF_3$ , and aryloxy;



-- 331 --

$R^{d'}$  and  $R^{e'}$  are independently selected from hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, Cy and Cy  $C_{1-10}$ alkyl, wherein alkyl, alkenyl, alkynyl and Cy are optionally substituted with one to four substituents independently selected from  $R^{e'}$ ; or  $R^{d'}$  and  $R^{e'}$  together with the atoms to which they are  
5 attached form a heterocyclic ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen;

$R^{f'}$  and  $R^{g'}$  are independently selected from hydrogen,  $C_{1-10}$  alkyl, Cy and Cy- $C_{1-10}$  alkyl; or  $R^{f'}$  and  $R^{g'}$  together with the carbon to which they are attached form a ring of 5 to 7 members containing 0-2 heteroatoms  
10 independently selected from oxygen, sulfur and nitrogen;

$R^{h'}$  is selected from the group consisting of hydrogen,  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, cyano, aryl, aryl  $C_{1-10}$  alkyl, heteroaryl, heteroaryl  $C_{1-10}$  alkyl, or  $-SO_2R^{i'}$ ;

wherein alkyl, alkenyl, and alkynyl are optionally substituted with one  
15 to four substituents independently selected from  $R^{a'}$ ; and aryl and heteroaryl are each optionally substituted with one to four substituents independently selected from  $R^{b'}$ ;

$R^{i'}$  is selected from the group consisting of  $C_{1-10}$  alkyl,  $C_{2-10}$  alkenyl,  $C_{2-10}$  alkynyl, and aryl;

20 wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from  $R^{c'}$ ;

Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;

$X''$  is selected from the group consisting of  $-C(O)OR^{d'}$ ,  $-P(O)(OR^{d'})(OR^{e'})$ ,  $-P(O)(R^{d'})(OR^{e'})$ ,  $-S(O)_mOR^{d'}$ ,  $-C(O)NR^{d'}R^{h'}$ , and -5-  
25 tetrazolyl;

$m$  is an integer from 1 to 2;

$n$  is an integer from 1 to 10;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;

-- 332 --

and further wherein the compounds of formula VIa and/or VIb have a binding affinity to VLA-4 as expressed by an  $IC_{50}$  of about 15  $\mu$ M or less.

68. The pharmaceutical composition of Claim 67, wherein  $R^1$  and  $R^2$ , together with the carbon atom and W to which they are bound respectively, are joined to form a heteroaryl or substituted heteroaryl group having two nitrogen atoms in the heteroaryl ring.

69. The pharmaceutical composition of Claim 68, wherein  $R^1$  and  $R^2$ , together with the carbon atom and W to which they are bound respectively, are joined to form a pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring; wherein the pyridazine, pyrimidine, pyrazine, 1-oxo-1,2,5-thiadiazole or 1,1-dioxo-1,2,5-thiadiazole ring is optionally substituted with 1 to 3 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen.


70. The pharmaceutical composition of Claim 67, wherein  $X''$  is  $-C(O)OR^d$ .

71. The pharmaceutical composition of Claim 67, wherein  $R^{24}$  is  $-CH_2-Ar^2-Ar^1$  and  $R^{25}$  is hydrogen.

72. The pharmaceutical composition of Claim 67, wherein the compound has formula VIIa, VIIc, VIId, VIIe or VIIf:


-- 333 --

5




VIIa

10



VIIc


15



VIIId

20

25




VIIe

30

-- 334 --

5



VIII f

wherein

$R^5$  is selected from the group consisting of alkyl, substituted alkyl,  
 10 alkenyl, substituted alkenyl, aryl, substituted aryl, cycloalkyl, substituted  
 cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted  
 heterocyclic, heteroaryl and substituted heteroaryl;

$R^6$  is selected from the group consisting of hydrogen, alkyl,  
 substituted alkyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted  
 15 cycloalkenyl, heterocyclic, substituted heterocyclic, aryl, substituted aryl,  
 heteroaryl, substituted heteroaryl, and  $-SO_2R^{10}$  where  $R^{10}$  is selected from the  
 group consisting of alkyl, substituted alkyl, cycloalkyl, substituted  
 cycloalkyl, cycloalkenyl, substituted cycloalkenyl, heterocyclic, substituted  
 heterocyclic, aryl, substituted aryl, heteroaryl, substituted heteroaryl; and

20  $R^7$  and  $R^8$  are independently selected from the group consisting of  
 hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl,  
 substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted  
 heterocyclic and halogen;

$R^{16}$  and  $R^{17}$  are independently selected from the group consisting of  
 25 hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino,  
 substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl,  
 heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and  
 halogen; and

$R^{18}$  is selected from the group consisting of alkyl, substituted alkyl,  
 30 alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted

-- 335 --

cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic;

5  $R^{20}$  is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic and halogen;

$R^{21}$  is selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, amino, substituted amino, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heterocyclic and substituted heterocyclic;

10  $b$  is 1 or 2;

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof.

73. The pharmaceutical composition of Claim 72, wherein the  
15 compound is selected from formula VIId, VIIe or VIIf.

74. A method for binding VLA-4 in a biological sample which  
method comprises contacting the biological sample with a compound of  
Claims 27, 38, 43 or 50 under conditions wherein said compound binds to  
20 VLA-4.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT 00/01686

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/42 A61K31/505 A61P7/00 C07D401/12 A61K31/495  
C07D405/12 C07D409/12 C07D285/10 A61K31/41 C07D417/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, Y	WO 99 37618 A (ARCHIBALD SARAH CATHERINE ;CELLTECH THERAPEUTICS LTD (GB); WARRELL) 29 July 1999 (1999-07-29) claim 1	50
P, Y	WO 99 37605 A (NOVARTIS ERFINDUNGEN VERWALTUN ;NOVARTIS AG (CH); VON MATT PETER J) 29 July 1999 (1999-07-29) claim 1; example 1	50
P, Y	WO 99 10313 A (HOFFMANN LA ROCHE) 4 March 1999 (1999-03-04) claim 1	50
P, Y	WO 99 10312 A (HOFFMANN LA ROCHE) 4 March 1999 (1999-03-04) claim 1	50
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

29 May 2000

Date of mailing of the international search report

09.06.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Gettins, M

## INTERNATIONAL SEARCH REPORT

International Application No

T/US 00/01686

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	WO 99 06390 A (LOMBARDO LOUIS JOHN ; SEMKO CHRISTOPHER M (US); THORSETT EUGENE D () 11 February 1999 (1999-02-11) claims 1,16 ---	50
P,Y	WO 99 06391 A (ATHENA NEUROSCIENCES INC) 11 February 1999 (1999-02-11) claims 4,6,13 ---	50
A	EP 0 288 176 A (AJINOMOTO KK) 26 October 1988 (1988-10-26) claim 1 ---	50
A	DE 196 54 483 A (MERCK PATENT GMBH) 2 January 1998 (1998-01-02) claim 1 ---	50
A	DE 195 48 709 A (MERCK PATENT GMBH) 3 July 1997 (1997-07-03) claim 1 -----	50

# INTERNATIONAL SEARCH REPORT

International application No.  
/US 00/01686

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☒ Claims Nos.: 1-49, 51-74  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
  
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
  
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-49,51-74

Present claims 1-10, 13-16, 19-23, 26-35, 38-40, 43-47, 51-59, 62-64, 67-71 relate to an extremely large number of possible compounds. In fact, the claims contain so many options, variables, possible permutations and provisos that a lack of clarity (and/or conciseness) within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search of the claims impossible. In particular the proviso relating to the binding affinity to VLA-4 is a functional definition which can only be established by empirical means. Combining this with the fact that the scope of the claims is huge (see in particular the use of unlimited terms such as "substituted" which is not further defined) it is impossible for the skilled person to readily determine which compounds fall under the scope of formulae Ia and Ib. Consequently, the search has been carried out for those parts of the application which do appear to be clear (and/or concise), namely compound claim 50 which does not refer to the said proviso. It is noted that all of the compounds in claim 50 appear to be phenylalanine derivatives and that the vast majority of the compounds are 4-(N,N-dimethylcarbamoyloxy)-phenylalanine derivatives (the only exceptions are phenylalanine derivatives). Since a search limited to phenylalanine derivatives would represent an overgeneralisation of the examples the search has been limited to the compounds of claim 50 containing the 4-(N,N-dimethylcarbamoyloxy)-phenylalanine group (in other words where X is OH or alkoxy, Q is nitrogen and R3 is 4-(N,N-dimethylcarbamoyloxy)-phenyl). Since no claim has been limited thereto it was not possible to limit the search other than on the basis of claim 50.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC 00/01686

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9937618 A	29-07-1999	AU 2432099 A	09-08-1999
WO 9937605 A	29-07-1999	AU 2829299 A	09-08-1999
WO 9910313 A	04-03-1999	AU 9341998 A	16-03-1999
WO 9910312 A	04-03-1999	AU 9262098 A	16-03-1999
		NO 20000841 A	21-02-2000
WO 9906390 A	11-02-1999	AU 8584998 A	22-02-1999
		AU 8605098 A	22-02-1999
		EP 1000051 A	17-05-2000
		EP 0954519 A	10-11-1999
		NO 20000413 A	28-03-2000
		WO 9906391 A	11-02-1999
WO 9906391 A	11-02-1999	AU 8584998 A	22-02-1999
		AU 8605098 A	22-02-1999
		EP 1000051 A	17-05-2000
		EP 0954519 A	10-11-1999
		NO 20000413 A	28-03-2000
		WO 9906390 A	11-02-1999
EP 0288176 A	26-10-1988	JP 2049974 C	10-05-1996
		JP 7084424 B	13-09-1995
		JP 63258841 A	26-10-1988
DE 19654483 A	02-01-1998	AU 3343097 A	21-01-1998
		CA 2259224 A	08-01-1998
		CZ 9804249 A	17-03-1999
		WO 9800395 A	08-01-1998
		EP 0907637 A	14-04-1999
		NO 986090 A	23-12-1998
		PL 330915 A	07-06-1999
		SK 176898 A	07-05-1999
		BR 9709953 A	10-08-1999
DE 19548709 A	03-07-1997	AU 1301697 A	17-07-1997
		BR 9612201 A	13-07-1999
		CA 2241149 A	03-07-1997
		CZ 9801951 A	11-11-1998
		WO 9723451 A	03-07-1997
		EP 0879227 A	25-11-1998
		JP 2000502664 T	07-03-2000
		NO 982907 A	22-06-1998
		PL 327185 A	23-11-1998
		SK 78398 A	04-11-1998